arXiv:1211.6073v2 [cond-mat.str-el] 27 Nov 2012

Elementary objects of the 1D Hubbard model from its symmetry and exact solution II

J. M. P. Carmelo[a,b,c] and P. D. Sacramento[c,d]
[a]*Center of Physics, University of Minho, Campus Gualtar, P-4710-057 Braga, Portugal*
[b]*Institut für Theoretische Physik III, Universität Stuttgart, D-70550 Stuttgart, Germany*
[c]*Beijing Computational Science Research Center, Beijing 100084, China and*
[d]*CFIF, Instituto Superior Técnico, TU Lisbon, P-1049-001 Lisboa, Portugal*
(Dated: 27 November 2012)

This paper presents the second part of a study on the elementary objects of the 1D Hubbard model that emerge from the interplay of its global symmetry with the exact Bethe-ansatz (BA) solution. Here we introduce and study the corresponding scattering theory and its scatterers and scattering centers that naturally emerge from the present formulation elementary objects. The theory refers to the excited states of ground states with arbitrary values of the densities and finite repulsive interaction. The occurrence of only zero-momentum forward-scattering results from the infinite number of conservation laws associated with the model integrability in the BA solution subspace. The unbound spinons and unbound η-spinons whose occupancies generate the energy eigenstates outside the BA solution subspace are found to be neither scatterers nor scattering centers. Those are rather the c pseudofermions and composite spin-neutral 2ν-spinon pseudofermions and η-spin-neutral 2ν-η-spinon pseudofermions. Here $\nu = 1, 2, ...$ is the number of spinon or η-spinon pairs. Their discrete momentum values are slightly shifted relative to those of the c fermions, $s\nu$ fermions, and $\eta\nu$ fermions introduced in the first paper. Otherwise, they have exactly the same properties. Each ground-state - excited-state transition is associated with a well defined set of elementary zero-momentum forward-scattering events. The pseudofermion scatterers dressed S matrix is expressed as a *commutative* product of S matrices, each corresponding to an elementary two-pseudofermion scattering event. This commutative factorization is stronger than the factorization associated with Yang-Baxter equation for the original spin-1/2 electron bare S matrix. The momentum dependence of the exponents characterizing the finite-energy correlation function singularities of the metallic phases of a wide class of 1D integrable and non-integrable systems is in the present exactly solvable model found to be fully controlled by the phase shifts and corresponding dressed S matrix introduced in this paper. Furthermore, we find that the elementary spin and charge currents carried by the present formulation elementary objects are controlled by the dressed phase shifts. The relation of the elementary objects considered here and in the first paper to the traditional holon and spinon descriptions is clarified. Our study confirms that there is no contradiction whatsoever between those and the present formulation. The latter is normal ordered relative to the electron and rotated-electron vacuum, whereas the traditional holon and spinon descriptions are normal ordered relative to an initial ground state. They refer to a different basis of scattering states and thus a different choice of scatterers and scattering centers. The possibility of such distinct basis choices stems from the degeneracy of the excited energy eigenstates that span the subspaces of the corresponding scattering theories. The consequences of the elementary objects choice for the description of the microscopic processes that control the model dynamical and spectral properties is also investigated.

PACS numbers: 03.65.Ca, 71.10.Pm, 71.27.+a, 03.65.Nk

I. INTRODUCTION

Since the 1D Hubbard model Bethe-ansatz (BA) solution is exact [1–4], most results of studies by means of that solution are correct independently of the choice of elementary objects whose occupancy configurations generate the model excitations. However, the operational definition of these elementary objects in terms of the original electron operators and how do they scatter with each other are two issues of highest importance for the physical interpretation of such results. The studies of the first paper, Ref. [5], focus on the former issue. In this paper we handle the second issue, which concerns the properties of the scatterers and scattering centers that emerge from the c fermions, spinons, and η-spinons of the operator formulation introduced in the first paper. We find that it leads to a uniquely defined choice of scattering states basis. Such states are found to be in one-to-one correspondence to the excited energy eigenstates of ground states with arbitrary values of the electronic and spin densities and finite repulsive interaction.

Our results reveal that due to the degeneracy of the energy spectrum of excited energy eigenstates of ground states with zero spin density and/or electronic density one (half filling) there are for the subspaces spanned by such states distinct choices of scattering states. Furthermore, as further discussed below we show that the choices of scattering states other than that of the pseudofermion scattering theory introduced in this paper refer to the scattering theories previously introduced in the literature. The validity of such theories is though limited to the above subspaces for which there are degeneracies that allow the use of alternative choices of scattering states and thus of scatterers and scattering centers. However, the major advantage of the scattering theory introduced in this paper is found to refer to the explicit description of the microscopic processes that control the model dynamical and spectral properties. Such an advantage follows form that theory accounting for both the phase shifts of the scatterers that pre-exist in the initial ground state and those that are created under the transitions from it to the excited states. Other scattering theories account only for the latter phase shifts.

Indeed, recently there has been a renewed interest on the dynamical and spectral properties of 1D correlated systems, particularly for finite energy, where methods beyond linear Luttinger-liquid theory [6] are now available. This includes the pseudofermion dynamical theory (PDT) [7–9] for the 1D Hubbard model, which is a generalization to arbitrary values of the repulsive interaction of the large-interaction method of Ref. [10], as well as alternative methods valid for both integrable and non-integrable 1D correlated problems [11–18]. (While above *finite energy* means an energy larger than the typical linear Luttinger-liquid theory excitation energy, in this paper finite energy often refers to both low energy and high energy in the sense that there are no restrictions to its values, unlike in that theory.)

One of our goals is the description of the scattering mechanisms behind the exotic dynamical and spectral properties studied within the PDT. Such a description is much simpler in terms of the scattering of the scatterers and scattering centers that emerge naturally from the elementary objects of the present formulation than in terms of electronic scattering. Indeed, the occupancy configurations that generate the energy eigenstates are much simpler in terms of such elementary objects than of the original electrons. The simplicity of the scattering events studied in this paper directly profits from the occurrence of an infinite number of conservation laws [19–21] associated with the 1D Hubbard model integrability [1–4]. Consistent with the elementary objects of the present formulation emerging from the interplay of the model global symmetry with its exact BA solution, such conservation laws are explicit in the scattering theory introduced in this paper, in that its scatterers and scattering centers only undergo zero-momentum forward scattering.

The studies of this paper confirm that such scattering events play a major role in the model dynamical and spectral properties. Specifically, the corresponding dressed S matrix phase shifts determine and control the momentum, interaction, and density dependence of the correlation-function exponents associated with the finite-energy behavior described in the metallic phase by the PDT [7–9]. Our present study reveals that indeed the microscopic scattering processes behind the corresponding finite-energy spectral properties are simple in terms of the elementary objects of the present operational formulation. Moreover, the elementary spin and charge currents are found to be controlled by the scattering theory phase shifts. The exotic finite-energy behavior first found for the 1D Hubbard model by the PDT is similar to that predicted by the studies of Refs. [11–13]. Those use a method valid for a wider class of both integrable and non-integrable 1D correlated systems. From the relation of the PDT to the scattering theory introduced in the following, one confirms that the corresponding exotic scattering centers of the 1D Hubbard model are observed by angle-resolved photoelectron spectroscopy in quasi-1D metals [22–24]. The model describes as well the effects of correlations in semiconductor - metal transitions of doped quasi-1D materials [25] and is of interest for systems of ultra-cold atoms on 1D optical lattices [26].

The scattering theory introduced in this paper goes beyond the BA solution, in that it accounts for both excited states inside and outside that solution subspace, which is spanned by the Bethe states. The latter energy eigenstates can be either lowest-weight states (LWSs) or highest-weight states (HWSs) of the η-spin and spin $SU(2)$ algebras [27]. The η-spin (and spin) and η-spin projection (and spin projection) of the energy eigenstates are denoted by S_η and $S_\eta^{x_3}$ (and S_s and $S_s^{x_3}$), respectively. The S_α and $S_\alpha^{x_3}$ values of the LWSs and HWSs are such that $S_\alpha = -S_\alpha^{x_3}$ and $S_\alpha = S_\alpha^{x_3}$, respectively. Here $\alpha = \eta$ for η-spin and $\alpha = s$ for spin. We use the BA solution LWS representation, alike in the first paper.

The two global $SU(2)$ symmetries of the Hubbard model on a bipartite lattice, including the present 1D lattice, have been known for a long time [28, 29]. For about twenty years [30] it was believed that for finite on-site interaction values the model global symmetry was $SO(4) = [SU(2) \otimes SU(2)]/Z_2$. The studies of Ref. [31] reveal that the global symmetry of the Hubbard model on a bipartite lattice is larger: It is given by $[SO(4) \otimes U(1)]/Z_2 = SO(3) \otimes SO(3) \otimes U(1)$. As discussed in the first paper, the occurrence of a global c hidden $U(1)$ symmetry beyond $SO(4)$ must be accounted for in studies of the Hubbard model on any bipartite lattice, such as the 1D lattice. That global symmetry may be rewritten as $[SU(2) \times SU(2) \times U(1)]/Z_2^2$. Indeed, it stems from the local gauge $SU(2) \times SU(2) \times U(1)$ symmetry of the Hubbard model on a bipartite lattice with vanishing transfer integral, $t = 0$ [32]. For finite U and t values the latter local symmetry becomes a group of permissible unitary transformations. The corresponding local $U(1)$ canonical transformation is not the ordinary gauge $U(1)$ subgroup of electromagnetism. It is rather a “nonlinear” transformation [32].

It is confirmed in the first paper that the BA solution accounts for the quantum number occupancy configurations that generate the representations of the c hidden $U(1)$ symmetry algebra in $[SU(2) \otimes SU(2) \otimes U(1)]/Z_2^2 = [SO(4) \otimes U(1)]/Z_2$ beyond $SO(4)$. Furthermore, the energy eigenstates outside the BA solution subspace are found to have exactly the same c hidden $U(1)$ symmetry algebra representations quantum number occupancy configurations as the

Bethe states from which they are generated by the off-diagonal η-spin and spin operator algebras. This explains why there is no contradiction whatsoever between the global $[SU(2)\otimes SU(2)\otimes U(1)]/Z_2^2 = [SO(4)\otimes U(1)]/Z_2$ symmetry found in Ref. [31] for the Hubbard model on any bipartite lattice and the results of Ref. [27], concerning the counting of the 4^{N_a} energy eigenstates of the Hubbard model in the bipartite 1D lattice. The studies of that reference have not explicitly considered the model c hidden $U(1)$ symmetry algebra beyond $SO(4)$, yet they have used the BA solution, which includes the quantum number occupancy configurations found in the first paper to generate the representations of the c hidden $U(1)$ symmetry algebra.

The studies of the first paper, Ref. [5], have profited from the relation of the 1D Hubbard model global symmetry to its exact BA solution. The elementary objects considered in it emerge from an operator formulation that makes explicit such a global symmetry. The first step of their emergence is an electron - rotated-electron unitary transformation uniquely defined in that paper, which is directly related to such a symmetry. The generator of the global c hidden $U(1)$ symmetry beyond $SO(4)$ is the number of rotated-electron singly occupied sites operator,

$$2\tilde{S}_c \equiv \hat{V}^{\dagger}\,\hat{Q}\,\hat{V}\,. \tag{1}$$

Here $\hat{Q}$ and the related operator $\hat{D}$ count the number of electron singly-occupied sites and doubly-occupied sites and read,

$$\hat{Q} = \sum_{j=1}^{N_a}\sum_{\sigma=\uparrow,\downarrow}\hat{n}_{j,\sigma}\,(1-\hat{n}_{j,-\sigma})\;; \qquad \hat{D} = (\hat{N}-\hat{Q})/2\,, \tag{2}$$

respectively. Moreover, $\hat{V}$ is the electron - rotated-electron unitary operator, which is uniquely defined in the first paper within the complete basis generated by the model 4^{N_a} energy and momentum eigenstates. The eigenvalues $2S_c = 0, 1, ...$ of the operator provided in Eq. (1) refer to the number of rotated-electron singly occupied sites [31]. The interplay of the BA solution with the model global symmetry was shown in the first paper to be consistent with the related object representation preliminarily considered in Refs. [33, 34]. Furthermore, the objects of the general formulation introduced in the first paper are an extension to the whole Hilbert space of those of Refs. [35, 36] for the low-energy subspace.

The transformation laws of the former objects under the electron - rotated-electron unitary transformation play an important role in the identification of the scatterers and scattering centers of the theory introduced here. The latter are found in this paper to be the c pseudofermions and $\alpha\nu$ pseudofermions. Here $\alpha = \eta, s$ for η-spin and spin, respectively, and $\nu = 1, ..., \infty$. Except for a slight shift of their discrete momentum values, such objects have exactly the same properties as the corresponding c fermions and $\alpha\nu$ fermions, respectively, from which they are generated. The studies of the first paper have revealed that the $\eta\nu$ fermions (and $s\nu$ fermions) are η-spin-neutral (and spin-neutral) composite objects containing $\nu = 1, ..., \infty$ pairs of η-spin-1/2 anti-bound η-spinons with opposite η-spin projection (and of spin-1/2 bound spinons with opposite spin projection). An additional well-defined number of η-spinons (and spinons) remains invariant under the electron - rotated-electron unitary transformation defined in that paper. Since they remain unbound, they are called unbound $\pm 1/2$ η-spinons (and unbound $\pm 1/2$ spinons). The values of the numbers $M^{un}_{\eta,\pm 1/2}$ of unbound $\pm 1/2$ η-spinons and $M^{un}_{s,\pm 1/2}$ of unbound $\pm 1/2$ spinons are fully controlled by the η-spin S_η and η-spin projection $S_\eta^{x_3}$ and spin S_s and spin projection $S_s^{x_3}$, respectively, of the subspace or state under consideration as follows,

$$M^{un}_{\alpha} = [M^{un}_{\alpha,-1/2} + M^{un}_{\alpha,+1/2}] = 2S_\alpha\,; \quad M^{un}_{\alpha,\pm 1/2} = [S_\alpha \mp S_\alpha^{x_3}]\,; \quad \alpha = \eta\,,s\,. \tag{3}$$

Thus the η-spin S_η, η-spin projection $S_\eta^{x_3}$, spin S_s, and spin projection $S_s^{x_3}$ of an energy eigenstate are fully determined by the η-unbound spinons and unbound spinons occupancies. For Bethe states with finite spin S_s and/or η-spin S_η all unbound spinons and/or unbound η-spinons have spin up and η-spin up, respectively.

Scattering theories of BA solvable models involve dressed S matrices [37–39]. A property that simplifies the pseudofermion scattering theory introduced in this paper is that unbound η-spinons and unbound spinons are found to be neither scatterers nor scattering centers. However, they play an important role in the model physics, in that they are found in this paper to fully control the effects of the transverse η-spin and spin fluctuations onto the non-LWSs η-spin and spin currents, respectively. The latter processes generate the η-spin and spin multiplet towers of non-LWSs, outside the BA solution subspace. In spite of at zero vector potential both the unbound spinons and unbound η-spinons being found to be neither scatterers nor scattering centers, they couple to uniform vector potentials. Such a coupling is behind their above effects onto the spin and η-spin currents. That in the absence of uniform vector potentials they are neither scatterers nor scattering centers follows from their momentum values remaining the same for all energy eigenstates. This property results from their invariance under the electron - rotated-electron unitary transformation studied in the first paper. That invariance simplifies the extension of the present scattering theory to excited states outside the BA solution subspace.

The spin-1/2 unbound spinons and η-spin-1/2 unbound η-spinons are the only elementary objects of the present operator formulation introduced in the first paper, Ref. [5], that have internal degrees of freedom associated with the corresponding $SU(2)$ algebras, but are neither scatterers nor scattering centers. All remaining elementary objects of that formulation are spin-neutral or η-spin neutral composite objects or spin-less and η-spin-less objects. The theory scatterers and scattering centers are slightly modified versions of the latter neutral or spin-less and η-spin-less elementary objects. Thus they have no internal degrees of freedom associated with spin or η-spin, so that their scattering is associated with fully diagonal dressed S matrices. In addition, the scatterers of the pseudofermion scattering theory include both objects created under the ground-state - excited-state transitions and those that pre-exist in the initial ground state. It is found in this paper that the phase shifts of both these two types of scatterers appear in the quantities that control the model dynamical and spectral properties. This is why the scattering theory introduced in this paper is suitable to describe such properties. On the other hand, the traditional holon and spinon scattering theories considered below account only for the phase shifts of the few scatterers created under the transitions to the excited states. Thus they miss many of the microscopic scattering events that contribute to the 1D Hubbard model dynamical and spectral properties.

The studies of this paper address the relation of the 1D Hubbard model elementary objects of the formulation used here and in the first paper, Ref. [5], to such traditional spinon and holon descriptions. As discussed in the latter paper, the elementary-object representations of the 1D Hubbard model may be classified into two large groups. One of such groups refers to elementary-object descriptions normal ordered relative to the electron and rotated-electron vacuum, which is the case of the present formulation. The corresponding elementary-object occupancy configurations generate the energy eigenstates from the electron and rotated-electron vacuum. A second and more common group of elementary-object representations are normal ordered relative to a well-defined initial ground state. This includes the traditional spinon and holon descriptions [6]. The spin-1/2 color spinors introduced in Ref. [37] for the solvable 1D Gross-Neveu model [38] whose spectrum is associated with a "hole" emerging under a transition from a $S_s = 0$ ground state to an excited state in a sequence of BA spin quantum numbers is an example of spin-1/2 spinons with no charge. The spin-1/2 spin waves introduced in Ref. [39] for the also BA solvable 1D isotropic Heisenberg antiferromagnetic model have a similar definition.

The spin-1/2 spinon description of Refs. [40–44] is a generalization for the 1D Hubbard model of such spin-1/2 objects. On the other hand, the traditional spinon and holon representations of that model are of two types in what the concept of a holon and an antiholon is concerned. Here we use as reference of such two alternative traditional holon and spinon descriptions those of Ref. [40] and Refs. [41–43], respectively. For those of which that of Ref. [40] is a representative, a holon and an antiholon refers to a "hole" and a "particle", respectively, created in the $k_j's$ BA distribution of Ref. [2] under the transition from a $S_s = 0$ ground state of arbitrary electronic density n to an excited state. Thus such descriptions refer to well-defined subspaces spanned by the excited states of ground states with arbitrary electronic density n and zero spin density, $m = 0$. On the other hand, for the descriptions of which that of Refs. [41–43] is a representative, a η-spin-1/2 holon and a η-spin-1/2 antiholon is an excitation of the $S_s = 0; S_\eta = 0; 2S_c = N_a$ absolute ground state with η-spin projection $+1/2$ and $-1/2$, respectively, associated with a "hole" created in the $k_j's$ distribution of Ref. [2] under the transition to an excited state. Hence such descriptions correspond to a well-defined subspace spanned by the excited states of that ground state.

In the case of the 1D Hubbard model, the possibility of using distinct elementary-object descriptions, such as that used here and those of Refs. [40–43] in terms of traditional holons and spinons, is found in this paper to follow from the possibility of making different choices of scattering states associated with the phase shifts extracted from the BA solution. For all the corresponding scattering theories the asymptotic scattering states must have a well-defined energy. This condition is fulfilled provided that such asymptotic scattering states refer to energy eigenstates of the 1D Hubbard model. For finite repulsive interaction values, the degeneracy of the excited energy eigenstates of $S_s = 0$ (and $S_\eta = 0$) ground states is confirmed in this paper to be behind the possibility of distinct choices of scattering states and thus of scatterers, scattering centers, and dressed S matrices. The unitary transformations relating the scattering states of the present elementary-object formulation to those of the traditional holons and spinons of the descriptions of Refs. [40] and [41–43] is derived in this paper for the 1D Hubbard model in subspaces in which the distinct scattering theories are valid. Such studies confirm that there is no contradiction whatsoever between the present elementary-object formulation and those in terms of traditional spinons and holons as defined in Refs. [40–43].

The Hubbard model Hamiltonian, under periodic boundary conditions, on a 1D lattice with a site number $N_a \gg 1$ very large and even and in a chemical potential μ and magnetic field H is given by,

$$\hat{H} = \hat{H}_{symm} - \sum_{\beta=\eta,s} \mu_\alpha \, \hat{S}_\alpha^{x_3} \, , \tag{4}$$

where

$$\hat{H}_{symm} = t\,[\hat{T} + 4u\,\hat{V}_D]\,; \quad u = U/4t\,; \quad \hat{S}^{x_3}_{\eta} = -\frac{1}{2}[N_a - \hat{N}]\,; \quad \hat{S}^{x_3}_{s} = -\frac{1}{2}[\hat{N}_{\uparrow} - \hat{N}_{\downarrow}]\,,$$

$$\hat{T} = -\sum_{\sigma=\uparrow,\downarrow}\sum_{j=1}^{N_a}\Big[c^{\dagger}_{j,\sigma}\,c_{j+1,\sigma} + c^{\dagger}_{j+1,\sigma}\,c_{j,\sigma}\Big]\,; \quad \hat{V}_D = \sum_{j=1}^{N_a}(\hat{n}_{j,\uparrow} - 1/2)\,(\hat{n}_{j,\downarrow} - 1/2)\;. \tag{5}$$

Here $\hat{T}$ is the kinetic-energy operator in units of t, $\hat{V}_D$ is the electron on-site repulsion operator in units of U, $u = U/4t$ is the electron on-site interaction in units of $4t$, which is often used in this paper, $\mu_c = 2\mu$, $\mu_s = 2\mu_B H$, μ_B is the Bohr magneton, and $\hat{S}^{x_3}_{\eta}$ and $\hat{S}^{x_3}_{s}$ are the diagonal generators of the η-spin and spin $SU(2)$ symmetry algebras [28, 30], respectively. The operator $c^{\dagger}_{j,\sigma}$ (and $c_{j,\sigma}$) that appears in the above equations creates (and annihilates) a spin-projection σ electron at lattice site $j = 1, ..., N_a$. The operator $\hat{n}_{j,\sigma} = c^{\dagger}_{j,\sigma}\,c_{j,\sigma}$ counts the number of spin-projection σ electrons at such a lattice site. The electronic number operators read $\hat{N} = \sum_{\sigma=\uparrow,\downarrow}\hat{N}_{\sigma}$ and $\hat{N}_{\sigma} = \sum_{j=1}^{N_a}\hat{n}_{j,\sigma}$. The momentum operator is given by $\hat{P} = \sum_{\sigma=\uparrow,\downarrow}\sum_k \hat{n}_{\sigma}(k)\,k$, where the spin-projection σ momentum distribution operator reads $\hat{n}_{\sigma}(k) = c^{\dagger}_{k,\sigma}\,c_{k,\sigma}$ and the operator $c^{\dagger}_{k,\sigma}$ (and $c_{k,\sigma}$) creates (and annihilates) a spin-projection σ electron of momentum k.

Throughout this paper we use in general units of both Planck constant $\hbar$ and lattice constant a one. We denote the electronic charge by e and the lattice length by $L = N_a\,a = N_a$. The LWSs have electronic densities $n = N/L$ and spin densities $m = [N_{\uparrow} - N_{\downarrow}]/L$ whose ranges obey the inequalities $0 \leq n \leq 1$ and $0 \leq m \leq n$, respectively. The description of the states corresponding to densities such that $0 \leq n \leq 1$; $1 \leq n \leq 2$ and $-n \leq m \leq n$; $-(2-n) \leq m \leq (2-n)$, respectively, is achieved by application onto the LWSs of off-diagonal generators of the η-spin and spin $SU(2)$ symmetry algebras.

The paper is organized as follows: In Section II it is found that a pseudofermion scattering theory and corresponding pseudofermion dressed S matrices naturally emerge from the operator formulation introduced in the first paper. The corresponding pseudofermion phase shifts are studied in Section III. In Section IV important issues concerning the effects of the pseudofermion transformation laws under the electron - rotated-electron unitary transformation on their scattering properties are addressed. The complementary roles played in the transport of spin by the $s\nu$ pseudofermions (and c pseudofermions at finite spin densities, due to scattering events) on the one hand and the unbound spinons on the other hand is investigated in Section V. In Section VI the relation to traditional holon and spinon scattering theories and corresponding dressed S matrices derived is investigated. Moreover, that section contains an analysis of the role played by the dressed S matrix of the present pseudofermion scattering theory in the metallic-phase finite-energy spectral distributions described by the PDT of Refs. [7, 8]. Finally, the concluding remarks are given in Section VII. Complementary useful information is provided in four appendices.

II. THE PSEUDOFERMION SCATTERING THEORY

In this paper we use the index β, which alike in the first paper refers both to the $\beta = c$ fermions and $\beta = \alpha\nu$ fermions. Here $\alpha = \eta$ and $\alpha = s$ correspond to η-spin and spin and $\nu = 1, 2, ..., \infty$ to the number of η-spinon and spinon pairs, respectively.

In this section we introduce the β fermion - β pseudofermion unitary transformation and corresponding pseudofermion and pseudofermion-hole dressed S matrices. Pseudofermion and pseudofermion-hole dressed S matrices refer here merely to β band occupied and unoccupied discrete momentum values, respectively. Symmetry implies that in spite of the spinons and η-spinons having spin 1/2 and η-spin 1/2, respectively, the scatterers and scattering centers of the present theory are spin-neutral and η-spin-neutral objects. Consistent, our analysis of the problem follows the standard quantum non-relativistic scattering theory of spin-less particles [45]. For simplicity and without loss in generality, in this section we consider densities in ranges defined by the inequalities $0 < n \leq 1$ and $0 < m \leq n$.

In Appendix A some general results useful for the studies of this paper are introduced. That includes extension of some of the results presented in the first paper to the 1D Hubbard model in a uniform vector potential.

A. The 1D Hubbard model in the pseudofermion subspace: The pseudofermion canonical momentum, associated functionals, and exotic pseudofermion algebra

The 1D Hubbard model in the pseudofermion subspace (PS) is the quantum problem that plays the major role in the studies of this paper. Such a subspace is spanned by a given initial ground state with arbitrary values for the electronic density n and spin density m and all excited energy eigenstates whose generation from it involve changes in the occupancy configurations of a finite number of β fermions, unbound spinons, and unbound η-spinons [5]. For the excited states belonging to the PS the following ratios vanish, $\delta N_{\beta}/N_a \rightarrow 0$, $\delta M_s^{un}/N_a \rightarrow 0$, and $\delta M_{\eta}^{un}/N_a \rightarrow 0$, as $N_a \rightarrow \infty$. Here δN_{β}, δM_s^{un}, and δM_{η}^{un} denote the deviations in the numbers N_{β}, $M_s^{un} = 2S_s$, and $M_{\eta}^{un} = 2S_{\eta}$ of β fermions, unbound spinons, and unbound η-spinons, respectively, under the ground-state - excited-state transitions. The numbers of unbound η-spinons and unbound spinons are defined in Eq. (3). We emphasize that within the definition of the PS there are no restrictions on the value of the excitation energy and excitation momentum.

Several subspaces contained in a PS subspace play an important role in the pseudofermion scattering theory introduced here. The following three types of subspaces are contained in well-defined subspaces considered in the first paper.

– An *electronic ensemble space* is a subspace spanned by all energy eigenstates with the same values for the electronic numbers $N_{\uparrow}$ and $N_{\downarrow}$. Such a subspace is contained in the *unbound η-spinon and unbound spinon subspace* considered in the first paper. The latter is spanned by the set of energy eigenstates with fixed values for the numbers $N_c = 2S_c$, $M_{\eta}^{un} = 2S_{\eta}$, and $M_s^{un} = 2S_s$. On the other hand, an electronic ensemble space is spanned by the set of energy eigenstates with fixed values for the numbers of unbound η-spinons and unbound spinons $M_{\eta,\pm 1/2}^{un} = [S_{\eta} \mp S_{\eta}^{x_3}]$ and $M_{s,\pm 1/2}^{un} = [S_s \mp S_s^{x_3}]$, respectively, defined in Eq. (3).

– A *CPHS ensemble space* is a subspace spanned by all energy eigenstates with the same values for the numbers $\{M_{\alpha,\pm 1/2}\}$ of $\pm 1/2$ η-spinons ($\alpha = \eta$) and $\pm 1/2$ spinons ($\alpha = s$). (The designation CPHS is that used in the studies of Refs. [7, 8, 33, 34]. Within the notation of such references, it refers to c pseudofermions (CP), holons (H), and spinons (S). The holons are the η-spinons of the present formulation.) Such a subspace is contained in the *η-spinon and spinon subspace* considered in the first paper. The latter is spanned by the set of energy eigenstates with fixed values for the corresponding numbers $M_{\eta} = [N_a - 2S_c]$ and $M_s = 2S_c$, respectively. The CPHS ensemble space considered here is spanned by the set of energy eigenstates with fixed values for the numbers $M_{\eta,\pm 1/2} = [M_{\eta,\pm 1/2}^{un} + M_{\eta}^{bo}/2]$ and $M_{s,\pm 1/2} = [M_{s,\pm 1/2}^{un} + M_s^{bo}/2]$.

– Finally, a *CPHS ensemble subspace* is spanned by all energy eigenstates with the same values for the sets of numbers $\{M_{\alpha,-1/2}^{un}\}$ where $\alpha = \eta, s$ and $\{N_{\beta}\}$ where $\beta = c, \alpha\nu$. Here $\alpha = \eta, s$ and $\nu = 1, 2, ..., \infty$. Such a subspace is contained in the *extended Takahashi subspace* considered in the first paper. The latter is spanned by the set of energy eigenstates with fixed values for the sets of numbers $\{M_{\alpha}^{un}\} = \{2S_{\alpha}\}$ where $\alpha = \eta, s$ and $\{N_{\beta}\}$ where $\beta = c, \alpha\nu$ and again $\alpha = \eta, s$ and $\nu = 1, 2, ..., \infty$.

At zero absolute temperature the pseudofermion description corresponds to a ground-state normal-ordered theory. Thus, there is a pseudofermion theory for each initial ground state. For densities $n \in [0, 1]$ and $m \in [0, n]$ the minimum excitation energy value of the energy eigenstates that span a given CPHS ensemble subspace is given by,

$$\omega_0 = \omega_0(D_r, S_r) = 2|\mu|\, D_r + 2\mu_B\, |H|\, S_r\,. \tag{6}$$

The numbers D_r and S_r appearing here are defined in Eq. (A19) of Appendix A. For the ground state, $D_r = S_r = 0$ and thus $\omega_0 = 0$. The application onto the latter state of a one-, two-, or any other finite-number-electron operator $\mathcal{O}^{\dagger}$ generates an excitation that can be described as a well-defined superposition of PS excited energy eigenstates. For some of the finite-energy line shape branches in the one- and two-electron spectral weight distributions, the PDT provides spectral weights associated with the coefficients of such a superposition [7, 8].

An excitation $\mathcal{O}^{\dagger}|GS\rangle$ associated with small values of the deviations $\delta N_{\uparrow}$ and $\delta N_{\downarrow}$ is contained in a well defined direct sum of CPHS ensemble subspaces,

$$\mathcal{S}_{cphs}^{1} \oplus \mathcal{S}_{cphs}^{2} \oplus \mathcal{S}_{cphs}^{3} \oplus \mathcal{S}_{cphs}^{4} \oplus \ldots\,. \tag{7}$$

Here $\mathcal{S}_{cphs}^{i}$ with $i = 1, 2, 3, ...$ corresponds to different CPHS ensemble subspaces. The β pseudofermion, unbound η-spinon, and unbound spinon number deviations of all the CPHS ensemble subspaces of such a direct sum obey the sum rules provided in Eqs. (18) and (19) of Ref. [7] and the selection rules given in Eq. (21) of the same reference. (The c and $c\nu$ branches of such equations are denoted here by c and $\eta\nu$, respectively, whereas the numbers $L_{c,-1/2}$ and $L_{s,-1/2}$ read within our notation $M_{\eta,-1/2}^{un}$ and $M_{s,-1/2}^{un}$, respectively.)

For the energy eigenstates that span the PS the rapidity functionals defined by the set of coupled integral thermodynamic BA equations, Eqs. (A2)-(A4) of Appendix A for $\phi_{\uparrow} = \phi_{\downarrow} = 0$, have the following exact property,

$$\begin{aligned}
k_c(q_j) &= k_c^0\Big(\bar{q}(q_j)\Big)\,; \quad j = 1, ..., N_a\,, \\
\Lambda_c(q_j) &= \sin k_c(q_j) = \sin k_c^0\Big(\bar{q}(q_j)\Big)\,; \quad j = 1, ..., N_a\,, \\
\Lambda_{\beta}(q_j) &= \Lambda_{\beta}^0\Big(\bar{q}(q_j)\Big)\,; \quad j = 1, ..., N_{a_{\alpha\nu}}\,.
\end{aligned} \tag{8}$$

The set of discrete numbers $\bar{q}_j = \bar{q}(q_j)$ where $j = 1, ..., N_{a_\beta}$ in the arguments of the functions $k_c^0\Big(\bar{q}(q_j)\Big)$ for $\beta = c$ and $\Lambda_\beta^0\Big(\bar{q}(q_j)\Big)$ for $\beta = \alpha\nu$ appearing in Eq. (8) play a central role in the pseudofermion scattering theory. Their spacing is to first order in $1/N_a$,

$$\bar{q}_{j+1} - \bar{q}_j = \frac{2\pi}{N_a} + \text{h.o.}\,. \tag{9}$$

Such discrete numbers $\bar{q}_j = \bar{q}(q_j)$ are the β pseudofermion discrete canonical-momentum values. A β pseudofermion carries discrete canonical momentum $\bar{q}_j = \bar{q}(q_j)$ whereas the corresponding β fermion carries the discrete momentum q_j. Often in the remaining of this paper we call q_j bare momentum, to distinguish it from the corresponding canonical momentum, $\bar{q}_j = \bar{q}(q_j)$.

For $\beta = \eta\nu$ (and $\beta = s\nu \neq s1$) branches, the relation $\Lambda_\beta(q_j) = \Lambda_\beta^0\Big(\bar{q}(q_j)\Big)$ given in Eq. (8) is valid provided that the hole concentration $(1-n)$ (and the spin density m) is finite. For an initial ground state with $S_\eta = 0$ and thus electronic density $n = 1$ (and $S_s = 0$ and spin density $m = 0$) one has that the number of $\eta\nu$ (and $s\nu \neq s1$) band discrete momentum values $N_{a_{\eta\nu}}$ (and $N_{a_{s\nu}}$) given in Eqs. (A10) and (A11) of Appendix A vanishes, $N_{a_{\eta\nu}} = 0$ (and $N_{a_{s\nu}} = 0$). Therefore, the corresponding pseudofermion branch does not exist. Hence the ground-state rapidity Λ_β^0 appearing in Eq. (8) is undefined. For the excited states of such a $n = 1$ (and $m = 0$) ground state the value of $\Lambda_\beta(q_j)$ for $\beta = \eta\nu$ (and $\beta = \alpha\nu \neq s1$) is an issue addressed below in Section IV B.

By use of the expressions provided in Eq. (8) in the thermodynamic BA equations, Eqs. (A2)-(A4) of Appendix A at $\phi_\uparrow = \phi_\downarrow = 0$, one uniquely finds that, to leading orders in $1/N_a$, the discrete canonical-momentum values have the following form,

$$\bar{q}_j = \bar{q}(q_j) = q_j + \frac{Q_\beta^\Phi(q_j)}{N_a} = \frac{2\pi}{N_a} I_j^\beta + \frac{Q_\beta^\Phi(q_j)}{N_a}\,; \quad \beta = c, \alpha\nu\,; \quad j = 1, ..., N_{a_\beta}\,. \tag{10}$$

Here N_{a_β} reads $N_{a_c} = N_a$ for $\beta = c$ and is given in Eqs. (A10) and (A11) of Appendix A for $\beta = \alpha\nu$. The relation provided in Eq. (10) uniquely defines a one-to-one correspondence between the two sets $\{q_j\}$ and $\{\bar{q}_j\}$ such that $j = 1, ..., N_{a_\beta}$, which refers to the β fermion - β pseudofermion unitary transformation. As given in the first paper, the β band discrete momentum value q_j can be written as $q_j = [2\pi/N_a] I_j^\beta$, where I_j^β with $j = 1, ..., N_{a_\beta}$ are BA quantum numbers. Depending on uniquely defined BA solution boundary conditions, those can have either integer or half-odd integer values.

Except for the slightly shifted discrete momentum values given in Eq. (10), the β pseudofermions have the same properties as the corresponding β fermions. The quantity $Q_\beta^\Phi(q_j)$ in $Q_\beta^\Phi(q_j)/N_a$ appearing in that equation plays a key role in the pseudofermion scattering theory. It reads,

$$Q_\beta^\Phi(q_j) = \sum_{\beta'} \sum_{j'=1}^{N_{a_{\beta'}}} 2\pi\, \Phi_{\beta,\beta'}(q_j, q_{j'})\, \delta N_{\beta'}(q_{j'})\,. \tag{11}$$

The β' band momentum distribution function deviation $\delta N_{\beta'}(q_{j'})$ appearing here is that defined in Eq. (A18) of Appendix A. Since our elementary-object formulation refers to the limit of a very large system, $N_a \gg 1$, we approximate the discrete momentum values q_j, such that $[q_{j+1} - q_j] = 2\pi/N_a$, by the a continuum variable q. Alike in the first paper, the general β band momentum distribution function deviation then reads,

$$\begin{aligned}
\delta N_\beta(q) &= \frac{2\pi}{N_a}\sum_{p=1}^{N_\beta^p} \delta(q - q_p) - \frac{2\pi}{N_a}\sum_{h=1}^{N_\beta^h} \delta(q - q_h)\,; \;\; \text{if } \delta G_\beta \text{ even for } \beta = c, s1\,, \\
&= \frac{2\pi}{N_a}\sum_{p=1}^{N_\beta^p} \delta(q - q_p) - \frac{2\pi}{N_a}\sum_{h=1}^{N_\beta^h} \delta(q - q_h) \pm \frac{\pi}{N_a}\sum_{j=\pm 1} j\, \delta(q - j\, q_{F\beta})\,; \;\; \text{if } \delta G_\beta \text{ odd for } \beta = c, s1\,, \\
\delta N_\beta(q) &= \frac{2\pi}{N_a}\sum_{p=1}^{N_\beta^p} \delta(q - q_p)\,; \;\; \text{for } \beta = \alpha\nu \neq s1\,,
\end{aligned} \tag{12}$$

Here and throughout this paper $\delta(x)$ denotes the usual Dirac delta-function distribution, $q_1, ..., q_{N_\beta^p}$ are the momentum values of the N_β^p added β fermions, $q_1, ..., q_{N_\beta^h}$ those of the N_β^h added β fermion holes, and δG_β stands for the deviation in the value of the number G_β given by,

$$G_\beta = \delta_{\beta,c}\sum_{\beta=\eta,s} B_\alpha + \delta_{\beta,s1}[N_c + N_{s1}]\,; \qquad B_\alpha = \sum_{\nu=1}^{\infty} N_{\alpha\nu}\,, \quad \alpha = \eta, s\,. \tag{13}$$

The additional term in the expression given in Eq. (12) for the deviation δG_β being a odd integer number relative to that for it being an even integer number results from the corresponding $\beta = c, s1$ band momentum shift $\pm\pi/N_a$.

Moreover, the quantity $\pi\,\Phi_{\beta,\beta'}(q_j, q_{j'})$ in Eq. (11) is a function of both the bare-momentum values q_j and $q_{j'}$, given by,

$$\pi\,\Phi_{\beta,\beta'}(q_j, q_{j'}) = \pi\,\bar{\Phi}_{\beta,\beta'}\left(\frac{\Lambda^0_\beta(q_j)}{u}, \frac{\Lambda^0_{\beta'}(q_{j'})}{u}\right)\,. \tag{14}$$

The function $\pi\,\bar{\Phi}_{\beta,\beta'}(r, r')$ appearing here is the unique solution of the integral equations (B1)-(B15) of Appendix B and the ground-state rapidity functions $\Lambda^0_\beta(q_j)$, where $\Lambda^0_c(q_j) = \sin k^0_c(q_j)$ for $\beta = c$, are defined in terms of their inverse functions in Eq. (A14) of Appendix A.

It is found below that $\pi\,\Phi_{\beta,\beta'}(q, q')$ [or $-\pi\,\Phi_{\beta,\beta'}(q, q')$] is an elementary *two-pseudofermion phase shift*. It is such that q is the bare-momentum value of a β pseudofermion or β pseudofermion hole scattered by a β' pseudofermion [or β' pseudofermion hole] of bare-momentum q' created under a ground-state - excited-energy-eigenstate transition. (Since there is a one-to-one correspondence between the canonical momentum $\bar{q}(q)$ and the bare-momentum q, *the bare momentum q of a pseudofermion* is meant as that corresponding to its canonical momentum $\bar{q}(q)$.)

An important functional related to that defined in Eq. (11) reads,

$$Q_\beta(q_j) = Q^0_\beta + Q^\Phi_\beta(q_j)\,. \tag{15}$$

Here Q^0_β is such that the quantity Q^0_β/N_a is the shift in the $\beta = c, \alpha\nu$ band discrete bare-momentum value q_j under the ground-state - excited-state transition. Its values are,

$$\begin{aligned} Q^0_c &= 0\,; \quad \sum_{\beta=\eta,s}\sum_{\nu=1}^{\infty}\delta N_{\alpha\nu} \text{ even}\,; \qquad Q^0_c = \pm\pi\,; \quad \sum_{\beta=\eta,s}\sum_{\nu=1}^{\infty}\delta N_{\alpha\nu} \text{ odd}\,; \\ Q^0_{\alpha\nu} &= 0\,; \quad \delta N_c + \delta N_{\alpha\nu} \text{ even}\,; \qquad Q^0_{\alpha\nu} = \pm\pi\,; \quad \delta N_c + \delta N_{\alpha\nu} \text{ odd}\,; \quad \alpha = \eta, s\,. \end{aligned} \tag{16}$$

When under such a transition the BA quantum numbers I^β_j in the β band discrete momentum value $q_j = [2\pi/N_a]I^\beta_j$ change from integers (or half-odd integers) to half-odd integers (or integers), the shift Q^0_β/N_a has a finite value, $Q^0_\beta/N_a = \pm\pi/N_a$.

To first order in $1/N_a$, the discrete canonical-momentum values spacing is, $\bar{q}_{j+1} - \bar{q}_j = 2\pi/N_a$, as given in Eq. (9). The one-to-one correspondence between the discrete canonical momentum values and the discrete bare momentum values given in Eq. (10) is associated with the β fermion - β pseudofermion unitary transformation. That transformation is defined within the PS. There is a pseudofermion representation for each initial ground state and corresponding PS. Analysis of the the discrete canonical-momentum value expression defined by Eqs. (10) and (11) reveals that for the initial ground state the discrete canonical-momentum value $\bar{q}_j$ and corresponding discrete bare-momentum value q_j are such that $\bar{q}_j = q_j$. Hence the ground-state limiting β canonical momenta and c and $s1$ Fermi canonical momenta equal the corresponding bare momenta. Those are given in Eqs. (A12) and (A13) of Appendix A, respectively.

As mentioned above, there are no $\eta\nu$ (and $s\nu \neq s1$) bare-momentum bands for $n = 1$ (and $m = 0$) ground states. Indeed, from use of Eqs. (A10) and (A11) of Appendix A one confirms that $N_{a_{\eta\nu}} = 0$ (and $N_{a_{s\nu}} = 0$) for such states. Then the corresponding ground-state rapidity functions $\Lambda^0_{\eta\nu}(q)$ (and $\Lambda^0_{s\nu}(q)$) cannot be defined. Fortunately, the expression given in Eq. (14) remains valid in that case provided that the ground-state rapidity functions are suitably replaced by those of the excited states. (We find below that the functions provided in Eq. (14) are phase shifts originated by well-defined ground-state - excited-state transitions. Thus, in the particular case of the initial $n = 1$ and/or $m = 0$ ground states, the quantities given in Eq. (14) are functionals rather than functions, with the rapidity functions for the $\eta\nu$ and/or $s\nu \neq s1$ branches being those of the excited state under consideration.)

The energy spectrum of Eq. (A17) of Appendix A can be expressed in terms of the corresponding β pseudofermion canonical-momentum distribution-function deviations as follows,

$$\delta E_{l_o,l_\triangle} = \sum_{\beta=c,s1}\sum_{j=1}^{N_{a_\beta}} \varepsilon_\beta(\bar{q}_j)\delta\mathcal{N}_\beta(\bar{q}_j) + \sum_{\beta\nu\neq s1}\sum_{j=1}^{N_{a_{\alpha\nu}}} \varepsilon^0_{\alpha\nu}(\bar{q}_j)\delta\mathcal{N}_{\alpha\nu}(\bar{q}_j) + 2|\mu|D_r + 2\mu_B|H|S_r\,. \tag{17}$$

Alike in the first paper and in Appendix A, here $l_{\triangle}$ stands for the set of numbers $[2S_c, S_\eta, S_s, M^{un}_{\eta,-1/2}, M^{un}_{s,-1/2}]$ and l_o for all remaining quantum numbers beyond $l_{\triangle}$ needed to uniquely define an energy eigenstate $|\Psi_{l_o,l_{\triangle},u}\rangle$ of general form given in Eq. (A6) of that appendix. For the Bethe states one has that $M^{un}_{\eta,-1/2} = M^{un}_{s,-1/2} = 0$, so that the above set of number values read $[2S_c, S_\eta, S_s, 0, 0]$. In that case one says that $l_{\triangle} = l^0_{\triangle}$.

In the expression provided in Eq. (17) we have used the results of the first paper to express the finite-energy part of the energy spectrum given in Eq. (A17) of Appendix A in terms of the number of rotated-electron doubly occupied sites, D_r, and that of spin-down rotated-electron singly occupied sites whose rotated electrons are not associated with the $s1$ fermions and thus $s1$ pseudofermions, S_r, given in Eq. (A19) of Appendix A. They vanish for any initial ground state, so that their deviations are given by $\delta D_r = D_r$ and $\delta S_r = S_r$.

Importantly, when expressed in terms of β fermion momentum distribution-function deviations, Eq. (A18) of Appendix A, the energy spectrum given in Eq. (17) contains all the contributions to second order in such deviations. Those are beyond the corresponding first-order expression, Eq. (A17) of Appendix A. Indeed, such a β fermion energy spectrum has higher-order energy functional terms in the deviations $\delta N_\beta(q_j)$, beyond those given in that equation. Within the $N_a \gg 1$ limit that the current operator formulation refers to only the terms up to second order in such deviations are physically meaningful. Such terms stem from contributions arising from expanding the PS rapidity functions $k^0_c(q_j + Q^\Phi_c(q_j)/N_a)$ and $\Lambda^0_\beta(q_j + Q^\Phi_\beta(q_j)/N_a)$ associated with the functions $k_c(q_j) = k^0_c(\bar{q}(q_j))$ and $\Lambda_\beta(q_j) = \Lambda^0_\beta(\bar{q}(q_j))$ of Eq. (8) around q_j. Within the β pseudofermion representation the quantity $Q^\Phi_c(q_j)/N_a$ (and $Q^\Phi_\beta(q_j)/N_a$) in the argument of the rapidity function $k^0_c(q_j + Q^\Phi_c(q_j)/N_a)$ (and $\Lambda^0_\beta(q_j + Q^\Phi_\beta(q_j)/N_a)$) is rather incorporated in the canonical momentum $\bar{q}(q_j) = q_j + Q^\Phi_c(q_j)/N_a$ (and $\bar{q}(q_j) = q_j + Q^\Phi_\beta(q_j)/N_a$).

That the energy spectrum given in Eq. (17) contains all the contributions of second order in the β fermion momentum distribution-function deviations is behing a property of major importance for the PDT. It is that within the $N_a \gg 1$ limit specific to that theory the c and $s1$ pseudofermions have no residual energy interactions and thus are not energy entangled [7, 8]. This follows from the second-order deviation contributions energy functional terms in the deviations $\delta N_\beta(q_j)$ beeing transferred to the $\beta = c, s1$ band discrete canonical momentum values. Such a lack of β pseudofermion energy entanglement then allows the one- and two-electron spectral functions to be expressed as convolutions of suitable c and $s1$ fermion spectral functions [7, 8]. In contrast, the related c and $s1$ fermions have residual energy interactions and thus are energy entangled.

To confirm that for $N_a \gg 1$ the c and $s1$ pseudofermions have no residual energy interactions we consider for simplicity PS excited states for which $M^{un}_{\eta,-1/2} = M^{un}_{s,-1/2} = 0$ in Eq. (3) and the momentum distribution function deviations $\delta N_\beta(q_j)$ in Eq. (A17) of Appendix A vanish for the $\beta \neq c, s1$ branches. Since our analysis refers to the limit of a very large system, $N_a \gg 1$, we approximate again the discrete momentum values q_j and now also the discrete canonical momentum values $\bar{q}_j$ by the continuum variables q and $\bar{q} = q + Q^\Phi_\beta(q)/N_a$, respectively. The corresponding deviation $\delta N_\beta(q)$ has the general form given in Eq. (12).

Next we extend the expression of the energy functional provided in Eq. (A17) of Appendix A to second order in the $\beta = c, s1$ momentum distribution function deviations $\delta N_\beta(q)$. To reach that goal, we use such momentum distribution function deviations in the functional BA thermodynamic equations, Eqs. (A2)-(A4) of Appendix A at $\phi_\uparrow = \phi_\downarrow = 0$, and corresponding general energy functional given in Eq. (A8) of Appendix A. The combined and consistent solution of the obtained equations up to second order in the deviations $\delta N_\beta(q)$ then leads to the energy functional,

$$\begin{aligned}\delta E_{l_o,l_{\triangle}} &= \frac{N_a}{2\pi}\sum_{\beta=c,s1}\int_{-q_\beta}^{q_\beta} dq\,\varepsilon_\beta(q)\,\delta N_\beta(q) \\ &+ \sum_{\beta=c,s}\left[\frac{1}{2\pi}\int_{-q_\beta}^{q_\beta} dq\,v_\beta(q)\,Q^\Phi_\beta(q)\delta N_\beta(q) + \frac{v_\beta}{2\pi N_a}\sum_{\iota=\pm 1}\left(Q^\Phi_\beta(\iota q_{F\beta})\right)^2\right]. \qquad (18)\end{aligned}$$

The energy dispersions $\varepsilon_\beta(q)$ and velocities $v_\beta(q)$ and $v_\beta \equiv v_\beta(q_{F\beta})$ appearing here are those defined in Eqs. (A20)-(A22) and Eq. (A23) of Appendix A, respectively.

The use of the functional $Q^\Phi_\beta(q)$ expression given in Eq. (11) reveals that the energy functional of Eq. (18) can be rewritten as,

$$\begin{aligned}\delta E_{l_o,l_{\triangle}} &= \frac{N_a}{2\pi}\sum_{\beta=c,s1}\int_{-q_\beta}^{q_\beta} dq\,\varepsilon_\beta(q)\delta N_\beta(q) \\ &+ \frac{1}{N_a}\left(\frac{N_a}{2\pi}\right)^2\sum_{\beta=c,s1}\sum_{\beta'=c,s}\int_{-q_\beta}^{q_\beta} dq\int_{-q_{\beta'}}^{q_{\beta'}} dq'\frac{1}{2}\,f_{\beta,\beta'}(q,q')\,\delta N_\beta(q)\delta N_{\beta'}(q'), \qquad (19)\end{aligned}$$

where the f functions read,

$$f_{\beta,\beta'}(q,q') = v_\beta(q)\,2\pi\,\Phi_{\beta,\beta'}(q,q') + v_{\beta'}(q')\,2\pi\,\Phi_{\beta',\beta}(q',q) + \frac{1}{2\pi}\sum_{\beta''=c,s1}\sum_{\iota=\pm 1} v_{\beta''}\,2\pi\,\Phi_{\beta'',\beta}(\iota q_{F\beta''},q)\,2\pi\,\Phi_{\beta'',\beta'}(\iota q_{F\beta''},q')\,. \tag{20}$$

Interestingly, such f functions involve only the β fermion group velocities and the two-pseudofermion phase shifts of the scattering theory introduced in this paper.

For low-energy excitations, both the energy terms in Eq. (18) of first-order and second-order in the β fermion momentum distribution function deviations $\delta N_\beta(q)$ become of order $1/N_a$. In that case one needs both the first-order and second-order energy terms in such deviations to reach the finite-size spectrum of conformal-field theory [8]. The corresponding finite-size corrections must be accounted for before taking the $N_a\rightarrow\infty$ thermodynamic limit. Within that limit, only the residual energy interactions of second order in the β fermion momentum distribution function deviations $\delta N_\beta(q)$ are physically meaningful, as mentioned above. Their occurrence implies that the $\beta = c, s1$ fermions have residual interactions and thus are energy entangled.

We now confirm that up to energy contributions of order $1/N_a$, which are those revelant within the present $N_a\rightarrow\infty$ thermodynamic limit, the c and $s1$ pseudofermions have no residual energy interactions and thus are not energy entangled. Expression of the general energy functional given in Eq. (18) in terms of the $\beta = c, s1$ pseudofermion canonical momenta leads to,

$$\delta E_{l_o,l_\triangle} = \frac{N_a}{2\pi}\sum_{\beta=c,s1}\int_{-q_\beta}^{q_\beta} d\bar{q}\,\varepsilon_\beta(\bar{q})\,\delta\bar{N}_\beta(\bar{q})\,. \tag{21}$$

Here $\bar{q}$ is the β pseudofermion canonical momentum defined in Eq. (10) and to first order in $1/N_a$ the deviation $\delta\bar{N}_\beta(\bar{q})$ accounts for the two types of contribution reported in the following.

The first contribution type results from the shift $q\rightarrow q + Q^\Phi_\beta(q)/N_a$ in the arguments of the δ-functions of the deviation $\delta N_\beta(q)$ expression provided in Eq. (12). Such a contribution is taken into account by considering that $\varepsilon_\beta(\bar{q}) = \varepsilon_\beta(q)$ and $\delta\bar{N}_\beta(\bar{q}) = \delta N_\beta(q + Q^\Phi_\beta(q)/N_a)$ in Eq. (21). It leads to the energy terms $\frac{N_a}{2\pi}\sum_{\beta=c,s1}\int_{-q_\beta}^{q_\beta} dq\,\varepsilon_\beta(q)\,\delta N_\beta(q)$ and $\frac{1}{2\pi}\sum_{\beta=c,s}\int_{-q_\beta}^{q_\beta} dq\,v_\beta(q)\,Q^\Phi_\beta(q)\delta N_\beta(q)$ of Eq. (18). To first order in $1/N_a$, exactly the same energy terms are obtained if one considers instead that $\varepsilon_\beta(\bar{q}) = \varepsilon_\beta(q + Q^\Phi_\beta(q)/N_a)$ and $\delta\bar{N}_\beta(\bar{q}) = \delta N_\beta(q)$ in Eq. (21). This latter choice is the most convenient for adding to $\delta\bar{N}_\beta(\bar{q})$ the second contribution reported below.

The first contribution refers to small changes in the momenta of the original β fermion creation and annihilation processes described by the deviations of form given in Eq. (12). On the other hand, the second contribution to the β pseudofermion canonical-momentum distribution function deviation $\delta\bar{N}_\beta(\bar{q})$ does not refer to small changes of processes contained in the the original β band momentum distribution function deviation $\delta N_\beta(q)$. Specifically, it involves the quantity $\iota\,Q^\Phi_\beta(\iota\,q_{F\beta})/N_a = \iota[\delta\bar{q}^\iota_{F\beta} - \delta q^\iota_{F\beta}]$ where $\delta\bar{q}^\iota_{F\beta}$ and $\delta q^\iota_{F\beta}$ are the deviations of the β pseudofermion and β fermion right ($\iota = 1$) and left ($\iota = -1$) Fermi points, which read,

$$\delta\bar{q}^\iota_{F\beta} = \delta q^\iota_{F\beta} + \frac{Q^\Phi_\beta(\iota q_{F\beta})}{N_a}\,,$$
$$\delta q^\iota_{F\beta} = \iota\,\frac{2\pi}{N_a}\delta N^F_{\beta,\iota}\,, \tag{22}$$

respectively. The quantity $\delta N^F_{\beta,\iota} = [\delta N^{0,F}_{\beta,\iota} + \iota\,Q^0_\beta/2\pi]$ denotes the deviation in the number of $\beta = c, s1$ fermions at the right ($\iota = +1$) and left ($\iota = +1$) $\beta = c, s1$ Fermi points under the ground-state - excited-state transition. Such a deviation includes the effects from the possible shift of the BA quantum numbers I^β_j considered in the first paper. That shift involves the quantity Q^0_β for the $\beta = c, s1$ branches considered here, which is defined for all β branches in Eq. (16). On the other hand, $\delta N^{0,F}_{\beta,\iota}$ does not account for such effects. Indeed the value of $\delta N^{0,F}_{\beta,\iota}$ follows only from the change in the β fermion occupancies at their $\iota = \pm 1$ Fermi points. We emphasize that the second term $Q^\Phi_\beta(\iota q_{F\beta})/N_a$ in the deviation $\delta\bar{q}^\iota_{F\beta}$ expression provided in Eq. (22), which is that associated with the second contribution type to $\delta\bar{N}_\beta(\bar{q})$, occurs in that expression even when $\delta q^\iota_{F\beta} = 0$, and thus there is no change in the $\iota = \pm 1$ β fermion Fermi-point occupancies under the ground-state - excited-state transition.

To first order in $1/N_a$, the above two types of contribution are accounted for provided that one uses the following expressions for $\varepsilon_\beta(\bar{q})$ and $\delta\bar{N}_\beta(\bar{q})$ in the energy functional of Eq. (21),

$$\begin{aligned}
\varepsilon_\beta(\bar{q}) &= \varepsilon_\beta(q + Q^\Phi_\beta(q)/N_a) \approx \varepsilon_\beta(q) + v_\beta(q)\,\frac{Q^\Phi_\beta(q)}{N_a} \\
\delta\bar{N}_\beta(\bar{q}) &= \delta N_\beta(q) + \iota\frac{Q^\Phi_\beta(\iota\, q_{F\beta})}{N_a}\,\delta(q - \iota\, q_{F\beta})\,. \qquad (23)
\end{aligned}$$

Consistent with our above discussion, the extra term $\iota[Q^\Phi_\beta(\iota\, q_{F\beta})/N_a]\,\delta(q-\iota\, q_{F\beta})$ in the $\delta\bar{N}_\beta(\bar{q})$ expression accounts for the second type of contribution to it. The use on the right-hand side of Eq. (21) of the expressions provided in Eq. (23) readily leads to the full energy expression given in Eq. (18), which is of second order in the β fermion momentum distribution function deviations $\delta N_\beta(q)$. (We recall that $\varepsilon_\beta(\iota\, q_{F\beta}) = 0$, so that the second type of contribution leads indeed only to the energy term $\sum_{\beta=c,s}\frac{v_\beta}{2\pi N_a}\sum_{\iota=\pm1}(Q^\Phi_\beta(\iota q_{F\beta}))^2$ in Eq. (18).)

On the one hand, within the β fermion representation the quantity $Q^\Phi_c(q)/N_a$ (and $Q^\Phi_\beta(q)/N_a$) in the argument of the rapidity function $k^0_c(q + Q^\Phi_c(q)/N_a)$ (and $\Lambda^0_\beta(q + Q^\Phi_\beta(q)/N_a)$) is behind the β fermion residual interactions. This is confirmed by the form of the second-order energy terms in the first expression of Eq. (18). Those would vanish if $Q^\Phi_\beta(q) = 0$. On the other hand, we have just confirmed that since within the β pseudofermion representation that quantity is rather incorporated in the canonical momentum, such objects are not energy entangled: The momentum shift $Q^\Phi_\beta(q)/N_a$ in the β pseudofermion canonical momentum $\bar{q} = q + Q^\Phi_\beta(q)/N_a$ of Eq. (10) exactly cancels the second-order energy terms in the expression given in Eq. (18).

That within the β pseudofermion representation the quantity $Q^\Phi_\beta(q)$ is incorporated in the canonical momentum $\bar{q} = q + Q^\Phi_\beta(q)/N_a$ of Eq. (10) renders the β pseudofermions being not energy entangled yet has consequences in the exotic algebra obeyed by their creation and annihilation operators. Consider a β pseudofermion of canonical momentum $\bar{q}$ and a β' pseudofermion of canonical momentum $\bar{q}'$ such that the canonical-momentum values $\bar{q}$ and $\bar{q}' = q'$ correspond to an excited energy eigenstate and the initial ground state, respectively. That the canonical momentum $\bar{q} = q + Q^\Phi_\beta(q)/N_a$ of the former β pseudofermion includes the momentum shift $Q^\Phi_\beta(q)/N_a$ is behind the effective anticommutators involving the creation and/or annihilation operators of these two pseudofermions having the general form [7],

$$\begin{aligned}
\{f^\dagger_{\bar{q},\beta}, f_{\bar{q}',\beta'}\} &= \delta_{\beta,\beta'}\frac{1}{N_{a_\beta}}\,e^{-i(\bar{q}-\bar{q}')/2}\,e^{iQ_\beta(q)/2}\,\frac{\sin\Big(Q_\beta(q)/2\Big)}{\sin([\bar{q}-\bar{q}']/2)}\,, \\
\{f^\dagger_{\bar{q},\beta}, f^\dagger_{\bar{q}',\beta'}\} &= \{f_{\bar{q},\beta}, f_{\bar{q}',\beta'}\} = 0\,. \qquad (24)
\end{aligned}$$

Here $Q_\beta(q_j) = Q^0_\beta + Q^\Phi_\beta(q_j)$ is the value of the functional defined in Eq. (15) specific to the excited energy eigenstate under consideration. The related quantity $Q_\beta(q_j)/N_a = [Q^0_\beta + Q^\Phi_\beta(q_j)]/N_a$ is the overall shift in the discrete canonical-momentum value that results from the ground-state - excited-energy-eigenstate transition.

The exotic β pseudofermion algebra associated with the effective anticommutators of form provided in Eq. (24) plays a key role in the one- and two-electron finite-energy spectral weight distributions [7, 8]. It follows from that equation that the functional $Q_\beta(q)$ controls the quantum overlaps associated with such spectral weight distributions. Indeed, the one- and two-electron matrix elements between the initial ground state and the excited energy eigenstates are within the PDT expressed in terms of the corresponsing effective anticommutators, Eq. (24) [7, 8].

The effective character of the anticommutators of Eq. (24) results from they involving two operators acting onto subspaces with different discrete canonical momentum values. Such subspaces are those of the initial ground state and excited state, respectively. That they connect two different Hilbert spaces, whose discrete quantum-number values are different due to shake-up effects, makes them to be effective anticommutators. Indeed, the standard operator commutators involve operators acting onto the same Hilbert space. However, such effective anticommutators are physically meaningful. They control the orthogonal-catrastrophe quantum overlaps associated with the PDT one- and two-electron spectral-weight distributions [7, 8].

B. The ground-state - virtual-state transition

From now on and until Section IV B our analysis refers to initial ground states with density values in the ranges $n \in [0,1[$ and $m \in]0,n]$. (The specific properties of the scattering theory for initial ground states corresponding to densities $n = 1$ and/or $m = 0$ are considered in that section.) Each transition from the initial ground state to a PS excited energy eigenstate can be divided into three steps. The first process is a scatter-less finite-energy and finite-momentum excitation that transforms the ground state onto a well defined virtual state. For the $\alpha\nu$ branches, that excitation can involve a change in the number of discrete bare-momentum values given by,

$$\delta N_{a_{s1}} = \delta N_c - \delta N_{s1} - 2\sum_{\nu=2}^{\infty}\delta N_{s\nu}\,; \qquad \delta N_{a_{\alpha\nu}} = \delta M^{un}_{\alpha} + 2\sum_{\nu'=\nu+1}^{\infty}(\nu'-\nu)\delta N_{\alpha\nu'}\,, \quad \alpha\nu \neq s1\,. \tag{25}$$

For the initial ground state these numbers read,

$$N^0_{a_{s1}} = N_{\uparrow}\,; \qquad N^0_{a_{s\nu}} = (N_{\uparrow} - N_{\downarrow})\,, \quad s\nu \neq s1\,; \qquad N^0_{a_{\eta\nu}} = (N_a - N)\,, \tag{26}$$

and $N^0_{a_c} = N_{a_c}$ is given by $N_{a_c} = N_a$ for the whole Hilbert space. Although the $\beta \neq c, s1$ branches have no finite pseudofermion occupancy in the initial ground state, for the present densities ranges one can define the discrete momentum number values $N_{a_\beta} = N^h_\beta$ of the corresponding unoccupied bands. For the $\beta \neq c, s1$ branches, those are the numbers $N^0_{a_{\eta\nu}}$ and $N^0_{a_{s\nu}}$ given in Eq. (26). Thus, for $\beta \neq c, s1$ branches with finite pseudofermion occupancy in the virtual state the discrete bare-momentum shifts given in Eq. (16) and deviations provided in Eq. (25) are relative to the values of such unoccupied bands.

In addition and following the change in the number of discrete bare-momentum values, this excitation also involves the pseudofermion creation and annihilation processes and pseudofermion particle-hole processes associated with PS excited states. The excitation momentum of such a ground-state - virtual-state transition reads,

$$\begin{aligned}
\delta P_{l_o,l_\triangle} &= \sum_{j=1}^{N_a} q_j\,\delta N_c(q_j) + \sum_{\nu=1}^{\infty}\sum_{j'=1}^{N_{a_{s\nu}}} q_j\,\delta N_{s\nu}(q_j) + \sum_{\nu=1}^{\infty}\sum_{j=1}^{N_{a_{\eta\nu}}} [\pi - q_j]\,N_{\eta\nu}(q_j) + \pi\,M_{\eta,-1/2}\,, \\
M_{\eta,-1/2} &= M^{un}_{\eta,-1/2} + \sum_{\nu=1}^{\infty}\nu N_{\eta\nu}\,; \quad \delta N_{s\nu}(q_j) = N_{s\nu}(q_j) \text{ for } s\nu \neq s1\,.
\end{aligned} \tag{27}$$

Here $M^{un}_{\eta,-1/2}$ is the η-spin-projection $-1/2$ unbound η-spinon number given in Eq. (3).

Since the momentum spectrum is of first order in the β momentum distribution function deviations, it is convenient to express it in terms of the corresponding occupancies of the β band bare momentum values q_j, rather than of those of the β band canonical momentum values $\bar{q}_j = \bar{q}(q_j)$, as given here. On the other hand, the excitation energy is that provided in Eq. (17).

In this first scatter-less step the pseudofermions acquire the excitation momentum and energy needed for the second and third steps.

C. Pseudofermion scattering processes, dressed S matrices, and phase shifts

In order to study the second and third processes of the ground-state - excited-energy-eigenstate transition, it is useful to express the many-pseudofermion states and operators in terms of one-pseudofermion states and operators, respectively. The PS energy and momentum eigenstates can be written as direct products of states. Those are generated by the occupancy configurations of each of the $\alpha\nu$ branches with finite pseudofermion occupancy. Moreover, the many-pseudofermion states generated by occupancy configurations of each $\alpha\nu$ branch can be expressed as a direct product of N_{a_β} one-pseudofermion states. Each of the latter states refers to one discrete bare-momentum value q_j, where $j = 1, ..., N_{a_\beta}$.

For the $N_a \gg 1$ pseudofermion description only momentum and energy contributions of order zero and one in $1/N_a$ are physically meaningful. Alike in the remaining of this paper, our analysis refers to periodic boundary conditions and large values of N_a such that $N_a \gg 1$. The Hamiltonian of the quantum problem described by the 1D Hubbard model in the PS, whose energy spectrum is given in Eq. (17), has a uniquely defined expression of the general form,

$$:\hat{H} := \sum_{\beta}\sum_{j=1}^{N_{a_\beta}} \hat{H}_{\beta,\bar{q}_j} + \sum_{\beta}\hat{H}_{\alpha}\,. \tag{28}$$

Here we have denoted the ground-state normal ordered Hamiltonian by $:\hat{H}:$, $\hat{H}_{\beta,\bar{q}_j}$ is the one-pseudofermion Hamiltonian associated with excited-state β pseudofermion or β pseudofermion hole of canonical momentum $\bar{q}_j$, and $\hat{H}_\alpha$ refers to the unbound η-spinons ($\alpha = \eta$) and unbound spinons ($\alpha = s$), which are found to be scatter-less objects. Alike the corresponding energy spectrum of Eq. (17), the ground-state normal ordered Hamiltonian is expressed as a sum of β-pseudofermion Hamiltonian terms $\sum_{j=1}^{N_{a_\beta}} \hat{H}_{\beta,\bar{q}_j}$, each of which involves a sum over the excited-state β band canonical momentum values $\bar{q}_j$.

For each many-pseudofermion PS virtual state reached in the first step of the transition from the ground state to the excited energy eigenstate, the number of active Hamiltonian terms, $\hat{H}_{\beta,\bar{q}_j}$, equals that of one-pseudofermion states of the virtual state. This number reads,

$$N_{a_c} + N_{a_{s1}} + \sum_{\beta\neq c,s1} \theta(|\delta N_\beta|)\, N_{a_\beta} \, . \tag{29}$$

Here $\theta(x) = 1$ for $x > 0$ and $\theta(x) = 0$ for $x = 0$. The numbers $N_{a_c} = N_c + N_c^h = N_a$, $N_{a_{s1}} = N_{s1} + N_{s1}^h$, and $N_{a_\beta} = N_\beta + N_\beta^h$ refer to the virtual state and corresponding excited energy eigenstate under consideration. (For the $\beta = \alpha\nu$ branches the pseudofermion-hole numbers of these expressions are provided in Eq. (A11) of Appendix A.)

The second scatter-less process generates the "in" state. The one-pseudofermion states belonging to the many-pseudofermion "in" state are the "in" asymptote states of the pseudofermion scattering theory. The generator of the virtual-state - "in"-state transition is of the form,

$$\hat{S}^0 = \prod_\beta \prod_{j=1}^{N_{a_\beta}} \hat{S}^0_\beta(q_j)\, ; \qquad \hat{S}^0_\beta(q_j) = e^{\sum_{j=1}^{N_{a_\beta}} f^\dagger_{q_j - Q^0_\beta/N_a,\beta} f_{q_j,\beta}} \, , \tag{30}$$

where $f^\dagger_{q_j,\beta}$ and $f_{q_j,\beta}$ are the β fermion operators considered in the first paper. The unitary one-pseudofermion operator $\hat{S}^0_\beta(q_j)$ is labelled by the virtual-state bare momentum q_j. Application of that unitary one-pseudofermion operator onto the corresponding one-pseudofermion state of the many-pseudofermion virtual state, shifts its virtual-state discrete bare-momentum value q_j to the excited-state discrete bare-momentum value $q_j + Q^0_\beta/N_a$, where Q^0_β is given in Eq. (16). (Note that Q^0_β may vanish, as given in that equation.)

Finally, the third step consists of a set of two-pseudofermion scattering events. It corresponds to the "in"-state - "out"-state transition, where the latter state is the PS excited energy eigenstate under consideration. The generator of that transition is the following operator,

$$\hat{S}^\Phi = \prod_\beta \prod_{j=1}^{N_{a_\beta}} \hat{S}^\Phi_\beta(q_j)\, ; \qquad \hat{S}^\Phi_\beta(q_j) = e^{\sum_{j=1}^{N_{a_\beta}} f^\dagger_{q_j - Q^\Phi_\beta(q_j)/N_a,\beta} f_{q_j,\beta}} \, . \tag{31}$$

For a PS "in" state the eigenvalue of this operator is given by,

$$S^\Phi_T = e^{i2\delta^\Phi_T} = \prod_\beta \prod_{j=1}^{N_{a_\beta}} S^\Phi_\beta(q_j)\, ; \qquad \delta^\Phi_T = \sum_\beta \sum_{j=1}^{N_{a_\beta}} Q^\Phi_\beta(q_j)/2 \, . \tag{32}$$

The one-pseudofermion states belonging to the many-pseudofermion "out" state are the "out" asymptote pseudofermion scattering states. Application of the unitary operator $\hat{S}^\Phi_\beta(q_j)$ given in Eq. (31) onto the corresponding one-pseudofermion state of the many-pseudofermion "in" state, shifts its discrete bare-momentum value $q_j + Q^0_\beta/N_a$ to the "out"-state discrete canonical-momentum value $q_j + Q_\beta(q_j)/L$. It follows that the generator of the overall virtual-state - "out"-state transition is the unitary operator,

$$\hat{S}_T \equiv \hat{S}^\Phi \hat{S}^0 = \prod_\beta \prod_{j=1}^{N_{a_\beta}} \hat{S}_\beta(q_j)\, ; \qquad \hat{S}_\beta(q_j) = \hat{S}^\Phi_\beta(q_j)\hat{S}^0_\beta(q_j) = e^{\sum_{j=1}^{N_{a_\beta}} f^\dagger_{q_j - Q_\beta(q_j)/N_a,\beta} f_{q_j,\beta}} \, . \tag{33}$$

Application of the one-β-pseudofermion or one-β-pseudofermion-hole unitary operator $\hat{S}_\beta(q_j)$ onto the corresponding one-pseudofermion state of the many-pseudofermion virtual state shifts its discrete bare-momentum value q_j directly into the "out"-state discrete canonical-momentum value $q_j + Q_\beta(q_j)/L$. The eigenvalue of the operator $\hat{S}_T$ reads,

$$S_T = e^{i2\delta_T} = \prod_\beta \prod_{j=1}^{N_{a_\beta}} S_\beta(q_j)\, ; \qquad \delta_T = \sum_\beta \sum_{j=1}^{N_{a_\beta}} Q_\beta(q_j)/2 \, . \tag{34}$$

The "in" state and "out" state are different representations of the same PS excited energy eigenstate. Specifically, they refer to the alternative β fermion and β pseudofermion representations of that state. Thus, that the one-pseudofermion states of the many-pseudofermion "in" state and "out" state are the "in" and "out" asymptote pseudofermion scattering states, respectively, implies that the one-pseudofermion Hamiltonian $\hat{H}_{\beta,\bar{q}_j}$ plays the role of the unperturbed Hamiltonian $\hat{H}_0$ of the spin-less one-particle nonrelativistic scattering theory [45]. The scattering unitary operator $\hat{S}^{\Phi}_{\beta}(q_j)$ commutes with the Hamiltonian $\hat{H}_{\beta,\bar{q}_j}$. Thus the one-pseudofermion "in" and "out" asymptote scattering states are energy eigenstates of both $\hat{H}_{\beta,\bar{q}_j}$ and $\hat{S}^{\Phi}_{\beta}(q_j)$. It follows that the matrix elements between one-pseudofermion states of $\hat{S}^{\Phi}_{\beta}(q_j)$ are diagonal. Therefore, these operators are fully defined by the set of their eigenvalues of such states. The same applies to the generator $\hat{S}^{\Phi}$ given in Eq. (31). The matrix elements of that generator between many-pseudofermion "in" states are also diagonal and thus it is fully defined by the set of its eigenvalues of such states.

Consistent, the "in" state and "out" state only differ by a mere overall phase factor. That the many-pseudofermion "in" and "out" states, which are a direct product of one-pseudofermion "in" and "out" asymptote pseudofermion scattering states, respectively, are PS excited energy eigenstates plays a major role in the pseudofermion scattering theory.

Since the operators $\hat{S}^{\Phi}_{\beta}(q_j)$ and $\hat{S}_{\beta}(q_j)$ are unitary, each of their eigenvalues has modulus one and can be written as the exponent of a purely imaginary number given by,

$$\begin{aligned} S^{\Phi}_{\beta}(q_j) &= e^{iQ^{\Phi}_{\beta}(q_j)} = \prod_{\beta'}\prod_{j'=1}^{N_{a_{\beta'}}} S_{\beta,\beta'}(q_j,q_{j'})\,; \quad j=1,...,N_{a_{\beta}}\,, \\ S_{\beta}(q_j) &= e^{iQ_{\beta}(q_j)} = e^{i\,Q^0_{\beta}} S^{\Phi}_{\beta}(q_j) = e^{i\,Q^0_{\beta}}\prod_{\beta'}\prod_{j'=1}^{N_{a_{\beta'}}} S_{\beta,\beta'}(q_j,q_{j'})\,; \quad j=1,...,N_{a_{\beta}}\,. \end{aligned} \tag{35}$$

Here $Q^{\Phi}_{\beta}(q_j)$ and $Q_{\beta}(q_j)$ are the functionals defined by Eqs. (11) and (15), respectively. By use of the former functional we find that,

$$S_{\beta,\beta'}(q_j,q_{j'}) = e^{i2\pi\,\Phi_{\beta,\beta'}(q_j,q_{j'})\,\delta N_{\beta'}(q_{j'})}\,, \tag{36}$$

where the functions $\pi\,\Phi_{\beta,\beta'}(q_j,q_{j'})$ are given in Eq. (14).

Except for the occupancy configuration changes produced by the ground-state - virtual-state transition, the only effect of, under a ground-state - excited-energy-eigenstate transition, moving the β pseudofermion or β pseudofermion hole of initial virtual-state canonical-momentum $\bar{q}_j = q_j$ once around the length L lattice ring is that its wave function acquires the overall phase factor $S_{\beta}(q_j)$ given in Eq. (35). This property follows from the form of the energy spectrum of the β pseudofermions. In contrast to that of the corresponding β fermions, it has no residual interaction terms. The procedure of moving the β pseudofermion or β pseudofermion hole once around the length L lattice ring refers to a method to derive the corresponding dressed S matrix. It is precisely the overall phase factor $S_{\beta}(q_j)$, Eq. (35), acquired by its wave function.

The phase factor $S_{\beta,\beta'}(q_j,q_{j'})$, Eq. (36), in the wave function of the β pseudofermion or β pseudofermion hole of bare-momentum q_j results from an elementary two-pseudofermion zero-momentum forward-scattering event whose scattering center is a β' pseudofermion ($\delta N_{\beta'}(q_{j'}) = 1$) or β' pseudofermion hole ($\delta N_{\beta'}(q_{j'}) = -1$) created under the ground-state - excited-state transition. The third step of that transition involves a well-defined set of elementary two-pseudofermion scattering events where all β pseudofermions and β pseudofermion holes of bare-momentum $q_j + Q^0_{\beta}/N_a$ of the "in" state are the scatterers. This leads to the overall scattering phase factor $S^{\Phi}_{\beta}(q_j)$ in their wave function given in Eq. (35). The scattering centers are the β' pseudofermions or β' pseudofermion holes of bare momentum $q_{j'} + Q^0_{\beta}/N_a$ created under the ground-state - "in"-state transition. This is confirmed by noting that $S_{\beta,\beta'}(q_j,q_{j'}) = 1$ for $\delta N_{\beta'}(q_{j'}) = 0$. Thus, out of the scatterers whose number equals that of the one-pseudofermion states given in Eq. (29), the scattering centers are only those whose bare-momentum distribution-function deviation is finite. However, out of the above scatterers, only a subclass of scatterers significantly contributes to the spectral properties [7].

The following properties play an important role in the pseudofermion scattering theory:

1. The elementary two-pseudofermion scattering processes associated with the phase factors, Eq. (36), conserve the total energy and total momentum. This stems from the occurrence of an infinite number of conservation laws [19–21] associated with the model integrability [1–4] being explicit in the present pseudofermion scattering theory. Its scatterers and scattering centers only undergo indeed zero-momentum forward scattering.

2. That the elementary two-pseudofermion scattering processes are of zero-momentum forward-scattering type also implies that they conserve the individual "in" asymptote β pseudofermion momentum value $q_j + Q^0_{\beta}/N_a$ and energy.

3. These processes also conserve the β branch, usually called *channel* in the scattering language [45].
4. The scattering amplitude does not connect quantum objects with different η spin or spin. (All such objects are η-spin neutral, spin neutral, or η-spin less and spin less anyway.)
5. For each β pseudofermion or β pseudofermion hole of virtual-state bare-momentum q_j, the dressed S matrix associated with the ground-state - excited-energy-eigenstate transition is simply the phase factor $S_\beta(q_j)$ given in Eq. (35).

The one-β-pseudofermion or one-β-pseudofermion-hole phase factor $S^{\Phi}_{\beta}(q_j)$ of the present 1D quantum problem corresponds to the usual one-particle phase factor $s_l(E)$ of similar three-dimensional quantum problems. The latter depends on the energy E and angular-momentum quantum numbers l and m. (See, for example, Eq. (6.9) of Ref. [45].) In the present 1D case, the energy E and the quantum numbers l and m are replaced by the bare-momentum q_j in the phase factor $S^{\Phi}_{\beta}(q_j)$. The β pseudofermion or β-pseudofermion hole energy is uniquely defined by the absolute bare-momentum value $|q_j|$. In 1D the sign of q_j corresponds to the three-dimensional angular-momentum quantum numbers. Another difference is that $s_l(E)$ is associated with a single scattering event. Here, $S^{\Phi}_{\beta}(q_j)$ results in general from several scattering events. Each of such events corresponds to a well defined factor $S_{\beta,\beta'}(q_j, q_{j'})$, Eq. (36), in the $S^{\Phi}_{\beta}(q_j)$ expression given in Eq. (35). There are as many of such factors as β' pseudofermion and β' pseudofermion hole scattering centers created under the transition to the virtual state and corresponding excited energy eigenstate under consideration. The factor 2 in the phase factor of Eq. (6.9) of Ref. [45] corresponds to the phase-shift definition of the standard nonrelativistic scattering theory for spin-less particles. We use in general here such a definition. It introduces the overall scattering phase shift $\delta^{\Phi}_{\beta}(q_j) = Q^{\Phi}_{\beta}(q_j)/2$ and overall phase shift $\delta_\beta(q_j) = Q_\beta(q_j)/2$. However, if instead we insert a factor 1, we would have an overall scattering phase shift $Q^{\Phi}_{\beta}(q_j)$ and overall phase shift $Q_\beta(q_j)$. (That is the phase-shift definition used in Refs. [40–43].)

The factorization of the BA bare S matrix for the original spin-1/2 electrons is associated with the so called Yang-Baxter equation (YBE) [40]. On the other hand, the factorization of the β pseudofermion or β pseudofermion-hole dressed S matrix $S_\beta(q_j)$ given in Eq. (35), in terms of the elementary two-pseudofermion S matrices $S_{\beta,\beta'}(q_j, q_{j'})$, Eq. (36), is *commutative*. Such commutativity is stronger than the symmetry associated with the YBE. This results from the c, $\eta\nu$, and $s\nu$ pseudofermions, which are the scatterers and scattering centers, being neutral objects. The $\beta = c$ fermions are both η-spin-less and spin-less objects. The composite $\eta\nu$ pseudofermions (and $s\nu$ pseudofermions) refer to η-spin-singlet (and spin-singlet) neutral configurations of 2ν anti-bound η-spin-1/2 η-spinons (and 2ν bound spin-1/2 spinons).

On the other hand, the momentum value of a unbound $\pm1/2$ η-spinon (and a unbound $\pm1/2$ spinon) was found in the first paper to be $q_{\eta,+1/2} = 0$ or $q_{\eta,-1/2} = \pi$ (and $q_{s,\pm1/2} = 0$) for all energy eigenstates with finite occupancy of such objects. That such a momentum value remains unchanged under the transition from any “in” state to the corresponding “out” state (excited energy eigenstate), confirms that such objects are not scatterers. Neither do the β pseudofermions scatterer off on them, so that they are not scattering centers. That is consistent with the unbound η-spinons (and unbound spinons) playing the passive role of unoccupied sites of both the η-spin (and spin) effective lattice and the corresponding $\eta\nu$ (and $s\nu$) effective lattices. Indeed, the objects that move around in the η-spin (and spin) effective lattice ring of length $L = N_{a_\eta}\, a_\eta = N_a\, a$ (and $L = N_{a_s}\, a_s = N_a\, a$) are the $\eta\nu$ pseudofermions (and $s\nu$ pseudofermions).

Moreover, the holes in the corresponding BA $\alpha\nu$ momentum bands exist in their own right and must not be treated as being the $M^{un}_{\alpha} = 2S_\alpha$ unbound η-spinons ($\alpha = \eta$) and ($\alpha = s$) unbound spinons and $\sum_{\nu'=\nu+1}^{\infty} 2(\nu' - \nu)N_{\alpha\nu'}$ bound η-spinons ($\alpha = \eta$) and ($\alpha = s$) bound spinons that play the role of unoccupied sites. Consistent with the results of the first paper, the unbound η-spinons and unbound spinons have their independent occupancies and the $\alpha\nu'$ pseudofermion with $\nu' > \nu$ pairs of bound η-spinons ($\alpha = \eta$) and ($\alpha = s$) bound spinons their own independent $\alpha\nu'$ momentum bands. The $\alpha\nu$ effective lattice object motion can alternatively be described in terms of $\alpha\nu$ pseudofermions and $\alpha\nu$ fpseudoermion holes, which in general have a finite velocity. Consistent, in addition to the c pseudofermions or c pseudofermion holes, the $\alpha\nu$ pseudofermions or $\alpha\nu$ pseudofermion holes are the scatterers and scattering centers of the pseudofermion scattering theory.

The BA bare S matrix refers to spin-1/2 electrons. Thus its form reflects the spin-1/2 $SU(2)$ symmetry of its scatterers and scattering centers. That of the β pseudofermions refers to neutral particles. This in spite of the η-spinons and spinons carrying η-spin 1/2 and spin 1/2, respectively. Here the M^{bo}_{η} anti-bound η-spinons (and M^{bo}_{s} bound spinons) have η-spin 1/2 (and spin 1/2) but are anti-bound (and bound) within composite neutral objects, which play the role of scatterers and scattering centers.

The dressed S matrix commutative factorization is that consistent with the form of the β pseudofermion occupancy configurations that describe the PS excited energy eigenstates. Those are superpositions of local occupancy configurations with the same number of β pseudofermions with different positions in each configuration. Hence the number of β pseudofermions belonging to β branches with finite occupancy in the virtual state is the same for all occupancy



FIG. 1: The elementary two-pseudofermion phase shift $\pi\,\Phi_{c,c}(q,q')$ in units of π as a function of q and q' for $n = 0.59$, $m = 0$, and (a) $U/t \rightarrow 0$, (b) $U/t = 0.3$, (c) $U/t = 4.9$, and (d) $U/t = 100$. The bare-momentum values q and q' correspond to the right and left axis of the figures, respectively.

configurations. The relative position of these quantum objects is different in each such a configuration. When under a specific ground-state - excited-energy-eigenstate transition a β pseudofermion or β pseudofermion hole moves once around the lattice ring, it scatters the same β' pseudofermion or β' pseudofermion-hole scattering centers, but in different order for different occupancy configurations. However, it is required that the phase factor $e^{iQ_\beta(q_j)}$ acquired by the β pseudofermion or β pseudofermion hole should be the same, independently of that order. This is consistent with the commutativity of the dressed S-matrix factors $S_{\beta,\beta'}(q_j,q_{j'})$ in the overall dressed S matrix $S_\beta(q_j)$, whose expression is given in Eq. (35). Such a commutativity follows from the elementary S matrices $S_{\beta,\beta'}(q_j,q_{j'})$, Eq. (36), being simple phase factors, instead of matrices of dimension larger than one.

As found in the first paper for the $\alpha\nu$ fermions, the $\alpha\nu$ pseudofermions generated from corresponding $\alpha\nu$ fermions with limiting bare-momentum values $q = \pm q_{\alpha\nu}$ and vanishing velocities $v_{\alpha\nu}(\pm q_{\alpha\nu}) = 0$ are invariant under the electron - rotated-electron unitary transformation. It is found below that for the excited energy eigenstates belonging to a given PS, the limiting canonical momenta have exactly the same values as those of the corresponding initial ground state provided in Eq. (A12) of Appendix A. Moreover, at the limiting momentum values $\pm q_{\alpha\nu}$, the energies of such $\alpha\nu$ pseudofermions equals those of the corresponding $\alpha\nu$ fermions and read [5],

$$\begin{aligned}\varepsilon_{\eta\nu}(\pm q_{\eta\nu}) &= \nu\,[\varepsilon_{\eta,-1/2}+\varepsilon_{\eta,+1/2}] = (1-\delta_{n,1})\,2\nu|\mu| + \delta_{n,1}\,2\nu\mu^0\,; \quad \varepsilon^0_{\eta\nu}(\pm q_{\eta\nu}) = 0\,,\\ \varepsilon_{s\nu}(\pm q_{s\nu}) &= \nu\,[\varepsilon_{s,-1/2}+\varepsilon_{s,+1/2}] = 2\nu\mu_B\,|H|\,; \quad \varepsilon^0_{s\nu}(\pm q_{s\nu}) = 0\,. \end{aligned} \tag{37}$$



FIG. 2: The elementary two-pseudofermion phase shift $\pi\,\Phi_{c,s1}(q,q')$ in units of π as a function of q and q' for $n = 0.59$, $m = 0$, and (a) $U/t \rightarrow 0$, (b) $U/t = 0.3$, (c) $U/t = 4.9$, and (d) $U/t = 100$. The bare-momentum values q and q' correspond to the right and left axis of the figures, respectively.

Here $2\mu^0$ is the $n = 1$ Mott-Hubbard gap, which for $u \ll 1$ and $u \gg 1$ behaves as,

$$\begin{aligned} 2\mu^0 &\approx \frac{8}{\pi}\sqrt{t\,U}\,e^{-2\pi\left(\frac{t}{U}\right)}\,, \quad m = 0\,; \quad 2\mu^0 \approx \frac{U^2}{8t}\,, \quad m = -1, 1\,, \quad u \ll 1\,, \\ 2\mu^0 &\approx [U - 4t]\,, \quad m \in [-1, 1]\,, \quad u \gg 1\,. \end{aligned} \tag{38}$$

Moreover, $\varepsilon_{\eta,\pm 1/2}$ and $\varepsilon_{s,\pm 1/2}$ denote in Eq. (37) the energies of the unbound η-spinons and spinons, respectively, which were found in the first paper to be given by,

$$\begin{aligned} \varepsilon_{\eta,\pm 1/2} &= 2|\mu|\,; \quad \varepsilon_{\eta,\mp 1/2} = 0\,, \quad \mathrm{sgn}\{(1-n)\}1 = \mp 1\,, \\ \varepsilon_{s,\pm 1/2} &= 2\mu_B\,|H|\,; \quad \varepsilon_{s,\mp 1/2} = 0\,, \quad \mathrm{sgn}\{m\}1 = \mp 1\,. \end{aligned} \tag{39}$$

The effects of such an additional pseudofermion symmetry are discussed below in Section IV A.

III. THE PSEUDOFERMION PHASE SHIFTS

In this section we study the β pseudofermion phase shifts associated with the dressed S matrix introduced above.

A. Phase-shift definition

As above, our analysis refers to periodic boundary conditions and $N_a \gg 1$. The c effective lattice considered in the first paper equals the original lattice. On the other hand, the spacing,

$$a_{\alpha\nu} = \frac{L}{N_{a_{\alpha\nu}}} = \frac{N_a}{N_{a_{\alpha\nu}}}\, a\,; \quad N_{a_{\alpha\nu}} \geq 1\,, \tag{40}$$

of the remaining $\beta = \alpha\nu$ effective lattices also considered in the first paper is for $n \neq 1$ and $m \neq 0$ larger than that of the original lattice. Its definition, Eq. (40), is such that the corresponding length, $L = N_a\, a = N_{a_\beta}\, a_\beta$, equals that the latter lattice. (In general we use in this paper units of lattice constant a, for which $N_a = L$.)

Depending on the asymptote coordinate choice, there are two possible definitions for the β pseudofermion phase shifts associated with the dressed S matrix $S_\beta(q_j)$ given in Eq. (35). The uniquely defined quantity is the dressed S matrix. The two choices of asymptote coordinates for the β pseudofermion or β pseudofermion hole correspond to $x \in [-L/2, +L/2]$ and $x \in [0, +L]$, respectively.

If when moving around the lattice ring the β pseudofermion (or β pseudofermion hole) departures from the point $x = -L/2$ and arrives to $x = L/2$, one finds that,

$$\lim_{x\to L/2} \bar{q}\, x = q\, L/2 + Q^0_\beta/2 + Q^\Phi_\beta(q)/2 = q\, L/2 + \delta_\beta(q)\,, \tag{41}$$

where

$$\delta_\beta(q) = Q_\beta(q)/2\,. \tag{42}$$

For this asymptote coordinate choice, $\delta_\beta(q)$ is the overall β pseudofermion or β pseudofermion-hole phase shift. Moreover, from analysis of Eqs. (11) and (15) it follows that $\pi\, \Phi_{\beta,\beta'}(q_j, q_{j'})$ is an elementary two-pseudofermion phase shift. This phase-shift definition corresponds to that used in standard quantum non-relativistic scattering theory [45]. It is such that the dressed S matrix, $S_\beta(q_j)$, given in Eq. (35) can be written as,

$$S_\beta(q_j) = e^{i2\delta_\beta(q_j)}\,, \qquad j = 1, ..., N_{a_\beta}\,. \tag{43}$$

The factor 2 appearing in the exponential argument of Eq. (43) corresponds to the usual form of the S matrix for that theory. This phase-shift definition is consistent with an exact Theorem due to Fumi [46]. For the phase-shift definition of Eq. (42), $Q^0_\beta/2 = 0, \mp\pi/2$ corresponds to the scatter-less term $-l\pi/2$ of the three-dimensional partial-wave problem of orbital angular momentum l [45, 46]. Although the orbital angular momentum vanishes in 1D, the scatter-less phase shift given in Eq. (16) plays a similar role. Here we follow the definition of the standard quantum non-relativistic scattering theory and choose the overall β pseudofermion phase shift definition, $Q_\beta(q)/2$, associated with Eq. (41).

On the other hand, if when moving around the lattice ring the β pseudofermion (or β pseudofermion hole) departures from the point $x = 0$ and arrives to $x = L$, one finds that,

$$\lim_{x\to L} \bar{q}\, x = q\, L + Q^0_\beta + Q^\Phi_\beta(q) = q\, L + Q_\beta(q)\,, \tag{44}$$

where q refers to the initial ground state. For this asymptote coordinate choice, $Q_\beta(q)$ is the overall β pseudofermion (or β pseudofermion hole) phase shift and $2\pi\, \Phi_{\beta,\beta'}(q_j, q_{j'})$ is an elementary two-pseudofermion phase shift.

The overall pseudofermion phase-shift choice, $Q_\beta(q) = Q^0_\beta + Q^\Phi_\beta(q)$, is associated with the asymptote condition of Eq. (44). It corresponds to a generalization of the conventional phase-shift definition previously used in the BA literature for both an initial $n = 1$ and $m = 0$ initial ground state [41, 42] and an initial ground state with electronic density in the range $n \in (0, 1)$ and zero spin density [40]. (All the discussions and analysis presented below in this paper also apply to the phase-shift definition, $Q_\beta(q) = Q^0_\beta + Q^\Phi_\beta(q)$, provided that the β phase shifts $\delta_\beta(q) = Q_\beta(q)/2$ are multiplied by two.)

The phase shifts of Refs. [47, 48] correspond to a particular case of the general elementary two-pseudofermion phase shifts considered in this paper. If one considers densities in the ranges $n \in [0, 1[$ and $m \in]0, n]$ and the specific PS subspace spanned by energy eigenstates such that $N_\beta = 0$ for the $\beta \neq c, s1$ branches and $M^{un}_{\alpha,-1/2} = 0$ for $\alpha = \eta, s$, the general integral equations (B1)-(B15) of Appendix B reduce to the integral equations (20)-(23) of Ref. [47] and (32)-(38) of Ref. [48].

An important property specific to the pseudofermion scattering theory is:



FIG. 3: The elementary two-pseudofermion phase shift $\pi\,\Phi_{s1,c}(q,q')$ in units of π as a function of q and q' for $n = 0.59$, $m = 0$, and (a) $U/t \rightarrow 0$, (b) $U/t = 0.3$, (c) $U/t = 4.9$, and (d) $U/t = 100$. The bare-momentum values q and q' correspond to the right and left axis of the figures, respectively.

The phase shift $\delta_{\beta}(q_j) = Q_{\beta}(q_j)/2 = Q^0_{\beta}/2 + Q^{\Phi}_{\beta}(q_j)/2$ defined in Eqs. (15) and (42) where $Q^{\Phi}_{\beta}(q_j)/2$ and $Q^0_{\beta}/2$ are the scattering phase shift expressed in Eq. (11) in terms of a suitable superposition of two-pseudofermion phase shifts $\pi\,\Phi_{\beta,\beta'}(q_j,q_{j'})$ uniquely defined for each excited state and the BA phase shift of Eq. (16), respectively, refers to all $j = 1,...,N_{a_{\beta}}$ β-band scatterers of the excited state. This includes both those created under the transition from the ground state to the excited state and the scatterers that pre-existed in the ground state.

This property does not hold for the traditional holon and spinon scattering theories of Refs. [40–42], whose phase shifts refer only to the scatterers created under the transition from the ground state to the excited states. Those are the scatterers that play as well the role of scattering centers. As discussed below in Section VI, this prevents such theories of describing the model dynamical and spectral properties. Indeed, such properties are found to be controlled by the phase shifts of both the scatterers created under the transition from the ground state to the excited states and pre-existing in the ground state.

We recall that the above $N_{a_{\beta}}$ β-band scatterers, include the N_{β} β pseudofermions and N^h_{β} β pseudofermion holes that populate the excited state, where $N_{a_{\beta}} = [N_{\beta} + N^h_{\beta}]$, as given in Eq. (A10) of Appendix A for $\beta = c, \alpha\nu$. For initial ground states with densities in the ranges $n \in [0,1]$ and $m \in [0,n]$ such numbers are for instance for the $\beta = c, s1$ bands of the PS excited states given by $N_c = [N + \delta N_c]$, $N^h_c = [N_a - N - \delta N_c]$, $N_{s1} = [N_{\uparrow} + \delta N_{s1}]$, and $N^h_{s1} = [N_{\uparrow} - N_{\downarrow} + \delta N^h_{s1}]$ where $\delta N_c/N_a \rightarrow 0$, $\delta N_{s1}/N_a \rightarrow 0$, and $\delta N^h_{s1}/N_a \rightarrow 0$ as $N_a \rightarrow \infty$.

Thus the pseudofermion scattering theory provides the phase shifts of a macroscopic number N_a of c band scatterers and $N+\delta N_{a_{s1}}$ of $s1$ band scatterers, which result from their scattering with the scattering centers emerging in one or several β' bands under the transition from the ground state to the excited state. The number of created β' scattering centers under a transition from the ground state to a excited state is much smaller than that of the theory scatterers. It given by,

$$N_{centers} = \sum_{\beta'} \sum_{j'=1}^{N_{a_{\beta'}}} |\delta N_{\beta'}(q_{j'})| \, , \tag{45}$$

where the deviations $\delta N_{\beta'}(q_{j'})$ are those on the right-hand side of the scattering phase-shift expression given in Eq. (11). They refer to the creation of one $\beta' = c, \alpha\nu$ pseudofermion scattering center of momentum $q_{j'}$ for $\delta N_{\beta'}(q_{j'}) = 1$ and one $\beta' = c, s1$ pseudofermion-hole scattering center of momentum $q_{j'}$ for $\delta N_{\beta'}(q_{j'}) = -1$. All $N_{a_\beta} = [N_\beta + N^h_\beta]$ β-band scatterers acquire a phase shift $\pi\,\Phi_{\beta,\beta'}(q_j, q_{j'})$ due to their interaction with each created β' scattering center, as confirmed by the form of the scattering phase shift $Q^\Phi_\beta(q_j)/2$ given in Eq. (11).

B. The two-pseudofermion phase shifts: Bare-momentum dependence and Levinson's Theorem

The bare-momentum two-pseudofermion phase shifts, $\pi\,\Phi_{\beta,\beta'}(q, q')$, are related to the rapidity two-pseudofermion phase shifts, $\pi\bar{\Phi}_{\beta,\beta'}(r, r')$, by Eq. (14). The latter phase shifts are defined by the integral equations (B1)-(B15) of Appendix B. In that Appendix we provide the general equations that define the rapidity elementary two-pseudofermion phase shifts, $\pi\bar{\Phi}_{\beta,\beta'}(r, r')$. In addition, in Appendix B corresponding simplified equations specific to the $m \to 0$ limit, closed-form expressions valid in the $m \to 0, n \to 1$ limit, and analytical expressions valid for $U/t \to 0$ as $m \to 0$ are provided.

In Figs. 1-6 the two-pseudofermion phase shifts in the expressions of the overall phase shifts associated with the excited states that mostly contribute to the spectral weights are plotted. The plots are as a function of the bare-momenta q and q' for electronic density $n = 0.59$, spin density $m \to 0$, and $U/t \to 0$ and $U/t = 0.3, 4.9$,100. (The electronic density $n = 0.59$ and the value $U/t = 4.9$ are those used in Refs. [9, 23, 24] for the description of the TCNQ photoemission dispersions observed in the quasi-1D organic compound TTF-TCNQ.)

The two-pseudofermion phase shift $\pi\,\Phi_{c,\eta 1}(q, q')$ plotted in Fig. 5 in units of π has values in the domain $\Phi_{c,\eta 1}(q, q') \in [-1, 1]$. Within the standard quantum non-relativistic scattering theory phase shift definition given in Eq. (41), this corresponds to the phase-shift range $\pi\,\Phi_{c,\eta 1}(q, q') \in [-\pi, \pi]$. Note that the width of this domain is 2π, whereas the definition of Eq. (44) would lead to a domain of width 4π.

The phase shifts $\pi\,\Phi_{\beta,\beta'}(q, q')$ fully control the expression of the f functions given in Eq. (20). Moreover, analysis of the expressions provided in Eqs. (A20) and (A21) of Appendix A reveals that also the β energy dispersions involve such phase shifts. Specifically, according to the expressions provided in Eq. (A20) of Appendix A the energy associated with the creation of a β pseudofermion has a contribution from the phase shifts felt by all ground-state c Fermi sea c pseudofermions, as a result of its creation. Consistent, the $\int_{-Q}^{Q} dk$ integration in such expressions may be written in terms of a corresponding bare-momentum integration $\int_{-q_{Fc}}^{q_{Fc}} dq' = \int_{-2k_F}^{2k_F} dq'$ and alternatively of a sum $\sum_{j'=1}^{N_a} N^0_c(q_{j'})$. Here $N^0_c(q_{j'}) = 0$ and $N^0_c(q_{j'}) = 1$ for $|q_{j'}| > 2k_F$ and $|q_{j'}| < 2k_F$, respectively. Under such a variable transformation, $\bar{\Phi}_{c,\beta}$ is replaced by a bare-momentum two-pseudofermion phase shift in units of π, $\Phi_{c,\beta}(q', q) = \Phi_{c,\beta}(q_{j'}, q_j)$, Eq. (14). In it the c pseudofermion and the β pseudofermion are the scatterer and the scattering center, respectively. (Here we have used the notation $\Phi_{c,\beta}(q_{j'}, q_j)$ rather than our usual notation $\Phi_{c,\beta}(q_j, q_{j'})$. That choice is that consistent with the argument of the corresponding energy dispersion $\varepsilon^0_\beta(q_j)$ in Eq. (A20) of Appendix A being denoted by q_j.)

The properties of the β pseudofermions are consistent with the absence of β pseudofermions bound states. Here we confirm that the corresponding phase shifts $\pi\,\Phi_{\beta,\beta'}(q, q')$ associated with the elementary two-pseudofermion scattering events obey Levinson's Theorem [49] provided that there are no such bound states. Such a theorem states that when in the reference frame of the scattering center the momentum of the scatterer tends to zero the phase shift is given by πN_b, where N_b is the number of bound states. In that frame, the phase shift $\pi\,\Phi_{\beta,\beta'}(q, q')$ reads $\pi\,\Phi_{\beta,\beta'}(q - q', 0)$.

In our case the, BA-solution $\alpha\nu$ momentum bands occupancy configurations are consistent with the absence of β pseudofermion bound states. On the other hand, Levinson's Theorem must hold for the present scattering theory. The two requirements of absence of β pseudofermion bound states and validity of Levinson's Theorem are fulfilled provided that the two-pseudofermion phase shifts of the present theory obey the following equation,

$$\lim_{q-q' \to 0} \pi\,\Phi_{\beta,\beta'}(q - q', 0) = 0 \, . \tag{46}$$



FIG. 4: The elementary two-pseudofermion phase shift $\pi\,\Phi_{s1,s1}(q,q')$ in units of π as a function of q and q' for $n = 0.59$, $m = 0$, and (a) $U/t \rightarrow 0$, (b) $U/t = 0.3$, (c) $U/t = 4.9$, and (d) $U/t = 100$. The bare-momentum values q and q' correspond to the right and left axis of the figures, respectively.

To check whether this equation is obeyed, after a straightforward algebra involving the integral equations (B1)-(B15) of Appendix B, we have found that $\pi\,\bar{\Phi}_{\beta,\beta'}\,(r,r') = -\pi\,\bar{\Phi}_{\beta,\beta'}\,(-r,-r')$. This result combined with the use of Eq. (14) and the odd character of the ground-state rapidity functions, $\Lambda^0_\beta(q) = -\Lambda^0_\beta(-q)$ [7], then leads to,

$$\pi\,\Phi_{\beta,\beta'}(q,q') = -\pi\,\Phi_{\beta,\beta'}(-q,-q')\,. \tag{47}$$

This latter symmetry implies that $\pi\,\Phi_{\beta,\beta'}(q-q',0)$ is an odd function of $q-q'$. This confirms both the validity of Levinson's Theorem, Eq. (46), and the absence of β pseudofermion bound states.

IV. EFFECTS OF THE PSEUDOFERMION INVARIANCE UNDER THE ELECTRON - ROTATED-ELECTRON UNITARY TRANSFORMATION ON THEIR SCATTERING PROPERTIES

The β pseudofermions have the same transformation laws under the electron - rotated-electron unitary transformation as the corresponding β fermions. For the latter objects such transformation laws and corresponding invariances were studied in the first paper.

Here we address the issue of the effects of the pseudofermion invariance under the electron - rotated-electron unitary transformation on their scattering properties. We consider two different cases of PSs whose initial ground states refer to (i) densities in the ranges obeying the inequalities $0 < n < 1$ and $0 < m < n$ and (ii) densities $n = 1$ and/or $m = 0$. The effects of the pseudofermion invariance under the electron - rotated-electron unitary transformation are different for these two PS types.

A. Effects of the pseudofermion invariances on their scattering properties for densities in the ranges obeying the inequalities $0 < n < 1$ and $0 < m < n$

The β pseudofermions are not in general invariant under the electron - rotated-electron unitary transformation. The exception is as the canonical momentum $\bar{q}$ of the composite $\beta = \alpha\nu$ pseudofermions approaches its limiting values, $\bar{q} \rightarrow \pm q_{\alpha\nu}$, and their energies obey Eq. (37). As discussed in the first paper for the composite $\alpha\nu$ fermions, this invariance occurs provided that the corresponding velocity, $v_{\alpha\nu}(q) = \partial\varepsilon^0_{\alpha\nu}(q)/\partial q = \partial\varepsilon_{\alpha\nu}(q)/\partial q$, vanishes at those limiting momentum values, $v_{\alpha\nu}(\pm q_\beta) = 0$. Below it is confirmed that the functional $\bar{q}_j = \bar{q}(q_j)$ of Eq. (10) is such that $\bar{q}(\pm q_{\alpha\nu}) = \pm q_{\alpha\nu}$ provided that $v_{\alpha\nu}(\pm q_\beta) = 0$. The latter condition is met for all $\alpha\nu \neq s1$ branches in all PSs. For the $s1$ branch it is met only for PSs whose initial ground state has a small or vanishing number, N_{s1}, of $s1$ pseudofermions such that $N_{s1}/N_a \rightarrow 0$ as $N_a \rightarrow \infty$. On the other hand, all ground states have zero occupancy of $\alpha\nu \neq s1$ pseudofermions.

Here we study the role of such a symmetry in the scattering properties of the $\alpha\nu$ pseudofermions for PSs whose initial ground states have densities in the ranges $0 < n < 1$ and $0 < m < n$. First we use the phase-shift expressions involving $\alpha\nu$ pseudofermions of canonical momentum $\bar{q} = \pm q_{\alpha\nu}$ and vanishing velocity $v_{\alpha\nu}(\pm q_{\alpha\nu}) = 0$ to extract useful information about their properties. Concerning the particular case of the $s1$ branch, below we account for the condition $v_{s1}(\pm q_{s1}) = 0$ being met provided that $N_{s1}/N_a \rightarrow 0$ as $N_a \rightarrow \infty$ being equivalent to consider that $r^0_s \rightarrow 0$ in the the integral equations, Eqs. (B1)-(B3) of Appendix B, for the two-pseudofermion phase shifts $\pi\,\Phi_{s1,\beta'}(q,q')$. Next we use such properties to clarify the effects of the $s1$ pseudofermion invariance under the electron - rotated-electron unitary transformation. (In this section we use in general the notations $\alpha\nu$ and β' to denote the branch of the pseudofermion or pseudofermion-hole scatterers and scattering centers, respectively.)

Our first goal is to show that the $\alpha\nu$ pseudofermions of limiting canonical momentum $\bar{q} = \pm\bar{q}_{\alpha\nu}$ and vanishing velocity $v_{\alpha\nu}(\pm\bar{q}_{\alpha\nu}) = 0$ are not *active scatterers*. Active scatterers are those whose overall phase shifts generated by the ground-state - excited-energy-eigenstate transitions lead to a shift of the corresponding canonical-momentum values. In the following we confirm that the ground-state limiting canonical-momentum values $\bar{q} = \pm\bar{q}_{\alpha\nu}$ of such $\alpha\nu$ pseudofermions are not shifted by the ground-state - excited-energy-eigenstate transitions, so that $\bar{q}_{\alpha\nu} = q_{\alpha\nu}$ and $v_{\alpha\nu}(\pm\bar{q}_{\alpha\nu}) = v_{\alpha\nu}(\pm q_{\alpha\nu}) = 0$.

In contrast to the band limiting momentum values of usual band particles and Fermi-liquid quasi-particles, the $\alpha\nu$ band bare-momentum limiting values $\pm q_{\alpha\nu}$ given in Eq. (A9) of Appendix A of the composite $\alpha\nu$ fermions can change due to shake-up effects. Those are caused by the ground-state - excited-energy-eigenstate transitions. Since $\pm\delta q_{\alpha\nu} = \pm\pi\,\delta N_{a_{\alpha\nu}}/N_a$, such an exotic behavior occurs when the deviation $\delta N_{a_{\alpha\nu}}$, Eq. (25), generated by the ground-state - excited-energy-eigenstate transition is finite.

On the other hand, in what the $\alpha\nu$ pseudofermions is concerned, we find from the use of the integral equations, Eqs. (B1)-(B3) of Appendix B, for the two-pseudofermion phase shifts, $\pi\,\Phi_{s1,\beta'}(q,q')$, and Eqs. (B10)-(B15) of the same Appendix for the two-pseudofermion phase shifts, $\pi\,\Phi_{\alpha\nu,\beta'}(q,q')$, involving the $\alpha\nu \neq s1$ branches that the two-pseudofermion phase shifts $\pi\,\Phi_{s1,\beta'}(\iota\,q_{s1},q)$ in the $r^0_s \rightarrow 0$ limit and the two-pseudofermion phase shifts $\pi\,\Phi_{\alpha\nu,\beta'}(\iota\,q_{\alpha\nu},q)$ of the $\alpha\nu \neq s1$ branches have the following expression,

$$\pi\,\Phi_{\alpha\nu,\beta'}(\iota\,q_{\alpha\nu},q) = \frac{\iota\,\pi}{2}\Big[\delta_{\beta',c}(\delta_{\alpha,\eta} - \delta_{\alpha,s}) + \delta_{\beta',\alpha'\nu'}\delta_{\alpha,\alpha'}(-\delta_{\nu,\nu'} + \nu + \nu' - |\nu - \nu'|)\Big]\,; \quad \iota = \pm 1\,. \tag{48}$$

The $r^0_s \rightarrow 0$ limit in which this expression is valid for the $s1$ branch is reached when $N_{s1}/N_a \rightarrow 0$ as $N_a \rightarrow \infty$. As found in the first paper, this is exactly the only limit in which the condition $v_{s1}(\iota\,q_{s1}) = 0$ is fulfilled.

The two-pseudofermion phase-shift expression given in Eq. (48) is valid provided that $q \neq \iota\,q_{\alpha\nu'}$ for $\beta' = \alpha\nu'$. Its use in the general overall scattering phase-shift expression given in Eq. (11) leads to,

$$\frac{Q^\Phi_{\alpha\nu}(\iota q_{\alpha\nu})}{2} = -\frac{\iota\,\pi}{2}\Big[\delta N_{\alpha\nu} - [\delta_{\alpha,\eta} - \delta_{\alpha,s}]\delta N_c - \sum_{\nu'=1}^{\infty}(\nu + \nu' - |\nu - \nu'|)\delta N_{\alpha\nu'}\Big]\,; \quad \iota = \pm 1\,. \tag{49}$$

Comparison of this expression with that of $\iota\delta q_{\alpha\nu} = \iota\pi\,\delta N_{a_{\alpha\nu}}/N_a$ where $N_{a_{\alpha\nu}} = [N_{\alpha\nu} + N^h_{\alpha\nu}]$ and $N^h_{\alpha\nu}$ is provided in Eq. (A11) of Appendix A, confirms that $Q^\Phi_{\alpha\nu}(\iota q_{\alpha\nu})/N_a = -\iota\delta q_{\alpha\nu}$. For the $s1$ pseudofermions such results are valid in the above limit in which $v_{s1}(\iota\,q_{s1}) = 0$.

For a $\beta' = \alpha\nu'$ pseudofermion scattering center the two-pseudofermion phase shift expression provided in Eq. (48) does not apply at $q = \iota\,q_{\alpha\nu'}$. However, from the form $\Lambda_{\alpha\nu}(q) = \Lambda^0_{\alpha\nu}(\bar{q}(q))$ of the PS excited energy-egenstates rapidity



FIG. 5: The elementary two-pseudofermion phase shift $\pi\,\Phi_{c,\eta 1}(q,q')$ in units of π as a function of q and q' for $n=0.59$, $m=0$, and (a) $U/t\rightarrow 0$, (b) $U/t=0.3$, (c) $U/t=4.9$, and (d) $U/t=100$. The bare-momentum values q and q' correspond to the right and left axis of the figures, respectively.

function given in Eq. (8), where $\Lambda^0_{\alpha\nu}(q)$ is the corresponding expression for the initial ground state, one confirms that the relation $Q^{\Phi}_{\alpha\nu}(\iota q_{\alpha\nu})/N_a=\iota\delta q_{\alpha\nu}$ is valid for all PS excited states provided that $v_{\alpha\nu}(\iota\, q_{\alpha\nu})=0$. The rapidity functions of the excited energy eigenstates of a given initial ground state equal those of the latter state with in the argument of such functions the ground-state bare momentum replaced by the excited-state canonical momentum. This property implies that the corresponding bare-momentum and canonical-momentum bands have precisely the same momentum width. Hence one has that $Q^{\Phi}_{\alpha\nu}(\iota q_{\alpha\nu})/N_a=-\iota\delta q_{\alpha\nu}$ for all PS excited states provided that $v_{\alpha\nu}(\iota\, q_{\alpha\nu})=0$.

We have just confirmed that whenever $v_{\alpha\nu}(\iota\, q_{\alpha\nu})=0$ the scattering phase shift leads to a canonical-momentum shift $Q^{\Phi}_{\alpha\nu}(\pm q_{\alpha\nu})/N_a=\mp\delta q_{\alpha\nu}$ that exactly cancels the bare-momentum shift $\pm\delta q_{\alpha\nu}$. Hence the overall canonical-momentum shift $\pm\delta\bar{q}_{\alpha\nu}=\pm\delta q_{\alpha\nu}+Q^{\Phi}_{\alpha\nu}(\pm q_{\alpha\nu})/N_a$ indeed vanishes. It follows that for the $\alpha\nu$ pseudofermions of vanishing velocity $v_{\alpha\nu}(\iota\, q_{\alpha\nu})=0$ the limiting canonical momenta have the same values, $\pm q_{\alpha\nu}$, both for the ground state and PS excited energy eigenstates. As a result, at $\bar{q}=\pm q_{\alpha\nu}$ such objects are not active scatterers.

Next we investigate the properties of the $\alpha\nu$ pseudofermions of canonical momentum $\bar{q}=\pm q_{\alpha\nu}$ and vanishing velocity $v_{\alpha\nu}(\pm q_{\alpha\nu})=0$ as scattering centers. In the following we use the PS property that for energy eigenstates belonging to such a subspace the limiting momentum $q_{\alpha\nu}$ of Eq. (A9) of Appendix A equals that of the initial ground state provided in Eq. (A12) of Appendix A. By use of the integral equations, Eqs. (B1)-(B15) of that appendix, and Eqs. (A.11)-(A.14) of Ref. [8], we find after some algebra that for $\alpha'\nu'\neq s1$ branches the two-pseudofermion phase



FIG. 6: The elementary two-pseudofermion phase shift $\pi\,\Phi_{s1,\eta 1}(q,q')$ in units of π as a function of q and q' for $n=0.59$, $m=0$, and (a) $U/t\rightarrow 0$, (b) $U/t=0.3$, (c) $U/t=4.9$, and (d) $U/t=100$. The bare-momentum values q and q' correspond to the right and left axis of the figures, respectively.

shift $\pi\,\Phi_{\beta,\alpha'\nu'}(q,\iota\,q_{\alpha'\nu'})$ may be expressed as follows,

$$\begin{aligned}\pi\,\Phi_{\beta,\alpha'\nu'}(q,\iota\,q_{\alpha'\nu'}) &= \frac{\iota\,\pi}{2}\Big[\delta_{\beta,c}-\delta_{\beta,\alpha\nu}\delta_{\alpha,\alpha'}(-\delta_{\nu,\nu'}+\nu+\nu'-|\nu-\nu'|)\Big]\\ &+ \frac{\iota}{2}\sum_{\iota'=\pm1}\iota'\Big[\pi\,\Phi_{\beta,c}(q,\iota'\,2k_F)-\delta_{\alpha',s}\,2\pi\,\Phi_{\beta,s1}(q,\iota'\,k_{F\downarrow})\Big]\,;\quad \iota=\pm1\,. \end{aligned}\tag{50}$$

Again, for $\alpha'\nu'=s1$ this expression is valid only in the limit of vanishing velocity $v_{s1}(\pm q_{s1})=0$ in which $2\pi\,\Phi_{\beta,s1}(q,\iota'\,k_{F\downarrow})=2\pi\,\Phi_{\beta,s1}(q,\iota'\,0)=0$. In that expression β stands for any of the branches with finite pseudofermion occupancy for the excited state under consideration. The values of q are such that $|q|<q_{\alpha'\nu'}$ for $\beta=\alpha'\nu$. Otherwise they can have any value and thus correspond to any excited-state active β scatterer.

The form of the two-pseudofermion phase-shift expression given in Eq. (50) reveals that, except for the constant phase-shift terms, creation of one $\eta\nu'$ pseudofermion (and one $s\nu'\neq s1$ pseudofermion) at canonical momentum $q_{\eta\nu'}=\iota\,[\pi-2k_F]$ (and $q_{s\nu'}=\iota\,[k_{F\uparrow}-k_{F\downarrow}]$ for $\nu'>1$ and $q_{s1}=\iota\,2k_F$ as $k_{F\uparrow}\rightarrow 2k_F$, $k_{F\downarrow}\rightarrow 0$, and thus $v_{s1}(\iota\,q_{s1})=0$) is felt by the β pseudofermion or β pseudofermion-hole active scatterers as a c pseudofermion (and a c and $s1$ pseudofermion) Fermi-points current excitation. Such an excitation is associated with a shift, $\iota\pi/N_a$, of both c bare-momentum Fermi points (and a shift, $\iota\pi/N_a$, of both c bare-momentum Fermi points and a shift, $-\iota 2\pi/N_a$, of both $s1$ bare-momentum Fermi points). Thus, such scatterers effectively feel that they are scattered by c Fermi-point current shifts (and c and $s1$ Fermi-point current shifts), rather than by the $\eta\nu'$ (and $s\nu'$) pseudofermion created at canonical momentum $q_{\eta\nu'} = \iota\,[\pi - 2k_F]$ (and $q_{s\nu'} = \iota\,[k_{F\uparrow} - k_{F\downarrow}]$ for $\nu' > 1$ and $q_{s1} = \iota\,2k_F$ as $k_{F\uparrow} \rightarrow 2k_F$ and $k_{F\downarrow} \rightarrow 0$).

Active scattering centers are those that contribute to the scattering phase shift given in Eq. (11). For instance, small-momentum and low-energy c and $s1$ pseudofermion particle-hole processes in the vicinity of the corresponding $\beta = c, s1$ Fermi points, called elementary processes (C) in Ref. [7], do not generate active scattering centers. Within such processes the phase shifts generated by the pseudofermion particle-like excitations exactly cancel those produced by the pseudofermion hole-like excitations.

The definitions of *active* $\beta = \alpha\nu$ *scatterers* and *active* $\beta' = \alpha'\nu'$ *scattering centers* refer to different phase shits. Active $\beta = \alpha\nu$ scatterers of canonical momentum $\bar{q}_j = \bar{q}(q_j)$ (and active $\beta' = \alpha'\nu'$ scattering centers of canonical momentum $\bar{q}'_j = \bar{q}(q'_j)$) are those whose overall phase shift $Q_\beta(q_j)/2 = Q^0_\beta/2 + Q^\Phi_\beta(q_j)/2$, Eq. (15), leads to a canonical momentum value $\bar{q}_j \neq q_j$ (and those that contribute to the scattering phase shift $Q^\Phi_\beta(q_j)/2$ defined by Eq. (11)).

The part of the β' bare-momentum distribution-function deviation generated by β' active scattering centers can be written as $\delta N^{NF}_{\beta'}(q') + \delta N^F_{\beta'}(q') + \delta N^L_{s1}(q') \times \delta_{v_{s1}(q_{s1}),0}$ for the $\beta' = c, s1$ branches and $\delta N^{NF}_{\beta'}(q') + \delta N^L_{\beta'}(q')$ for the $\beta' = \alpha'\nu' \neq s1$ branches. Here $\delta N^{NF}_{\beta'}(q')$ is generated by the processes called elementary processes (A) in Ref. [7]. Those create and annihilate (and create) $\beta' = c, s1$ pseudofermions (and $\beta' = \alpha'\nu' \neq s1$ pseudofermions) away from the $\beta' = c, s1$ Fermi points and $s1$ limiting values $\pm q_{s1} = \pm 2k_F$ as $k_{F\uparrow} \rightarrow 2k_F$ and $k_{F\downarrow} \rightarrow 0$ (and from the limiting values $\pm q_{\alpha'\nu'}$). On the other hand, $\delta N^F_{\beta'}(q')$ and $\delta N^L_{\beta'}(q')$ are generated by the processes called elementary processes (B) in the same reference. Those create and annihilate (and create) $\beta' = c, s1$ pseudofermions (and $\beta' = \alpha'\nu'$ pseudofermions of vanishing velocity) at the $\beta' = c, s1$ Fermi points (and at the limiting values $\pm q_{\alpha'\nu'}$). In this section we are mostly interested in the scattering centers associated with the deviation $\delta N^F_{\beta'}(q')$ and $\delta N^L_{\beta'}(q')$, whose general expression reads,

$$\begin{aligned}
\delta N^F_{\beta'}(q') &= \sum_{\iota=\pm1}\Big[\frac{\delta N^F_{\beta'}}{2} + \iota\,\delta J^F_{\beta'}\Big]\,\delta_{q',\iota\,q_{F\beta'}}\,; \quad \beta' = c, s1\,; \\
\delta N^L_{\beta'}(q') &= \sum_{\iota=\pm1}\Big[\frac{\delta N^L_{\beta'}}{2} + \iota\,\delta J^L_{\beta'}\Big]\,\delta_{q',\iota\,q_{\beta'}}\,; \quad \beta' = \alpha'\nu'\,.
\end{aligned} \tag{51}$$

For the $\beta' = s1$ branch the deviation $\delta N^L_{\beta'}(q')$ applies only to the vanishing-velocity $s1$ pseudofermion scattering centers with $s1$ limiting values $\pm q_{s1} = \pm 2k_F$, which corresponds to the $k_{F\uparrow} \rightarrow 2k_F$ and $k_{F\downarrow} \rightarrow 0$ limit. In the above equation the deviation numbers $\delta N^F_{\beta'}$ (and $\delta N^L_{\beta'}$) are such that $\delta N_{\beta'} = \delta N^F_{\beta'} + \delta N^{NF}_{\beta'}$ (and $\delta N_{\beta'} = \delta N^L_{\beta'} + \delta N^{NF}_{\beta'}$). They can be expressed as $\delta N^F_{\beta'} = \delta N^F_{\beta',+1} + \delta N^F_{\beta',-1}$ (and $\delta N^L_{\beta'} = \delta N^L_{\beta',+1} + \delta N^L_{\beta',-1}$). Here $\delta N^F_{\beta',\pm1}$ is the deviation in the number of $\beta' = c, s1$ pseudofermions at the right $(+1)$ and left (-1) $\beta' = c, s1$ Fermi point (and $\delta N^L_{\beta',\iota}$ is the deviation in the number of $\beta' = \alpha'\nu'$ pseudofermions created at $\iota\,q_{\beta'}$ with $\iota = \pm1$). The deviation current numbers (and deviation current numbers) read $2\delta J^F_{\beta'} = [\delta N^F_{\beta',+1} - \delta N^F_{\beta',-1}]$ for $\beta' = c, s1$ pseudofermions (and $2\delta J^L_{\beta'} = [\delta N^L_{\beta',+1} - \delta N^L_{\beta',-1}]$ for vanishing-velocity $\beta' = \alpha'\nu'$ pseudofermions). For the $\beta' = \alpha'\nu' \neq s1$ pseudofermions the deviations $\delta N_{\beta'} = \delta N^L_{\beta'} + \delta N^{NF}_{\beta'}$ equal the corresponding numbers $N_{\beta'} = N^L_{\beta'} + N^{NF}_{\beta'}$. Indeed the $\beta' = \alpha'\nu' \neq s1$ pseudofermion occupancy vanishes for initial ground states.

According to the PDT of Ref. [7], the elementary processes (A), (B), and (C) mentioned above lead to qualitatively different contributions to the spectral-weight distributions. The PDT studies of these references considered that creation of $\eta\nu'$ pseudofermions and $s\nu' \neq s1$ pseudofermions at the limiting bare-momentum values is felt by both the c and $s1$ scatterers as effective c scattering centers and effective c and $s1$ scattering centers, respectively. However, such studies applied only to $\beta = c$ scatterers and $\beta = s1$ scatterers of bare-momentum value $q = \pm q_{F\beta}$. Hence the general two-pseudofermion expression provided in Eq. (50) generalizes that result to *all* active $\beta = c, \alpha\nu$ scatterers of arbitrary bare momentum q.

From the linearity in the deviations of the overall scattering phase shift given in Eq. (11), one can write $Q^\Phi_\beta(q)/2 = [Q^{\Phi(NF)}_\beta(q)/2 + Q^{\Phi(F)}_\beta(q)/2]$. Also the part of the total momentum deviation of Eq. (27) associated with the elementary processes (A) and (B) can be written as $\delta P^{NF} + \delta P^F$. After some algebra involving the use of Eqs. (11), (27), (50), and (51), we reach the following expressions for such quantities,

$$\frac{Q^{\Phi(NF)}_\beta(q)}{2} = \sum_{\beta'}\sum_{q'} \pi\,\Phi_{\beta,\beta'}(q,q')\,\delta N^{NF}_{\beta'}(q')\,, \tag{52}$$

$$
\begin{aligned}
\frac{Q_{\beta}^{\Phi(F)}(q)}{2} &= \sum_{\beta'=c,s1}\sum_{\iota'=\pm 1}\pi\,\Phi_{\beta,\beta'}(q,\iota'\,q_{F\beta'})\,\frac{\delta N^F_{\beta'}}{2} \\
&+ \sum_{\iota'=\pm 1}\iota'\,\pi\,\Phi_{\beta,c}(q,\iota'\,2k_F)\Big[\delta J^F_c + \sum_{\nu'=1}^{\infty} J^L_{\eta\nu'} + \sum_{\nu'=2}^{\infty} J^L_{s\nu'} + \delta J^L_{s1}\times\delta_{v_{s1}(q_{s1}),0}\Big] \\
&+ \sum_{\iota'=\pm 1}\iota'\,\pi\,\Phi_{\beta,s1}(q,\iota'\,k_{F\downarrow})\Big[\delta J^F_{s1} - 2\sum_{\nu'=2}^{\infty} J^L_{s\nu'}\Big] \\
&+ \sum_{\beta'=\alpha'\nu'\neq s1}\pi\Big[\delta_{\beta,c} - \delta_{\beta,\alpha\nu}\delta_{\alpha,\alpha'}\left(-\delta_{\nu,\nu'}+\nu+\nu'-|\nu-\nu'|\right)\Big]J^L_{\alpha'\nu'} \\
&+ \pi\Big[\delta_{\beta,c} - \delta_{\beta,s\nu}\left(2-\delta_{\nu,1}\right)\Big]\delta J^L_{s1}\times\delta_{v_{s1}(q_{s1}),0}\,,
\end{aligned}
\tag{53}
$$

and

$$
\begin{aligned}
\delta P^{NF} &= \sum_{\beta'=c,s\nu'}\sum_{q'} q'\,\delta N^{NF}_{\beta'}(q') + \sum_{\eta\nu'}\sum_{q'}[\pi - q']\,\delta N^{NF}_{\eta\nu'}(q')\,; \\
\delta P^{F} &= \pi\,[M^{un}_{\eta,-1/2} + \sum_{\nu'=1}^{\infty}\nu' N^L_{\eta\nu'}] \\
&+ 4k_F\Big[\,\delta J^F_c + \sum_{\nu'=1}^{\infty} J^L_{\eta\nu'} + \sum_{\nu'=2}^{\infty} J^L_{s\nu'} + \delta J^L_{s1}\times\delta_{v_{s1}(q_{s1}),0}\Big] + 2k_{F\downarrow}\Big[\,\delta J^F_{s1} - 2\sum_{\nu'=2}^{\infty} J^L_{s\nu'}\Big]\,.
\end{aligned}
\tag{54}
$$

In the above expressions we used that $\delta J_{\beta'} = J_{\beta'}$ for the $\beta' = \alpha'\nu' \neq s1$ scattering centers.

The general expression of the phase shift $Q_{\beta}^{\Phi(F)}(q)/2$ given in Eq. (53) is valid for all active β scatterers. In the δP^F expression of Eq. (54) we have included the contribution from the unbound $-1/2$ η-spinons. (The momentum contributions from the unbound $+1/2$ η-spinons and $\pm 1/2$ unbound spinons vanish.) Note that the current contributions to the momentum spectrum δP^F given in Eq. (54), which multiply $4k_F$ and $2k_{F\downarrow}$, are identical to the current contributions to the scattering phase shift of Eq. (53), which multiply the phase shifts $\pi\,\Phi_{\beta,c}(q,\iota'\,2k_F)$ and $\pi\,\Phi_{\beta,s1}(q,\iota'\,k_{F\downarrow})$, respectively.

That the vanishing-velocity $\alpha'\nu'$ pseudofermions of limiting canonical-momentum values $\bar{q}\rightarrow\pm q_{\alpha'\nu'}$ are not active scatterers and are felt by the active scatterers as effective c scattering centers and both effective c and $s1$ scattering centers, respectively, follows from their invariance under the electron - rotated-electron unitary transformation. Such an invariance is related to the $2\nu' = 2, 4, ..., \infty$ η-spinons contained in a vanishing-velocity composite $\eta\nu'$ pseudofermion (and the $2\nu' = 2, 4, ..., \infty$ spinons contained in a vanishing-velocity composite $s\nu'$ pseudofermion) losing their anti-bound (and bound) character as $\bar{q}\rightarrow\pm q_{\eta\nu'}$ (and as $\bar{q}\rightarrow\pm q_{s\nu'}$). Consistent, in that limit the pseudofermion energy reaches its maximum (and minimum) magnitude, given in Eq. (37). Alike for the corresponding $\alpha'\nu'$ fermion, in that limit the $\alpha'\nu'$ pseudofermion energy is a mere sum of the $2\nu' = 2, 4, ..., \infty$ η-spinon ($\alpha = \eta$) or $2\nu' = 2, 4, ..., \infty$ spinon ($\alpha = s$) energies, so that Eq. (37) is obeyed. Each of the corresponding $2\nu'$ unbound η-spinons (and $2\nu'$ unbound spinons) has an energy, given in Eq. (39), equal to that of a unbound η-spinon (and a unbound spinon) with the same η-spin (and spin) projection.

The main difference between the $2\nu'$-η-spinon configurations (i) of $M^{un}_{\eta,-1/2} = \nu'$ unbound $-1/2$ η-spinons plus $M^{un}_{\eta,+1/2} = \nu'$ unbound $+1/2$ η-spinons and (ii) the $2\nu'$ anti-bound η-spinons of one vanishing-velocity $\eta\nu'$ pseudofermion of canonical momentum $\bar{q} = \pm q_{\eta\nu'}$ is that the former configuration has a total η-spin value, $S_\eta = \nu'$, whereas the latter is a η-spin-singlet configuration. (The same holds for vanishing-velocity $s\nu'$ pseudofermions of canonical momentum $\bar{q} = \pm q_{s\nu'}$ provided that one replaces the $2\nu'$ η-spinons by $2\nu'$ spinons.)

Hence for the $\alpha'\nu'$ branches the picture that emerges is that in the $\bar{q}\rightarrow\pm q_{\eta\nu'}$ (and $\bar{q}\rightarrow\pm q_{s\nu'}$) canonical-momentum limit the corresponding vanishing-velocity $\eta\nu'$ pseudofermion (and a vanishing-velocity $s\nu' \neq s1$ pseudofermion) decouples into $2\nu' = 2, 4, ..., \infty$ η-spinons of vanishing anti-binding energy (and $2\nu' = 2, 4, ..., \infty$ spinons of vanishing binding energy) and one $\eta\nu'$ (and $s\nu'$) FP current scattering center. By a $\eta\nu'$ (and $s\nu'$) FP current scattering center we mean here the elementary current $\delta J^L_{\eta\nu'} = \iota/2$ (and $\delta J^L_{s\nu'} = \iota/2$) generated by creation of one vanishing-velocity $\eta\nu'$ pseudofermion (and one vanishing-velocity $s\nu'$ pseudofermion) at canonical momentum $\bar{q} = \pm q_{\eta\nu'} = \pm[\pi - 2k_F]$ (and $\bar{q} = \pm q_{s\nu'} = \pm[k_{F\uparrow} - k_{F\downarrow}]$ for $s\nu' \neq s1$ and $\bar{q} = \pm q_{s1} = \pm 2k_F$ as $k_{F\uparrow}\rightarrow 2k_F$ and $k_{F\downarrow}\rightarrow 0$). As confirmed by the form of the phase shifts given in Eqs. (50) and (53), such elementary currents are felt by the β active scatterers as elementary shifts of both c Fermi-points (and both c and both $s1$ Fermi-points). This justifies the designation FP (from Fermi-points) current scattering center.

B. Effects of the pseudofermion invariances on their scattering properties for densities $n=1$ and/or $m=0$

The general pseudofermion scattering theory introduced in this paper also applies to PSs with initial $S_\eta=0$ and/or $S_s=0$ ground states, provided that the specific issues addressed here are accounted for. For an initial ground state with $S_\eta=0$ and thus electronic density $n=1$ (and $S_s=0$ and thus spin density $m=0$) one has that the number of $\eta\nu$ band (and $s\nu\neq s1$ band) discrete canonical momentum values $N_{a_{\eta\nu}}$ (and $N_{a_{s\nu}}$) vanishes. This means that the corresponding pseudofermion branch does not exist. If one takes into account that for a ground state the numbers $N_{\alpha\nu}$ vanish for all $\alpha\nu\neq s1$ branches, one finds from inspection of the $N^h_{\alpha\nu}$ expression provided in Eq. (A11) of Appendix A that for $S_\alpha=0$ the number $N_{a_{\alpha\nu}}=[N_{\alpha\nu}+N^h_{\alpha\nu}]$ vanishes as well.

For simplicity, we focus our attention onto excited energy eigenstates of such an initial ground state with a single $\eta\nu$ pseudofermion, $N_{\eta\nu}=1$, and vanishing number values, $N_{\eta\nu'}=0$, for $\nu'>\nu$ branches (and a single $s\nu\neq s1$ pseudofermion, $N_{s\nu}=1$, and vanishing number values, $N_{s\nu'}=0$, for $\nu'>\nu$ branches). For these excited states the use of Eqs. (A10) and (A11) of Appendix A leads to $N_{a_{\eta\nu}}=1$ (and $N_{a_{s\nu}}=1$). The corresponding $\eta\nu$ (and $s\nu$) band reduces to the bare momentum $q=\pm q_{\eta\nu}=0$ (and $q=\pm q_{s\nu}=0$.) Indeed, at $n=1$ (and $m=0$) one finds from Eq. (A12) of Appendix A that $q_{\eta\nu}=[\pi-2k_F]=0$ (and $q_{s\nu}=[k_{F\uparrow}-k_{F\downarrow}]=0$.)

The number $N^h_c=[N_a-2S_c]$ and the number N^h_{s1} whose general expression is given in Eq. (A11) of Appendix A may be written as $N^h_c=M^{un}_\eta+2\sum_{\nu=1}^\infty \nu N_{\eta\nu}$ and $N^h_{s1}=M^{un}_s+2\sum_{\nu=1}^\infty(\nu-1)N_{s\nu}$, respectively. $S_\eta=0$ and $n=1$ (and $S_s=0$ and $m=0$) initial ground states have full c bands (and $s1$ bands), so that $N^h_c=0$ (and $N^h_{s1}=0$.) Hence a property specific to excited energy eigenstates of $S_\eta=0$ and $n=1$ initial ground states is that their c band has 2ν c pseudofermion holes for each $\eta\nu$ pseudofermion and one c pseudofermion hole for each unbound η-spinon. Similarity, the $s1$ band of excited energy eigenstates of $S_s=0$ and $m=0$ initial ground states has $2(\nu-1)$ $s1$ pseudofermion holes for each $s\nu\neq s1$ pseudofermion and one $s1$ pseudofermion hole for each unbound spinon.

If one further considers the particular case of the initial $S_\eta=0; S_s=0; 2S_c=N_a$ absolute ground state, it follows from Eqs. (B26), (B35), (B36), and (B37) of Appendix B that $\pi\,\Phi_{c,s\nu'}(q,0)=\pi\,\Phi_{\eta\nu,s\nu'}(q,0)=\pi\,\Phi_{s\nu,\eta\nu'}(q,0)=0$ for $s\nu'\neq s1$. On the other hand, we find below that the value of the phase shifts $\pi\,\Phi_{c,\eta\nu}(q,0)$ (and $\pi\,\Phi_{s1,s\nu}(q,0)$) is fully determined by the 2ν (and $2(\nu-1)$) bare-momentum values of the c pseudofermion-hole (and $2(\nu-1)$ $s1$ pseudofermion-hole) scattering centers. Thus, for excited states of the initial $S_\eta=0; S_s=0; 2S_c=N_a$ absolute ground state the c and $s1$ scatterers feel the creation of one $\eta\nu$ pseudofermion (and one $s\nu\neq s1$ pseudofermion) as that of 2ν c effective scattering centers (and $s1$ effective scattering centers).

Such effective scattering centers are different from those considered above for the excited states of ground states with electronic density (and spin density) in the range obeying the inequality $0<n<1$ (and $0<m<n$). For the excited states of a $n=1$ (and $m=0$) initial ground state considered here the current number $J^L_{\eta\nu}$ (and $J^L_{s\nu}$) vanishes and thus there are no $\eta\nu$ (and $s\nu\neq s1$) FP current scattering centers. The occurrence of the type of c (and $s1$) effective scattering centers considered in this section follows from the non-scatterer character of the corresponding $\eta\nu$ pseudofermions (and $s\nu\neq s1$ pseudofermion), as discussed in the following.

In spite of the lack of ground-state $\eta\nu$ (and $s\nu\neq s1$) pseudofermion bands, the scattering theory can be generalized to an initial ground state with electronic density $n=1$ (and $m=0$). The point is that the "in" asymptote one-pseudofermion scattering states do not contribute to the direct-product expression of the initial ground state but rather to that of the "in" state defined in Section II C. For the excited states of $n=1$ (and $m=0$) ground states considered here, the $\eta\nu$ (and $s\nu\neq s1$) bare-momentum band corresponds to a single value at $q=\pm q_{\eta\nu}=0$ (and $q=\pm q_{s\nu}=0$). For initial ground states with densities $0<n<1$ and $0<m<n$ and except for non-physical higher-order $1/N_a$ contributions, the scattering canonical-momentum shift $Q^\Phi_\beta(q)/N_a$, Eq. (11), has the same value whether one uses the ground-state rapidity functions $\Lambda^0_\beta(q)$ and $\Lambda^0_{\beta'}(q')$ or the corresponding "out"-state (excited-energy-eigenstate) rapidity functions $\Lambda_\beta(q)$ and $\Lambda_{\beta'}(q')$ in the general expression provided in Eq. (14) for the two-pseudofermion phase shifts. Indeed, these two alternative definitions of the two-pseudofermion phase shifts lead to the same value of the functional $Q^\Phi_\beta(q)/N_a$ up to contributions of order $1/N_a$. The general pseudofermion scattering theory introduced above also applies to initial ground states with densities $n=1$ and/or $m=0$ provided that the following procedure is performed:

Two-pseudofermion $\pi\,\Phi_{\beta,\beta'}(q,q')$ phase-shift definition for transitions to the excited states of a $S_\alpha=0$ ground state where $\alpha=\eta$ and/or $\alpha=s$ – For such an initial ground state there are no $\beta=\alpha\nu\neq s1$ pseudofermion bands. Hence the two-pseudofermion expression of Eq. (14) should be replaced by $\pi\,\Phi_{\beta,\beta'}(q,q')=\pi\,\bar{\Phi}_{\beta,\beta'}(\Lambda_\beta(q)/u,\Lambda_{\beta'}(q')/u)$. Here the rapidity function $\Lambda_\beta(q)$ (and $\Lambda_{\beta'}(q')$) is that of the excited state if $\beta=\alpha\nu\neq s1$ (and $\beta'=\alpha'\nu'\neq s1$) and of the initial ground state if $\beta=c,s1$ (and $\beta'=c,s1$). Since the former rapidity function is that of the excited state under consideration, it follows that for the particular case of such an initial ground state the quantity given in Eq. (14) is a functional, rather than a function.

The energy of the $\eta\nu$ pseudofermion of bare momentum $q=\pm q_{\eta\nu}$ (and $s\nu\neq s1$ pseudofermion of bare momentum

$q = \pm q_{s\nu}$) of the excited states considered here obeys the equality of Eq. (37). Hence such objects are invariant under the electron - rotated-electron unitary transformation. This is alike for vanishing-velocity $\beta = \alpha\nu$ pseudofermions of canonical momentum $\pm q_{\alpha\nu}$ created under the transitions to the excited states of ground states with densities in the ranges $0 < n < 1$ and $0 < m < n$. Those were considered above in Section IV A. Consistent with that invariance and as discussed in the first paper, creation of such a $\eta\nu$ pseudofermion (and $s\nu$ pseudofermion) leads to a change ν in the number of lattice sites doubly occupied by both electrons and rotated electrons (and singly occupied by both spin-down electrons and spin-down rotated electrons). The 2ν η-spinons (and 2ν spinons) of that vanishing-velocity $\eta\nu$ pseudofermion (and vanishing-velocity $s\nu$ pseudofermion) are not energetically anti-bound (and bound).

In spite of their energetic non-anti-binding (and unbinding) character, such 2ν η-spinons (and 2ν spinons) remain being in a collective η-spin-singlet (and spin-singlet) configuration. The corresponding above mentioned c effective scattering centers (and $s1$ effective scattering centers) associated with the $\eta\nu$ pseudofermion (and $s\nu \neq s1$ pseudofermion) refer to such a η-spin-neutral (and spin-neutral) collective configuration.

Since $\alpha\nu \neq s1$ bands do not exist for a $S_\alpha = 0$ initial ground state, one has that $Q_{\alpha\nu}(0)/N_a = Q^{\Phi}_{\alpha\nu}(0)/N_a$ for a $\alpha\nu \neq s1$ pseudofermion created onto such a state. Moreover, the invariance under the electron - rotated-electron transformation of a $\alpha\nu \neq s1$ pseudofermion created onto a $S_\alpha = 0$ ground state implies that it is not an active scatterer. For the class of excited states considered here a necessary condition for such an object not being an active scatterer is that it is not a scatterer at all. Since the $\alpha\nu \neq s1$ bare-momentum band of the "in" state has a single value at $q = \pm q_{\alpha\nu} = 0$, it is required that the corresponding canonical-momentum band of the "out" state has also a single value at $\bar{q} = \pm q_{\alpha\nu} = 0$. This implies that $Q_{\alpha\nu}(0)/N_a = Q^{\Phi}_{\alpha\nu}(0)/N_a = 0$, and thus that the $\alpha\nu \neq s1$ fermion remains invariant under the fermion - pseudofermion unitary transformation. This is a confirmation that the corresponding $\alpha\nu \neq s1$ pseudofermion is not a scatterer.

However, the requirement that such a $\alpha\nu \neq s1$ pseudofermion is not a scatterer and thus that $Q^{\Phi}_{\alpha\nu}(0)/N_a = 0$ imposes a specific form to the corresponding effective two-peudofermion phase shifts $\pi\,\Phi_{\alpha\nu,\beta'}(0,q')$. Since such objects are not scatterers, the quantities $\pi\,\Phi_{\alpha\nu,\beta'}(0,q')$ are not actual two-peudofermion phase shifts. They may be seen as mere effective two-peudofermion phase shifts whose values are such that the overall scattering phase shift $Q^{\Phi}_{\alpha\nu}(0)/2$ vanishes. Thus, there is no requirement for them to obey Levinson's Theorem.

For simplicity, we consider $S_\eta = 0; S_s = 0; 2S_c = N_a$ excited energy eigenstates of the initial $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state. Hence such excited states have no unbound η-spinons and no unbound spinons. Specifically, we consider three types of such excited states with pseudofermion occupancy in the c and $s1$ bands, none or a finite number M^{un}_η of unbound η-spinons and M^{un}_s of unbound spinons plus (a) one $\eta\nu$ pseudofermion and one $s\nu' \neq s1$ pseudofermion, (b) one $\eta\nu$ pseudofermion, and (c) one $s\nu' \neq s1$ pseudofermion. The initial $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state is described by full c and $s1$ pseudofermion bands whose c and $s1$ Fermi momenta read $q_{Fc} = 2k_F = q_c = \pi$ and $q_{Fs1} = k_{F\downarrow} = q_{s1} = k_{F\uparrow} = k_F = \pi/2$, respectively. Thus, from the use of Eq. (A11) of Appendix A we have that $\delta N_c = -\delta N^h_c = -(M^{un}_\eta + 2\nu)$, $\delta N_{s1} = -[(M^{un}_\eta + M^{un}_s)/2 + (\nu + \nu')]$, $\delta N^h_{s1} = [M^{un}_s + 2(\nu' - 1)]$ for the excited states (a), $\delta N_c = -\delta N^h_c = -(M^{un}_\eta + 2\nu)$, $\delta N_{s1} = -[(M^{un}_\eta + M^{un}_s)/2 + \nu]$, $\delta N^h_{s1} = M^{un}_s$ for the excited states (b), and $\delta N_c = -\delta N^h_c = -M^{un}_\eta$, $\delta N_{s1} = -[(M^{un}_\eta + M^{un}_s)/2 + \nu']$, $\delta N^h_{s1} = [M^{un}_s + 2(\nu' - 1)]$ for the excited states (c). On the other hand, according to Eqs. (B26), (B27), and (B37) of Appendix B, within the simultaneous $m \to 0$ and $n \to 1$ limits the two-pseudofermion phase shifts that contribute to $Q^{\Phi}_{\eta\nu}(0)/2$ and $Q^{\Phi}_{s\nu'}(0)/2$ simplify to $\pi\,\bar{\Phi}_{\eta\nu,s\nu'}(r,r') = \pi\,\bar{\Phi}_{s\nu',c}(r',r) = \pi\,\bar{\Phi}_{s\nu',\eta\nu}(r',r) = 0$, $\pi\,\bar{\Phi}_{\eta\nu,c}(r,r') = \arctan\left(\frac{r-r'}{\nu}\right)$, and $\pi\,\bar{\Phi}_{s\nu',s1}(r',r) = \arctan\left(\frac{r'-r}{\nu'-1}\right)$ for $s\nu' \neq s1$. It follows then from the use of Eq. (11) that for the excited energy eigenstates (a)-(c) the equation $Q^{\Phi}_{\eta\nu}(0)/2 = 0$ and/or $Q^{\Phi}_{s\nu'}(0)/2 = 0$ leads to,

$$
\begin{aligned}
\frac{Q^{\Phi}_{\eta\nu}(0)}{2} &= \sum_{h=1}^{M^{un}_\eta + 2\nu} \pi\,\Phi_{\eta\nu,c}(0,q_h) = \sum_{h=1}^{M^{un}_\eta + 2\nu} \arctan\left(\frac{\Lambda_{\eta\nu}(0) - \Lambda^0_c(q_h)}{\nu\,u}\right) = 0\,, \\
\frac{Q^{\Phi}_{s\nu'}(0)}{2} &= \sum_{h=1}^{M^{un}_s + 2(\nu'-1)} \pi\,\Phi_{s\nu',s1}(0,{q'}_h) = \sum_{h=1}^{M^{un}_s + 2(\nu'-1)} \arctan\left(\frac{\Lambda_{s\nu'}(0) - \Lambda^0_{s1}({q'}_h)}{(\nu'-1)\,u}\right) = 0\,, \qquad s\nu' \neq s1\,. \qquad (55)
\end{aligned}
$$

Here the first and second equations refer to the c branch and both the states (a) and (b) and to the $s1$ branch and both the states (a) and (c), respectively. In these equations the set of $M^{un}_\eta + 2\nu = 2, 4, 6, ...$ values $\{q_h\}$ and of $M^{un}_s + 2(\nu' - 1) = 2, 4, 6, ...$ values $\{{q'}_h\}$ correspond to the excited-energy-eigenstate c pseudofermion holes and $s1$ pseudofermion holes, respectively. The ground-state rapidity functions $\Lambda^0_c(q) = \sin k_0(q)$ and $\Lambda^0_{s1}(q)$ are defined in terms of their inverse functions, as provided in Eq. (A14) of Appendix A. For the present $n = 1$ and $m = 0$ ground state such inverse functions may be written explicitly, as given in Eq. (A16) of that appendix. The form of the equalities given in Eq. (55) reveals that the corresponding solutions, $\Lambda_{\eta\nu}(0) = \Lambda_{\eta\nu}(0, \{q_h\})$ and/or $\Lambda_{s\nu'}(0) = \Lambda_{s\nu'}(0, \{{q'}_h\})$, are functions of the above sets of bare-momentum values $\{q_h\}$ and $\{{q'}_h\}$, respectively.

We obtained here Eq. (55), which the functions $\Lambda_{\eta\nu}(0) = \Lambda_{\eta\nu}(0, \{q_h\})$ and $\Lambda_{s\nu'}(0) = \Lambda_{s\nu'}(0, \{q'_h\})$ for the rapidities $\Lambda_{\eta\nu}(0)$ and $\Lambda_{s\nu'}(0)$, respectively, obey, from the invariance under that unitary transformation of the corresponding $\eta\nu$ and $s\nu' \neq s1$ pseudofermions. For the above excited states the solution of the thermodynamic BA equations, Eqs. (A2)-(A4) of Appendix A at $\phi_\uparrow = \phi_\downarrow = 0$, exactly leads to the same functions, thus also obeying Eq. (55). This confirms that for such excited states the exact solution of the BA equations is equivalent to imposing the symmetry requirement $Q^{\Phi}_{\eta\nu}(0)/2 = 0$ and $Q^{\Phi}_{s\nu'}(0)/2 = 0$ associated with the invariance under the electron - rotated-electron unitary transformation and corresponding non-scatterer character of the $\eta\nu$ pseudofermion and/or $s\nu'$ pseudofermion under consideration, respectively. Here the BA solution accounts for that symmetry.

The above functions $\Lambda_{\eta\nu}(0) = \Lambda_{\eta\nu}(0, \{q_h\})$ and $\Lambda_{s\nu'}(0) = \Lambda_{s\nu'}(0, \{q'_h\})$ are to be used in the following expressions,

$$\begin{aligned}
\pi\,\Phi_{\eta\nu,c}(0,q') &= \pi\,\bar{\Phi}_{\eta\nu,c}\Big(\frac{\Lambda_{\eta\nu}(0,\{q_h\})}{u}, \frac{\Lambda^0_c(q')}{u}\Big) = \arctan\Big(\frac{\Lambda_{\eta\nu}(0,\{q_h\}) - \Lambda^0_c(q')}{\nu\, u}\Big)\,; \\
\pi\,\Phi_{s\nu',s1}(0,q') &= \pi\,\bar{\Phi}_{s\nu',s1}\Big(\frac{\Lambda_{s\nu'}(0,\{q'_h\})}{u}, \frac{\Lambda^0_{s1}(q')}{u}\Big) = \arctan\Big(\frac{\Lambda_{s\nu'}(0,\{q'_h\}) - \Lambda^0_{s1}(q')}{(\nu'-1)\, u}\Big)\,. \qquad (56)
\end{aligned}$$

They assure that $Q^{\Phi}_{\eta\nu}(0)/2 = 0$ and/or $Q^{\Phi}_{s\nu'}(0)/2 = 0$. The simplest case corresponds to $M^{un}_\eta = M^{un}_s = 0$ and $\nu = 1$ and/or $\nu' = 2$. In that case the solution of Eq. (55) leads to $\Lambda_{\eta 1}(0, q_1, q_2) = [\Lambda^0_c(q_1) + \Lambda^0_c(q_2)]/2$ and/or $\Lambda_{s2}(0, q'_1, q'_2) = [\Lambda^0_{s1}(q'_1) + \Lambda^0_{s1}(q'_2)]/2$, respectively. Hence, the symmetry requirement that the $\eta\nu$ pseudofermion (and $s\nu \neq s1$ pseudofermion) considered here is not a scatterer implies that the corresponding rapidity function $\Lambda_{\eta\nu}(0) = \Lambda_{\eta\nu}(0, \{q_h\})$ (and $\Lambda_{s\nu'}(0) = \Lambda_{s\nu'}(0, \{q'_h\})$) does not in general vanish and is the unique solution of the first (and second) equation in Eq. (55).

On the other hand, combination of this result with the two-pseudofermion phase shift expressions provided in Eqs. (B35) and (B36) of Appendix B, reveals that the c scatterer two-pseudofermion phase shift, $\pi\,\Phi_{c,\eta\nu}(q,0)$, and the $s1$ scatterer two-pseudofermion phase shift, $\pi\,\Phi_{s1,s\nu'}(q,0)$, are such that,

$$\begin{aligned}
\pi\,\Phi_{c,\eta\nu}(q,0) &= \pi\,\bar{\Phi}_{c,\eta\nu}\Big(\frac{\Lambda^0_c(q)}{u}, \frac{\Lambda_{\eta\nu}(0,\{q_h\})}{u}\Big) = -\arctan\Big(\frac{\Lambda^0_c(q) - \Lambda_{\eta\nu}(0,\{q_h\})]}{\nu\, u}\Big)\,; \\
\pi\,\Phi_{s1,s\nu'}(q,0) &= \pi\,\bar{\Phi}_{s1,s\nu'}\Big(\frac{\Lambda^0_{s1}(q)}{u}, \frac{\Lambda_{s\nu'}(0,\{q'_h\})}{u}\Big) = \arctan\Big(\frac{\Lambda^0_{s1}(q) - \Lambda_{s\nu'}(0,\{q'_h\})]}{(\nu'-1)\, u}\Big)\,; \quad q \neq \pm k_F \\
&= \pm\frac{\pi}{\sqrt{2}}\,; \qquad q = \pm k_F\,. \qquad (57)
\end{aligned}$$

In addition to the c or $s1$ scatterer bare-momentum q, the two-pseudofermion phase shifts provided in Eq. (57) are functions of the set of $M^{un}_\eta + 2\nu = 2, 4, 6, ...$ bare-momentum values $\{q_h\}$ or $M^{un}_s + 2(\nu'-1) = 2, 4, 6, ...$ bare-momentum values $\{q'_h\}$ of the $M^{un}_\eta + 2\nu = 2, 4, 6, ...$ c pseudofermion-hole scattering centers or $M^{un}_s + 2(\nu'-1) = 2, 4, 6, ...$ $s1$ pseudofermion-hole scattering centers, respectively. The latter are created under the ground-state - excited-energy-eigenstate transition. This confirms that as a result of the creation of the $\eta\nu$ (and $s\nu' \neq s1$) pseudofermion, the c (and $s1$) scatterers acquire the phase shift $\pi\,\Phi_{c,\eta\nu}(q,0)$ (and $\pi\,\Phi_{s1,s\nu'}(q,0)$). Its value is fully controlled by the $M^{un}_\eta + 2\nu = 2, 4, 6, ...$ (and $M^{un}_s + 2(\nu'-1) = 2, 4, 6, ...$) bare-momentum values of the $M^{un}_\eta + 2\nu = 2, 4, 6, ...$ (and $M^{un}_s + 2(\nu'-1) = 2, 4, 6, ...$) c (and $s1$) pseudofermion-hole scattering centers. Thus, through the $\{q_h\}$ (and $\{q'_h\}$) momentum dependence of the two-pseudofermion phase shift $\pi\,\Phi_{c,\eta\nu}(q,0)$ (and $\pi\,\Phi_{s1,s\nu'}(q,0)$), the c (and $s1$) scatterers feel the created $\eta\nu$ (and $s\nu' \neq s1$) pseudofermion as $M^{un}_\eta + 2\nu = 2, 4, 6, ...$ c effective scattering centers (and $M^{un}_s + 2(\nu'-1) = 2, 4, 6, ...$ $s1$ effective scattering centers).

Similar results are obtained for excited states of initial ground states with density $n = 1$ and/or $m = 0$ and occupancy of a larger finite number of β pseudofermions belonging to several $\beta = \alpha\nu \neq s1$ branches. In that general case, the number of equations defining the set of rapidities $\{\Lambda_{\alpha\nu}\}$ is in general larger than above and each of these equations is more involved than the two equations given in Eq. (55). Moreover, the equations that follow from the $Q^{\Phi}_{s\nu'}(0)/2 = 0$ and $Q^{\Phi}_{\eta\nu}(0)/2 = 0$ symmetry requirements involve additional phase shifts $\pi\,\bar{\Phi}_{s\nu',s\nu''}(r, r'')$ and $\pi\,\bar{\Phi}_{\eta\nu,\eta\nu'}(r, r')$ defined by Eqs. (B15) and (B37) of Appendix B, respectively. The corresponding $\beta = \alpha\nu \neq s1$ pseudofermions are also invariant under both the electron - rotated-electron unitary transformation and fermion - pseudofermion unitary transformation.

Finally, the expression for $Q^{\Phi}_{\eta\nu}(0)/2$ (and $Q^{\Phi}_{s\nu'}(0)/2$) given in Eq. (55) is also valid for the type of excited states of $S_\eta = 0$ initial ground states with spin density $m \neq 0$ (and of $S_s = 0$ initial ground states with electronic density $n \neq 1$) with the following occupancies: Pseudofermion occupancy in the c and $s1$ bands, none or a finite number M^{un}_η of unbound η-spinons and M^{un}_s of unbound spinons plus one $\eta\nu$ pseudofermion (and one $s\nu' \neq s1$ pseudofermion). Indeed, within the $n \to 1$ limit for $m \neq 0$ (and the $m \to 0$ limit for $n \neq 1$) the two-pseudofermion phase shifts that contribute to $Q^{\Phi}_{\eta\nu}(0)/2$ (and $Q^{\Phi}_{s\nu'}(0)/2$) also simplify to $\pi\,\bar{\Phi}_{\eta\nu,s\nu'}(r, r') = 0$ and $\pi\,\bar{\Phi}_{\eta\nu,c}(r, r') = \arctan\Big(\frac{r-r'}{\nu}\Big)$ (and to $\pi\,\bar{\Phi}_{s\nu',c}\,(r',r) = \pi\,\bar{\Phi}_{s\nu',\eta\nu}\,(r',r) = 0$ and $\pi\,\bar{\Phi}_{s\nu',s1}\,(r',r) = \arctan\Big(\frac{r'-r}{\nu'-1}\Big)$ for $s\nu' \neq s1$.) Similarly, the expressions for $\pi\,\Phi_{\eta\nu,c}(0,q')$ and $\pi\,\Phi_{c,\eta\nu}(q,0)$ (and $\pi\,\Phi_{s\nu',s1}(0,q')$ and $\pi\,\Phi_{s1,s\nu'}(q,0)$) provided in Eqs. (56) and (57), respectively, are valid as well for the above type of excited states of $S_\eta = 0$ initial ground states with spin density $m \neq 0$ (and excited states of $S_s = 0$ initial ground states with electronic density $n \neq 1$).

V. COMPLEMENTARY ROLES OF THE $\beta = s\nu, c$ PSEUDOFERMIONS AND UNBOUND SPINONS IN THE TRANSPORT OF SPIN

In this section we continue the studies of the elementary objects of the 1D Hubbard model. Although the unbound η-spinons and unbound spinons are neither scatterers nor scattering centers, here they are found to play an important role in the model physics: They fully control the effects of the η-spin and spin transverse fluctuations on the charge and spin currents carried by the excited states.

The studies of this section also investigate the related problem of the complementary roles in the transport of spin played by the $s\nu$ pseudofermions for all spin densities and by the c pseudofermions for finite values of the spin density, due to their scattering events with the $s\nu$ pseudofermions, on the one hand, and by the unbound spinons on the other hand. Similar results apply to the complementary roles in the transport of charge played by the $\eta\nu$ pseudofermions and the c pseudofermions, on the one hand, and the unbound η-spinons on the other hand [50]. Although our study focuses mainly on the transport of spin, we consider as well the charge currents of some classes of energy eigenstates. The properties reported in this section for the c pseudofermions, $s\nu$ pseudofermions, and their band holes, apply as well to the corresponding c fermions, $s\nu$ fermions, and their band holes, respectively.

A. The general problem

It is shown in the first paper that the $s\nu$ fermion occupancy configurations generate the spin-singlet representations of the spin $SU(2)$ symmetry algebra. Within a PS, that applies as well to the corresponding $s\nu$ pseudofermion occupancy configurations. On the other hand, the unbound-spinon occupancies generate the spin-multiplet non-LWS tower of states from the Bethe states. Thus the $s\nu$ pseudofermion occupancy configurations and the unbound spinon occupancies refer to different degrees of freedom of the spin $SU(2)$ symmetry algebra. Consistent, their contribution to the spin currents has also a different nature, as discussed in this section. The spin-1/2 unbound spinons exist in their own right, besides playing the role of unoccupied sites of the $s\nu$ effective lattices of the $\nu = 1, ..., \infty$ branches. As given in Eq. (A11) of Appendix A for $\alpha = s$, the number of such sites reads $N^h_{s\nu} = [M^{un}_s + \sum_{\nu'=\nu+1}^{\infty} 2(\nu'-\nu)N_{s\nu'}]$ where $M^{un}_s = 2S_s$ denotes the number of unbound spinons. (For states with finite occupancy of $s\nu'$ pseudofermions with $\nu' > \nu$ spinon pairs, also $2(\nu'-\nu)$ of the spin-effective lattice $2\nu'$ sites occupied by each such $s\nu'$ pseudofermions is used by the $s\nu$ pseudofermions as unoccupied sites of their $s\nu$ effective lattice.)

Upon moving around the spin effective lattice, the neutral 2ν-spinon $s\nu$ pseudofermions interchange position with the unbound spinons. As discussed above in Section II C and in the first paper, the unbound spin-1/2 spinons play the passive role of unoccupied sites of both the spin effective lattice and the corresponding $s\nu$ effective lattices. Thus the objects that move around in the spin effective lattice are the $s\nu$ pseudofermions. In that motion they use all the $M^{un}_s = 2S_s$ unbound spinons and the $2(\nu'-\nu)$ bound spinons in the sets of $2(\nu'-\nu)$ sites of the spin effective lattice out of the $2\nu'$ sites occupied by each $s\nu'$ pseudofermion with $2\nu' > 2\nu$ bound spinons. Furthermore, the holes in the corresponding BA $s\nu$ momentum bands exist in their own right and must not be treated as being the $M^{un}_s = 2S_s$ unbound spinons and $\sum_{\nu'=\nu+1}^{\infty} 2(\nu'-\nu)N_{s\nu'}$ unbound spinons that play the role of unoccupied sites. The $M^{un}_s = 2S_s$ unbound spinons have their independent occupancies and the $s\nu'$ pseudofermion with $2\nu' > 2\nu$ bound spinons their own independent $s\nu'$ momentum bands. Consistent, the $s\nu$ effective lattice object motion can alternatively be described in terms of $s\nu$ pseudofermions and $s\nu$ pseudofermion holes.

Since the unbound spinons do not contribute to the states total momentum, the spin currents associated with the spin $U(1)$ symmetry sub-algebra longitudinal $S^{x_3}_s$ fluctuations are carried by the $s\nu$ band occupancy configurations. On the other hand, the unbound-spinon occupancies fully control the transverse fluctuations that change the $S^{x_3}_s$-spin currents of the spin $SU(2)$ tower non-LWSs relative to those of the initial LWSs, which are the Bethe states. Indeed application onto the latter states of the spin $SU(2)$ off-diagonal generators flips the spins of the unbound spinons yet leaves invariant the occupancy configurations of the remaining spinons, which are bound within the spin-neutral 2ν-spinon $s\nu$ pseudofermions.

The c pseudofermions and the spinons describe different degrees of freedom of the same original-lattice sites singly occupied by the rotated electrons. The bare c pseudofermions are spin-less objects that carry charge e. The bare two-spinon composite $s\nu$ pseudofermions are charge-less objects that refer to spin-singlet occupancy configurations. Nonetheless, as a result of the scattering processes studied in this paper, for finite spin density the c pseudofermions are found below to contribute to the spin currents. Moreover, for electronic densities $n \neq 1$ the $s1$ band occupancy configurations are found in the following to contribute to the charge currents. Such exotic effects confirm the important role played by the pseudofermion scattering theory in the transport of charge and spin.

The use of the 1D Hubbard model in a uniform vector potential, Eq. (A1), which can also be solved by the BA, simplifies the calculation of the expectation values of the spin and charge current operators of the Bethe states. These operators read,

$$\hat{J}^{\sigma_{x3}} = -\left(\frac{1}{2}\right) i\,t \sum_{\sigma} \sum_{j=1}^{N_a} \sigma \left[c^{\dagger}_{j,\sigma}\, c_{j+1,\sigma} - c^{\dagger}_{j+1,\sigma}\, c_{j,\sigma}\right] ,$$

$$\hat{J}^{\rho} = -(e)\, i\,t \sum_{\sigma} \sum_{j=1}^{N_a} \left[c^{\dagger}_{j,\sigma}\, c_{j+1,\sigma} - c^{\dagger}_{j+1,\sigma}\, c_{j,\sigma}\right] , \qquad (58)$$

respectively. Moreover, their Bethe-state expectation values are given by,

$$J_0^{\sigma_{x3}} = \left(\frac{1}{2}\right) \frac{\partial E_{l_o,l^0_{\triangle}}(\phi)}{\partial(\phi/N_a)}|_{\phi=0}\,; \quad \phi = \phi_{\uparrow} = -\phi_{\downarrow}\,; \qquad J_0^{\rho} = (e)\frac{\partial E_{l_o,l^0_{\triangle}}(\phi)}{\partial(\phi/N_a)}|_{\phi=0}\,, \quad \phi = \phi_{\uparrow} = \phi_{\downarrow}\,, \qquad (59)$$

respectively. Here $E_{l_o,l^0_{\triangle}}$ stands for the general energy eigenvalue spectrum given in Eq. (A8) of Appendix A.

B. Simple excited states spin and charge currents and corresponding effective charges and spin projections

In the following we consider the spin and charge currents carried by excited states of an initial ground state with densities in the ranges $n \in [0, 1]$ and $m \in [0, n]$. The object numbers of the latter state read [5],

$$M_s = M^{bo}_s + M^{un}_{s,+1/2} = N = n\, N_a\,; \quad M^{bo}_s = 2N_{\downarrow}\,; \quad M^{un}_{s,+1/2} = N_{\uparrow} - N_{\downarrow} = m\, N_a\,,$$

$$M_{\eta} = M^{un}_{\eta} = M^{un}_{\eta,+1/2} = N_a - N = (1-n)\, N_a\,,$$

$$N_c = N\,; \quad N_{s1} = N_{\downarrow}\,; \quad N^h_{s1} = N_{\uparrow} - N_{\downarrow} = m\, N_a\,. \qquad (60)$$

For that ground state the values of the following numbers vanish: $M^{un}_{\eta,-1/2}$, M^{bo}_{η}, $M^{un}_{s,-1/2}$, and set of all $\alpha\nu$ pseudofermion numbers $\{N_{\alpha\nu}\}$ except that of the $s1$ pseudofermions. In addition, the $s1$ band momenta are occupied for $q \in [-q_{Fs1}, q_{Fs1}] = [-k_{F\downarrow}, k_{F\downarrow}]$ and unoccupied for $|q| \in [q_{Fs1}, q_{s1}] = [k_{F\downarrow}, k_{F\uparrow}]$. The c band momenta are occupied for $q \in [-q_{Fc}, q_{Fc}] = [-2k_F, 2k_F]$ and unoccupied for $|q| \in [q_{Fc}, q_c] = [2k_F, \pi]$ where $2k_F = \pi\, n$. Although here we have ignored corrections of $1/N_a$ order, which vanish within the thermodynamic limit, we chose the number of electrons $N_{\uparrow}$ and $N_{\downarrow}$ values so that the right and left β Fermi points are exactly symmetrical for both the $\beta = c$ and $\beta = s1$ momentum bands. In that case both the spin and charge currents defined in Eq. (59) vanish for the initial ground state.

In general, the derivation of the spin and charge currents of Eq. (59) requires the numerical solution of Eqs. (A2)-(59) of Appendix A [51]. We start by considering a class of simple excited states whose numbers are provided in Eq. (60). Such states are generated from it by the elementary processes called processes (B) in Section IV A. Their $\beta = c, s1$ band momentum distribution function deviations are of the form $\delta N^F_{\beta}(q) = \sum_{\iota=\pm 1}[\delta N^F_{\beta}/2 + \iota\, \delta J^F_{\beta}]\, \delta_{q,\iota\, q_{F\beta}}$, as given in Eq. (51). An example of such excited states are all ground states with electron numbers different from those of the initial ground state. The corresponding transitions are described by this type of $\beta = c, s1$ band momentum distribution deviations, which involve only momenta in the vicinity of the right ($\iota = +1$) and left ($\iota = -1$) $\beta = c, s1$ Fermi points.

For such excited states, the solution of Eqs. (A2)-(59) of Appendix A is a relatively simpler problem, which gives a spin current of the general form,

$$J_0^{\sigma_{x3}} = \frac{1}{2} \sum_{\beta=c,s1} j^{\sigma_{x3}}_{\beta}\, \delta 2J^F_{\beta}\,. \qquad (61)$$

Here the elementary spin currents $j^{\sigma_{x3}}_c$ and $j^{\sigma_{x3}}_{s1}$ are given by,

$$j^{\sigma_{x3}}_c = v_c\, \xi^1_{c,c}\, [\xi^1_{c,c} - 2\xi^1_{c,s1}] + v_{s1}\, \xi^1_{s1,c}\, [\xi^1_{s1,c} - 2\xi^1_{s1,s1}]\,,$$

$$j^{\sigma_{x3}}_{s1} = v_{s1}\, \xi^1_{s1,s1}\, [\xi^1_{s1,c} - 2\xi^1_{s1,s1}] + v_c\, \xi^1_{c,s1}\, [\xi^1_{c,s1} - 2\xi^1_{c,c}]\,, \qquad (62)$$

where v_β is for the $\beta = c, s1$ branches the pseudofermion velocity at the β-band Fermi momentum $q_{F\beta}$ given in Eq. (A23) of Appendix A. The parameter $\xi^1_{\beta,\beta'}$ can be solely expressed in terms of two-pseudofermion phase shifts in units of π as follows,

$$\xi^1_{\beta,\beta'} = \delta_{\beta,\beta'} + \sum_{\iota=\pm 1} (\iota)\, \Phi_{\beta,\beta'}\left(q_{F\beta}, \iota q_{F\beta'}\right) , \quad \beta = c, s1 \, . \tag{63}$$

For the particular case of $\beta = \beta'$ and $\iota = 1$ in this expression, our notation assumes that the two β Fermi momenta in the argument of the two-pseudofermion phase shift, $\Phi_{\beta,\beta}\left(q_{F\beta}, q_{F\beta}\right)$, differ by $2\pi/N_a$. This is consistent with the Pauli principle concerning the BA β band discrete momentum values occupancies, which can be zero (β pseudofermion hole) or one (β pseudofermion) [5]. Moreover, a β pseudofermion or β pseudofermion hole does not scatter into itself, so that $\Phi_{\beta,\beta}\left(q_{F\beta}, q_{F\beta}\right)$ vanishes if the two β Fermi momenta in the argument of $\Phi_{\beta,\beta}\left(q_{F\beta}, q_{F\beta}\right)$ are exactly the same.

Creation of one β pseudofermion (and one β pseudofermion hole) at the $\iota = \pm 1$ β Fermi point generates a current deviation $\delta 2J^F_\beta$ given by $\delta 2J^F_\beta = \iota$ (and $\delta 2J^F_\beta = -\iota$) and thus a corresponding spin current $\iota(1/2) j^{\sigma_{x3}}_\beta$ (and $-\iota(1/2) j^{\sigma_{x3}}_\beta$). Hence for the present excited states, $j^{\sigma_{x3}}_c$ and $-j^{\sigma_{x3}}_{s1}$ are the elementary spin currents in units of 1/2 carried by the c pseudofermions and $s1$ pseudofermion holes, respectively.

The form of the spin currents given in Eqs. (61)-(63) confirms the major role played by the present scattering theory two-pseudofermion phase shifts in the transport of spin. For excited states of the type considered here whose initial ground state is a zero-spin-density state the values of the two-pseudofermion phase shifts appearing in the expressions provided in Eqs. (62) and (63) are such that $j^{\sigma_{x3}}_c = 0$. Hence as expected for an initial ground state with zero spin density, $m = 0$, the c pseudofermions do not contribute to the spin current. However, as a result of their scattering with the $s1$ pseudofermion holes, for excited states of $m \neq 0$ ground states the c pseudofermions carry with them a finite spin current $j^{\sigma_{x3}}_c$. For instance, in the $m \to 0$ and $m \to n$ spin-density limits and for electronic density in the range $n \in [0, 1]$ the expressions of the elementary spin currents $j^{\sigma_{x3}}_c$ and $j^{\sigma_{x3}}_{s1}$ given in Eq. (62) read,

$$\begin{aligned}
j^{\sigma_{x3}}_c &= 0 \, ; \quad j^{\sigma_{x3}}_{s1} = -v_{s1} \, , \quad m \to 0 \, , \\
j^{\sigma_{x3}}_c &= v_c + v_{s1} \left[1 - \frac{4}{\pi} \arctan\left(\frac{\sin \pi n}{u}\right)\right] \frac{2}{\pi} \arctan\left(\frac{\sin \pi n}{u}\right) \, , \quad m \to n \, , \\
j^{\sigma_{x3}}_{s1} &= -v_{s1} \left[2 - \frac{2}{\pi} \arctan\left(\frac{\sin \pi n}{u}\right)\right] \, . \quad m \to n \, ,
\end{aligned} \tag{64}$$

For the low-energy excited states considered here the solution of Eqs. (A2)-(59) of Appendix A provides as well a charge current of the general form,

$$J^\rho_0 = -e \sum_{\beta=c,s1} j^\rho_\beta \, \delta 2J^F_\beta \, , \tag{65}$$

where the elementary charge currents in units of e carried by the $\beta = c, s1$ pseudofermions are given by,

$$\begin{aligned}
j^\rho_c &= v_c \, [\xi^1_{c,c}]^2 + v_{s1} \, [\xi^1_{s1,c}]^2 \, , \\
j^\rho_{s1} &= v_c \, \xi^1_{c,c} \xi^1_{c,s1} + v_{s1} \, \xi^1_{s1,s1} \xi^1_{s1,c} \, .
\end{aligned} \tag{66}$$

Those are again fully controlled by the two-pseudofermion phase shifts produced by the scattering-theory events. For electronic densities in the range $n \in [0, 1]$ and the $m \to 0$ and $m \to n$ spin-density limits the expressions of the elementary charge currents, Eq. (66), simplify to,

$$\begin{aligned}
j^\rho_c &= v_c \, \xi_0^2 \, ; \quad j^\rho_{s1} = v_c \, \frac{\xi_0^2}{2} \, , \quad m \to 0 \, , \\
j^\rho_c &= v_c + v_{s1} \left[\frac{2}{\pi} \arctan\left(\frac{\sin \pi n}{u}\right)\right]^2 \, , \quad m \to n \, , \\
j^\rho_{s1} &= v_{s1} \left[\frac{2}{\pi} \arctan\left(\frac{\sin \pi n}{u}\right)\right] \, , \quad m \to n \, .
\end{aligned} \tag{67}$$

The dependence on the electronic density $n \in [0, 1]$ and on-site repulsion in units of $4t$, $u = U/4t$, of the $m = 0$ parameter ξ_0 appearing here is given by $\xi_0 = \xi_0(r^0_c)$ where the function $\xi_0(r)$ obeys the following integral equation,

$$\begin{aligned}
\xi_0(r) &= 1 + \int_{-r^0_c}^{r^0_c} dr' A(r - r') \, \xi_0(r') \, , \\
A(r) &= \frac{1}{\pi} \int_0^\infty d\omega \frac{\cos(\omega \, r)}{1 + e^{2\omega}} \, ,
\end{aligned} \tag{68}$$

and r_c^0 is defined in Eq. (A15) of Appendix A. The parameter ξ_0 has limiting values $\xi_0 = \sqrt{2}$ for $u \rightarrow 0$ and $\xi_0 = 1$ for $u \rightarrow \infty$.

For the present type of low-energy excited states, one may introduce an effective spin projection and an effective charge for the c pseudofermions and $s1$ pseudofermion holes with momenta at the corresponding Fermi points,

$$s_\beta^{\sigma_{x3}} = i_\beta \frac{1}{2}\frac{j_\beta^{\sigma_{x3}}}{v_\beta}\,; \qquad e_\beta = i_\beta (e) \frac{j_\beta^{\rho}}{v_\beta}\,, \quad \beta = c, s1\,. \tag{69}$$

Here $i_c = 1$ for the c pseudofermions and $i_{s1} = -1$ for the $s1$ pseudofermion holes. The concepts of an effective spin projection and an effective charge only apply provided that the β-band Fermi velocity v_β, Eq. (A23) of Appendix A, appearing in the expressions of Eq. (69) is finite.

In the $m \rightarrow 0$ and $m \rightarrow n$ spin-density limits and for electronic density in the range $n \in [0, 1]$ the effective spin projections read,

$$\begin{aligned} s_c^{\sigma_{x3}} &= 0\,; \quad s_{s1}^{\sigma_{x3}} = \frac{1}{2}\,, \quad m \rightarrow 0\,, \\ s_c^{\sigma_{x3}} &= \frac{1}{2}\left\{1 + \frac{v_{s1}}{v_c}\left[1 - \frac{4}{\pi}\arctan\left(\frac{\sin \pi n}{u}\right)\right]\frac{2}{\pi}\arctan\left(\frac{\sin \pi n}{u}\right)\right\}\,, \quad m \rightarrow n\,. \end{aligned} \tag{70}$$

That no expression for $s_c^{\sigma_{x3}}$ is given here for $m \rightarrow n$ results from $v_{s1} \rightarrow 0$ in that limit, so that the concept on a $s1$ pseudofermion-hole effective spin projection does not apply.

Moreover, in the $u \rightarrow 0$ limit the effective spin projections read for the whole densities ranges $n \in [0, 1]$ and $m \in [0, n]$,

$$s_c^{\sigma_{x3}} = 0\,; \qquad s_{s1}^{\sigma_{x3}} = \frac{1}{2}\,, \quad u \rightarrow 0\,. \tag{71}$$

On the other hand, for electronic densities in the range $n \in [0, 1]$ and the spin-density limits $m \rightarrow 0$ and $m \rightarrow n$ the effective charges are given by,

$$\begin{aligned} e_c &= -e\,\xi_0^2\,; \quad e_{s1} = -(e)\,\frac{v_c}{v_{s1}}\frac{\xi_0^2}{2}\,, \quad m \rightarrow 0\,, \\ e_c &= -e\left\{1 + \frac{v_{s1}}{v_c}\left[\frac{2}{\pi}\arctan\left(\frac{\sin \pi n}{u}\right)\right]^2\right\}\,, \quad m \rightarrow n\,. \end{aligned} \tag{72}$$

Again, the concept on a $s1$ pseudofermion-hole effective charge does not apply as $m \rightarrow n$ since $v_{s1} \rightarrow 0$ in that limit.

In the $u \rightarrow 0$ and $u \rightarrow \infty$ limits the effective charge carried by the c pseudofermions is for densities in the ranges $n \in [0, 1]$ and $m \in [0, n]$ given by,

$$\begin{aligned} e_c &= -2e\,, \quad u \rightarrow 0\,, \\ e_c &= -e\,, \quad u \rightarrow \infty\,. \end{aligned} \tag{73}$$

Moreover,, in the $u \rightarrow 0$ limit the effective charge carried by the $s1$ pseudofermion holes reads for such density ranges,

$$e_{s1} = -(e)\,, \quad u \rightarrow 0\,, \tag{74}$$

whereas for electronic densities in the range $n \in [0, 1]$ it has the following value for the $m \rightarrow 0$ and $u \gg 1$ limits,

$$e_{s1} = -(e)\,8\,n\,u\left(\frac{\sin \pi n}{2\pi n - \sin 2\pi n}\right)\,, \quad m \rightarrow 0\,,\ u \gg 1\,. \tag{75}$$

Since $v_{s1} \rightarrow 0$ as $u \rightarrow \infty$ the effective charge expression given here for e_{s1} does not apply in that limit. That it is very large for $u \gg 1$ just results from the smallness of $v_{s1} = \frac{\pi t}{2u}\left(1 - \frac{\sin(2\pi n)}{2\pi n}\right)$.

The above elementary spin and charge currents and corresponding effective spin projections and effective charges are valid within the class of low-energy excited states considered here. Such excited states have no 2ν-η-spinon composite $\eta\nu$ pseudofermions and no unbound η-spinons of η-spin projection $-1/2$. Hence their charge degrees of freedom are mainly described by the c pseudofermion occupancy configurations. The effective spin-projection values $s_c^{\sigma_{x3}} = 0$ and $s_{s1}^{\sigma_{x3}} = 1/2$ given in Eq. (70) for electronic density in the range $n \in [0, 1]$ and spin density $m \rightarrow 0$ are those consistent with the low-energy charge - spin separation occurring in that spin-density limit. However, concerning charge transport, such a charge - spin separation is a concept that should be carefully revisited. Indeed, as given in Eq. (67), one has that $j_c^\rho = v_c\,\xi_0^2$ and $j_{s1}^\rho = v_c\,\xi_0^2/2$ where $\xi_0^2 \rightarrow 2$ for $u \rightarrow 0$ and $\xi_0^2 \rightarrow 1$ for $u \rightarrow \infty$. Note however that although the $s1$ pseudofermion holes carry with them an effective charge current, $-j_{s1}^\rho = -v_c\,\xi_0^2/2$, the vanishing spin-density charge - spin separation is in part perserved. The reason is that both the c pseudofermion elementary charge current $j_c^\rho = v_c\,\xi_0^2$ and $s1$ pseudofermion hole elementary charge current $-j_{s1}^\rho = -v_c\,\xi_0^2/2$ propagate with the c pseudofermion velocities $\pm v_c$. Hence in the $m \rightarrow 0$ limit the charge currents do not propagate with the $s1$ pseudofermion velocities $\pm v_{s1}$. That all elementary charge currents propagate with the c pseudofermion velocities $\pm v_c$ masks the role played by the $s1$ pseudofermion holes in the transport of charge. Only a careful and suitable analysis of the problem in terms of the elementary effective transport charges as that performed here does reveal that in the $m \rightarrow 0$ limit there are indeed finite charge currents carried by the $s1$ pseudofermion holes.

That, as given in Eq. (73), for the densities ranges $n \in [0,1]$ and $m \in [0,n]$ the effective charge carried by the c pseudofermions equals as $u \rightarrow \infty$ the electronic charge is consistent within that limit the rotated electrons becoming electrons. The bare c pseudofermions describe the charge degrees of freedom of the rotated electrons that singly occupied sites. That electron single and double occupancy become good quantum numbers as $u \rightarrow \infty$ imposes that the effective charge carried by a c pseudofermion must equal the electronic charge e in that limit. On the other hand, for finite u values single and double occupancy are good quantum numbers only for the rotated electrons. Upon decreasing the u value a rotated-electron singly occupied site has increasing contributions from electron doubly-occupied sites. This is why upon decreasing the u value the effective charge carried by a c pseudofermion increases for all densities in the ranges $n \in [0,1]$ and $m \in [0,n]$ from e for $u \rightarrow \infty$ to $2e$ as $u \rightarrow 0$. However, that $e_c \rightarrow 2e$ as $u \rightarrow 0$ refers only to c pseudofermions near the c Fermi points and thus to low-energy rotated-electron contributions.

The low-energy one-electron excitations are an example of the type of excited states considered here. For the electronic density range $n \in [0,1]$ and the $m \rightarrow n$ spin-density limit one finds that the non-interacting one-electron spectrum of the fully polarized state is recovered by the energy dispersion of the c pseudofermions. As given in the first paper, in that limit it reads $\varepsilon_c(q) = -2t[\cos q - \cos 2k_F]$. That is consistent with such objects carrying an effective spin projection 1/2 in that limit, as given in Eq. (70). In it the spin-up rotated-electron single occupancy becomes a good quantum number so that the on-site repulsion does not play any role and a full charge - spin recombination is achieved. Upon increasing the spin density from $m = 0$ to $m = n$, the c pseudofermion effective spin projection smoothly changes from $s_c^{\sigma_{x3}} = 0$ to $s_c^{\sigma_{x3}} = 1/2$.

All the above reported exotic behaviors concerning the effective charges and effective spin projections carried with the objects of the present operator formulation are fully controlled by the events of the pseudofermion scattering theory. Indeed, in addition to such objects group velocities, the expressions of the corresponding elementary charge and spin currents involve only two-pseudofermion phase shifts.

C. More general states and the effects of the spin transverse fluctuations onto the spin currents

Next we consider a spin-singlet ground state with electronic density in the range $n \in [0,1]$ and vanishing spin density, $m = 0$. Such a state has no unbound spinons and no $s1$ fermion holes, $M_s^{un} = 0$ and $N_{s1}^h = 0$, respectively. Its object numbers are,

$$\begin{aligned} M_s &= M_s^{bo} = N = n\,N_a\,; \quad M_s^{bo} = N\,; \quad M_{s,\pm 1/2}^{un} = 0\,, \\ M_\eta &= M_\eta^{un} = M_{\eta,+1/2}^{un} = N_a - N = (1-n)\,N_a\,, \\ N_c &= N\,; \quad N_{s1} = N/2\,; \quad N_{s1}^h = 0\,. \end{aligned} \tag{76}$$

For simplicity, we consider that here $N/2$ is an odd integer number, so that $N_c = N$ and $N_{s1} = N/2$ are even and odd integer numbers respectively. For such an initial ground state both the spin and charge currents defined in Eq. (59) vanish. It is a particular case of the ground state whose numbers are provided in Eq. (60), for which all $s1$ band momenta $q \in [-q_{Fs1}, +q_{Fs1}] = [-k_F, +k_F]$ are occupied. (Here $k_F = \pi\,n/2$.) On the other hand, the c band has the same occupancies as that ground state. The $\mu = 0$, $n = 1$, and $m = 0$ absolute ground state is a particular case of the $m = 0$ ground state for which the number values of Eq. (76) apply with $n = 1$ and thus $N = N_a$. For the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state the c band is full and there are no unbound η-spinons.

As discussed in the first paper, the present $m = 0$ ground state has a branch of spin-triplet excited states and another of $N_{s1}^h = 2$ spin-singlet excited states whose energy spectra are degenerate. The number deviations of such spin-triplet excited states are given by,

$$\delta N_c = -\delta M_{\eta,+1/2}^{un} = 0\,; \quad \delta N_{s1} = -1\,; \quad \delta N_{s1}^h = 2\,; \quad \delta M_s^{un} = +2\,. \tag{77}$$

On the other hand, those of the spin-singlet excited states read,

$$\delta N_c = -\delta M_{\eta,+1/2}^{un} = 0\,; \quad \delta N_{s1} = -2\,; \quad \delta N_{s1}^h = 2\,; \quad \delta N_{s2} = +1\,. \tag{78}$$

Both such two types of excited states have two $s1$ pseudofermion holes. In contrast to the spin-triplet state, the spin-singlet state has a single four-spinon composite $s2$ pseudofermion.

The energy spectrum of the spin-triplet excited states of Eq. (77) and spin-singlet excited states of Eq. (78) is plotted in the first paper for $U/t = 4.9$ and two electronic densities, $n = 0.59$ and $n = 1.00$. In contrast to the class of low-energy excited states considered above in Section V B, whose $s1$ pseudofermion holes are under the ground-state excited-state transition created or annihilated at the $s1$ Fermi points, here the two emerging $s1$ pseudofermion holes may have any $s1$ band momentum values. The derivation of the spin and charge currents of Eq. (59) is now a more involved technical problem that requires the numerical solution of Eqs. (A2)-(59) of Appendix A.

The spin currents of the spin-triplet excited states of Eq. (77) and spin-singlet excited states of Eq. (78) were studied in Ref. [51] for the case of an initial $n = 1$ and $m = 0$ ground state. The spin-triplet excited states carry a finite spin current, whose spectrum is plotted in that reference. Alike in the case of the simpler low-energy excited states involving only $\beta = c, s1$ Fermi-point deviations considered in Section V B, such a finite spin current can be written as a sum of elementary spin currents carried by the two $s1$ pseudofermion holes. Also similarly to such low-energy excited states, the two $s1$ pseudofermion holes are found in Ref. [51] to carry with them an effective spin projection 1/2. On the other hand, in spite of having the same energy spectrum as the spin-triplet excited states, the studies of that reference revealed that the spin-singlet excited states carry no spin current. In contrast to the spin-triplet states, the former states have one spin-neutral four-spinon $s2$ pseudofermion, which couples to the vector potential. Such a coupling is described by the term $-4([\phi_\uparrow - \phi_\downarrow]/2N_a)$ added to the discrete momentum q_j on the right-hand side of Eq. (A3) of Appendix A for the $\alpha\nu = s2$ branch. Its role is to exactly cancel the spin currents carried by the two $s1$ pseudofermion holes, so that the spin-singlet state carries no spin current. Similarly and as confirmed in the following, in the case of the non-LWSs of a spin $SU(2)$ tower of states the unbound spinons couple to the vector potential, so that the spin current carried by the corresponding spin LWS Bethe state is changed.

The thermodynamic BA equations, Eqs. (A2)-(A3) of Appendix A, refer only to Bethe states. For $\phi_\uparrow = \phi_\downarrow = 0$ these equations can be extended to non-LWSs of form given in Eq. (A6) of that appendix. This is fulfilled by formally setting some of the rapidities $\Lambda_{\eta\nu}$ and $\Lambda_{s\nu}$ in such equations equal to infinity [3]. For the present LWS BA representation, the resulting equations describe LWSs and non-LWSs with $S_\alpha^{x_3} \leq 0$ for $\alpha = \eta, s$. (A corresponding HWS BA representation refers to HWSs and non-HWSs with $S_\alpha^{x_3} \geq 0$ for $\alpha = \eta, s$.) The spin ($\alpha = s$) or charge ($\alpha = s$) current expectation value of a $S_\alpha^{x_3} \leq 0$ state with unbound-spinon or unbound-η-spinon numbers, respectively, $M_\alpha^{un} = 2S_\alpha$ and $M_{\alpha,-1/2}^{un} = n_\alpha \leq S_\alpha$ is exactly minus that of the corresponding $S_\alpha^{x_3} \geq 0$ state belonging to the same $SU(2)$ tower and having numbers $M_\alpha^{un} = 2S_\alpha$ and $M_{\alpha,-1/2}^{un} = 2S_\alpha - n_\alpha \geq S_\alpha$. Hence without loss in generality one can consider only the thermodynamic BA equations for the LWSs and non-LWSs with $S_\alpha^{x_3} \leq 0$ for $\alpha = \eta, s$.

For such states it is useful to write the unbound-spinon ($\alpha = s$) or unbound-η-spinon ($\alpha = s$) number as $M_\alpha^{un} = 2(S_\alpha + S_\alpha^{x_3}) - 2S_\alpha^{x_3}$. Here the number $(S_\alpha + S_\alpha^{x_3})$ obeys the inequality $0 \leq (S_\alpha + S_\alpha^{x_3}) \leq S_\alpha$. It is the number of pairs of $+1/2$ and $-1/2$ unbound spinons ($\alpha = s$) or $+1/2$ and $-1/2$ unbound η-spinons ($\alpha = s$). On the other hand, the number $-2S_\alpha^{x_3}$ obeys the inequality $0 \leq -2S_\alpha^{x_3} \leq 2S_\alpha$. It is the number of unpaired $+1/2$ unbound spinons ($\alpha = s$) or $+1/2$ unbound η-spinons ($\alpha = s$) left over.

In the general thermodynamic BA equations for both the LWSs and non-LWSs with $S_\alpha^{x_3} \leq 0$, there emerge vector-potential coupling contributions of the form $2(S_\alpha + S_\alpha^{x_3})([\phi_\uparrow - e^{i\pi\delta_{\alpha,\eta}}\,\phi_\downarrow]/2N_a)$ for $\alpha = \eta, s$. They refer to rapidity values $\Lambda_{\alpha\nu} = \infty$ where $\nu = (S_\alpha + S_\alpha^{x_3})$ is now the number of $+1/2$ and $-1/2$ unbound-spinons pairs ($\alpha = s$) or $+1/2$ and $-1/2$ unbound-η-spinons pairs ($\alpha = s$). Such a vector-potential coupling has a form similar to that appearing in Eq. (A3) Appendix A, $2\nu([\phi_\uparrow - e^{i\pi\delta_{\alpha,\eta}}\,\phi_\downarrow]/2N_a)$. The latter refers to the composite $\alpha\nu$ pseudofermions and corresponding ν spin-neutral ($\alpha = s$) and η-spin-neutral ($\alpha = \eta$) spinon and η-spinon pairs, respectively. The Bethe states are LWSs, so that for them $(S_\alpha + S_\alpha^{x_3}) = 0$ and $-2S_\alpha^{x_3} = 2S_\alpha = M^{un}$ for $\alpha = \eta, s$. That they refer to $(S_\alpha + S_\alpha^{x_3}) = 0$ LWSs justifies why the thermodynamic BA equations, Eqs. (A2)-(A4) of Appendix A, do not include explicit vector-potential coupling terms associated with the unbound spinons ($\alpha = s$) and unbound η-spinon ($\alpha = \eta$).

The solution of the extended thermodynamic BA equations for both the LWSs and non-LWSs is technically a quite involved problem. Fortunately, there is an alternative exact way to handle that issue. Indeed, the effects on the Bethe-states spin currents of the transverse spin fluctuations generated by the unbound-spinon spin-flip processes can be exactly accounted for by use of a suitable symmetry operator algebra. The related problem of the effects on the Bethe-states charge currents of the transverse η-spin fluctuations generated by the unbound-η-spinon η-spin-flip processes was addressed in Ref. [50]. Here we apply the same method to the spin currents.

The expectation values of the spin current operator of the non-LWSs of a multiplet-spin $SU(2)$ tower of states can be written as,

$$J^{\sigma_{x_3}} = \frac{1}{\mathcal{C}_s}\,\langle\Psi_{l_o,l_\triangle^0,u}|(\hat{S}_s^-)^{M_{s,-1/2}^{un}}\,\hat{J}^{\sigma_{x_3}}\,(\hat{S}_s^+)^{M_{s,-1/2}^{un}}|\Psi_{l_o,l_\triangle^0,u}\rangle\,. \qquad (79)$$

where the normalization constant $\mathcal{C}_s$ is that given in Eq. (A7) of Appendix A for $\alpha = s$. One can then systematically use the commutators of the off-diagonal generators of the spin $SU(2)$ symmetry algebra, $\hat{S}_s^+$ and $\hat{S}_s^-$, given in the first

S_s	$M_{s,-1/2}=0$	$M_{s,-1/2}=1$	$M_{s,-1/2}=2$	$M_{s,-1/2}=3$	$M_{s,-1/2}=4$	$M_{s,-1/2}=5$	$M_{s,-1/2}=6$	$M_{s,-1/2}=7$
1/2	1	-1	-	-	-	-	-	-
1	1	0	-1	-	-	-	-	-
3/2	1	1/3	-1/3	-1	-	-	-	-
2	1	1/2	0	-1/2	-1	-	-	-
5/2	1	3/5	1/5	-1/5	-3/5	-1	-	-
3	1	2/3	1/3	0	-1/3	-2/3	-1	-
7/2	1	5/7	9/16	1/7	-1/7	-9/6	-5/7	-1

TABLE I: The spin current ratio $J^{\sigma_{x3}}/J_0^{\sigma_{x3}} = C(M_s^{un}, M_{s,-1/2}^{un})$ as obtained from the use of Eqs. (84)-(83) for the spin-tower states of spin up to $S_s = 7/2$.

paper and spin-current operator $\hat{J}^{\sigma_{x3}}$ defined in Eq. (58) to express the spin current $J^{\sigma_{x3}}$ of Eq. (79) in terms of the spin current $J_0^{\sigma_{x3}}$, Eq. (59), of the corresponding spin-tower LWS. Specifically, this involves the use of the following commutators,

$$\left[\hat{J}^{\sigma_{x3}}, \hat{S}_s^{\pm}\right] = \left[\hat{S}_s^{x_3}, \hat{J}^{\pm}\right] = \hat{J}^{\pm}\,; \qquad \left[\hat{J}^{\pm}, \hat{S}_s^{\pm}\right] = \pm 2\hat{J}^{\sigma_{x3}}\,. \tag{80}$$

In addition to the three generators of the spin $SU(2)$ symmetry algebra $\hat{S}_s^{x_3}$, $\hat{S}_s^{+}$, and $\hat{S}_s$ given in the first paper and the spin current operator $\hat{J}^{\sigma_{x3}}$, Eq. (58), the commutators of Eq. (80) involve the tranverse-spin current operators,

$$\begin{aligned} \hat{J}^{+} &= \left(\frac{1}{2}\right) i\,2t \sum_{j=1}^{N_a} \left[c_{j,\downarrow}^{\dagger}\, c_{j+1,\uparrow} - c_{j+1,\downarrow}^{\dagger}\, c_{j,\uparrow}\right]\,, \\ \hat{J}^{-} &= (\hat{J}^{+})^{\dagger} = \left(\frac{1}{2}\right) i\,2t \sum_{j=1}^{N_a} \left[c_{j,\uparrow}^{\dagger}\, c_{j+1,\downarrow} - c_{j+1,\uparrow}^{\dagger}\, c_{j,\downarrow}\right]\,. \end{aligned} \tag{81}$$

After a suitable operator algebra involving commutator manipulations, the spin current $J^{\sigma_{x3}}$ of Eq. (79) carried by a spin-S_s non-LWS with $M_s^{un} = 2S_s$ unbound spinons and $M_{s,-1/2}^{un} \leq 2S_s$ unbound spinons of spin projection $-1/2$ is expressed in terms of the spin current $J_0^{\sigma_{x3}}$ of the corresponding spin-S_s LWS Bethe state with $M_s^{un} = M_{s,+1/2}^{un} = 2S_s$ unbound spinons and no unbound spinons of spin projection $-1/2$ as follows,

$$J^{\sigma_{x3}} = C(M_s^{un}, M_{s,-1/2}^{un})\, J_0^{\sigma_{x3}}\,. \tag{82}$$

The coefficient $C(l,\tilde{l})$ obeys the exact relation,

$$C(l,\tilde{l}) = -C(l, 2S_s - \tilde{l})\,, \quad \tilde{l} \leq S_s\,. \tag{83}$$

Here we have denoted M_s^{un} and $M_{s,-1/2}^{un}$ by l and $\tilde{l}$, respectively. The spin current $J_0^{\sigma_{x3}}$ of the LWS Bethe state appearing on the right-hand side of Eq. (82) is obtained by solution of Eqs. (A2)-(59) of Appendix A. The coefficient $C(M_s^{un}, M_{s,-1/2}^{un})$ is derived for any value of $M_{s,-1/2}^{un} \leq 2S_s$ by the above-mentioned commutator manipulations. The general expression of that coefficient for $M_{s,-1/2}^{un} = 1,2,3$ and any spin value S_s such that $M_s^{un} = 2S_s \geq M_{s,-1/2}^{un}$ (for $M_{s,-1/2}^{un} > 3$ it becomes too cumbersome) is,

$$\begin{aligned} C(l,\tilde{l}) &= \frac{\prod_{j=1}^{\tilde{l}}[\,jl - 2^j\,] - (\tilde{l}-1)(2[(l+1) - (\tilde{l}-1)])^{\tilde{l}-1} - (1-\delta_{\tilde{l},1})(l\tilde{l} - 2^{\tilde{l}})(\tilde{l}-2)(2[(l+1) - (\tilde{l}-2)])^{\tilde{l}-2}}{[\tilde{l}!]\prod_{j=1}^{\tilde{l}}[\,l+1-j\,]}\,, \\ \tilde{l} &= 1,2,3\,, \quad l \geq \tilde{l}\,; \quad l \equiv M_s^{un}\,, \quad \tilde{l} \equiv M_{s,-1/2}^{un}\,. \end{aligned} \tag{84}$$

The use of Eqs. (82)-(84) provides the spin currents $J^{\sigma_{x3}}$ of all states with spin $S_s \leq 7/2$ and thus $M_s^{un} \leq 7$ unbound spinons and $M_{s,-1/2}^{un} \leq 7$ unbound spinons of spin projection $-1/2$. The corresponding spin-current ratios $J^{\sigma_{x3}}/J_0^{\sigma_{x3}} = C(M_s^{un}, M_{s,-1/2}^{un})$ are given in Table I. For instance, the number of $s1$ pseudofermion holes N_{s1}^h of excited states with no $s\nu$ pseudofermions with $\nu > 1$ spinon pairs of the $m = 0$ ground state whose numbers are given in Eq. (76) is given by $N_{s1}^h = M_s^{un}$. The spin of such excited states thus is $S_s = M_s^{un}/2 = N_{s1}^h/2$. The spin current ratios

$J^{\sigma_{x3}}/J_0^{\sigma_{x3}} = C(M_s^{un}, M_{s,-1/2}^{un})$ provided in Table I apply to such excited states with up to $N_{s1}^h = 7$ $s1$ pseudofermion holes.

The excited Bethe states whose $N_{s1}^h \leq 7$ $s1$ pseudofermion holes emerge under the transition from the $m = 0$ ground state in the vicinity of the $s1$ Fermi points carry with them a spin current $J_0^{\sigma_{x3}}$ of general form given in Eqs. (61) and (62). The maximum absolute value of the deviation $\delta 2J_\beta^F$ appearing in Eq. (62) occurs when all $N_{s1}^h \leq 7$ $s1$ pseudofermion holes emerge at the vicinity of one of the two $\iota = \pm 1$ $s1$ Fermi points, so that $\delta 2J_\beta^F = -\iota\, N_{s1}^h$. The spin currents carried by the corresponding $2S_s = M_s^{un} \leq 7$ non-LWSs generated by spin-flipping the $M_s^{un} \leq 7$ unbound spinons of spin projection 1/2 of the Bethe state are then given by $J^{\sigma_{x3}} = C(M_s^{un}, M_{s,-1/2}^{un})\, J_0^{\sigma_{x3}}$. Here the spin current $J_0^{\sigma_{x3}}$ has the form provided in Eqs. (61) and (62) and the magnitudes of the coefficient $C(M_s^{un}, M_{s,-1/2}^{un}) = J^{\sigma_{x3}}/J_0^{\sigma_{x3}}$ are given in Table I.

As reported in Section III A, the phase shifts of Refs. [47, 48] correspond to a particular case of the general elementary two-pseudofermion phase shifts studied in this paper. The related studies of Ref. [52] used a completely different yet equivalent method to derive the spin currents of Eqs. (61) and (62) and charge currents of Eqs. (65) and (66) of the low-energy excited states considered above. Rather than the Hubbard model in a vector potential, such investigations used charge and spin conservation laws to access the corresponding charge and spin current expectation values, respectively. Our present results correct those of that reference concerning the elementary charge current j_{s1}^ρ carried by the $s1$ pseudofermions provided in Eq. (66). Since the $s1$ pseudofermions (called s pseudoparticles in Ref. [52]) do not couple directly to charge, the studies of that reference assumed that $j_{s1}^\rho = 0$. However, our alternative method involving the direct use of the 1D Hubbard model in a uniform vector potential reveals that for the low-energy excited states under consideration the elementary charge current j_{s1}^ρ carried by the $s1$ pseudofermions is in general finite, as given in Eq. (66).

Also within our description of the problem in terms of the 1D Hubbard model in a uniform vector potential, the $s1$ pseudofermions and $s1$ pseudofermion holes do not couple directly to charge. Their coupling to the vector potential corresponds to the term $-2([\phi_\uparrow - \phi_\downarrow]/2N_a)$ on the right-hand side of Eq. (A3) Appendix A for the $\alpha\nu = s1$ branch, which adds to the discrete momentum value q_j. Such a term vanishes for the choice $\phi = \phi_\uparrow = \phi_\downarrow$ corresponding to the charge current operator, as given in Eq. (59). Thus the $s1$ pseudofermions and $s1$ pseudofermion holes do not couple directly to charge. Within our analysis involving a uniform vector potential, that the $s1$ pseudofermions and $s1$ pseudofermion holes feel the charge vector potential through their scattering with the c pseudofermions becomes more clear. This effect leads to the $s1$ pseudofermion elementary charge current j_{s1}^ρ provided in Eq. (66).

In summary, while the states spin currents are carried by the $s\nu$ pseudofermion occupancy configurations (and by the c pseudofermion occupancy configurations, provided that $m \neq 0$), the effects on such currents of the spin transverse fluctuations associated with spin-flipping the unbound spinons are fully controlled by such spin-1/2 objects, which exist on their own right. This is consistent with the results of the first paper according to which the $s\nu$ pseudofermions are spin-neutral 2ν-spinon composite objects whose occupancy configurations generate the spin-singlet representations of the spin $SU(2)$ symmetry algebra, whereas the spin-multiplet towers of states are generated by the independent unbound-spinon occupancies.

VI. RELATION TO THE TRADITIONAL SPINONS AND HOLONS AND TO THE FINITE-ENERGY SPECTRAL PROPERTIES

A review of the vast literature about the 1D Hubbard model is not among the goals of this paper. Many studies associate its ground-state excitations with spinons and holons [6, 40–44]. As mentioned in Section I, concerning the holon definition there are in the literature of the model two main different descriptions. For both of them the holons and antiholons refer to two well-defined object branches. (This is in spite of the objects of such branches having a completely different definition for each of the two descriptions.) In this section we often use the word *holon* to generally designate both the holons and antiholons of the descriptions of Refs. [40] and [41, 42]. It will become clear from the context whether the designation *holon* refers to holons and antiholons as a whole or rather to the holons, as counterpart to the antiholons. Results complementary to those of this section are presented in Appendices C and D.

It is confirmed here and in Appendix D that the distinct choices of elementary objects of the 1D Hubbard model refer actually to different choices of scattering states allowed by the degeneracy of the excited energy eigenstates that span the subspaces where the alternative descriptions apply. For instance, for finite u values the holons, antiholons, and spinons as defined in Ref. [40] are found to correspond to a scattering state choice valid in the subspaces spanned by the excited states of a $S_s = 0$ ground state with arbitrary electronic density n. Furthermore, for finite values of u the holons, antiholons, and spinons as defined in Refs. [41–43] are found to refer to a scattering state choice that holds in the subspace spanned by the excited states of the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state. As discussed in Appendix D, only in the $u \to \infty$ limit the full degeneracy of both η-spin and spin configurations renders the holons, antiholons, and spinons as defined in Ref. [40] and Refs. [41–43] well defined in the subspaces spanned by excited states of ground states with both arbitrary electronic density n and spin density m. On the other hand, the pseudofermion scattering theory is valid for the latter subspaces for all $u > 0$ values. Moreover, the dressed S matrix and phase shifts of the pseudofermion scattering theory are related here and in Appendix C to those of the two traditional spinon and holon descriptions. Concerning the physical role of such dressed matrices and phase shifts, that of the pseudofermion scattering theory is confirmed below to control the one- and two-electron spectral-weight distributions. Its relation to the PDT of Refs. [7, 8] is clarified.

A. Relation to the traditional spinon and holon representations

Within both the description of Andrei and Lowenstein for the 1D Gross-Neveu model [38] and that of Faddeev and Takhtajan for the 1D spin-1/2 isotropic Heisenberg antiferromagnetic model [39], the spin-1/2 color spinors and spin-1/2 spin waves, respectively, are constructed inherently to be excitations of such models $S_s = 0$ absolute ground state. Similarly, the traditional spinons and holons of Refs. [6, 40–43] are also constructed inherently to be excitations of a $S_s = 0$ ground state of the 1D Hubbard model. For both traditional descriptions, spinons are spin-1/2 excitations with no charge. In addition they have been constructed inherently to carry the momentum values of the "holes" that emerge under a transition from a $S_s = 0$ ground state to an excited state in a sequence of BA spin quantum numbers, which within the present notation refers to the $s1$ momentum band.

For one of such ground-state normal-ordered descriptions, of which that of Ref. [40] is a representative, holons and antiholons correspond within the present formulation to the c pseudofermion holes and c pseudofermions, respectively, created under the transitions from the ground state to the excited states. On the other hand, the present formulation accounts for all c pseudofermion holes and c pseudofermions. This includes both those that pre-exist in the ground state and are created under the transitions from it to the excited states, respectively. Alike the c pseudofermion holes and c pseudofermions, the traditional holons and antiholons of Ref. [40] have no internal degrees of freedom associated with the η-spin $SU(2)$ symmetry algebra. They populate excited states of ground states with arbitrary values of the electronic density n and zero spin density, $m = 0$. The spin-1/2 spinons, holons, and antiholons of Ref. [40] are excitations of such a ground state.

On the other hand, for the alternative also ground-state normal-ordered traditional description, of which that of Refs. [41–43] is a representative, holons and antiholons are η-spin-1/2 objects with $+1/2$ and $-1/2$ η-spin projection, respectively, created under the transitions from the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state to its excited states. Such holons and antiholons have been constructed inherently to carry the rapidity momentum values of the "holes" that emerge in the $k'_j s$ BA distribution of Ref. [2] under such transitions. These rapidity momentum values $k'_j s$ refer to the momentum values q_j of the holes that within the notation used in this paper emerge in the c momentum band. For rotated electrons as defined in the first paper, the holons and antiholons of Refs. [41–43] contain both the η-spin $SU(2)$ symmetry and c hidden $U(1)$ symmetry degrees of freedom of the rotated-electron unoccupied and doubly-occupied sites created under the above-mentioned transitions. The spin-1/2 spinons and η-spin-1/2 holons are excitations of the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state.

Within the operator formulation introduced in the first paper, all model energy eigenstates involve occupancy configurations of $M_s = 2S_c$ 1/2-spin spinons and $M_\eta = [N_a - 2S_c]$ η-spin-1/2 η-spinons. The number values of such objects may be rewritten as $M_s = [2S_s + 2\sum_{\nu=1}^{\infty} \nu\, N_{s\nu}]$ and $M_\eta = [2S_\eta + 2\sum_{\nu=1}^{\infty} \nu\, N_{\eta\nu}]$, respectively. Furthermore, such states involve a c momentum band with $N_c = 2S_c = M_s$ c fermions and $N_c^h = [N_a - 2S_c] = M_\eta$ c fermion holes. The corresponding spinon and η-spinon occupancy configurations generate representations of the spin $SU(2)$ symmetry and η-spin $SU(2)$ symmetry algebras, respectively.

The spinons refer to a normal-ordered description relative to the $S_\eta = N_a/2; S_s = 0; 2S_c = 0$ electron and rotated-electron vacuum. The number of spinons vanishes for that vacuum whereas for it the number of η-spinons reaches its maximum value, $M_\eta = N_a$. Indeed, there is a duality between spinons and η-spinons, the latter referring to a normal-ordered description relative to the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state. That state has no η-spinons and for it the number of spinons reaches its maximum value, $M_s = N_a$. However, since the present operator formulation accounts for the elementary-object occupancy configurations that generate all 4^{N_a} energy eigenstates from the $S_\eta = N_a/2; S_s = 0; 2S_c = 0$ electron and rotated-electron vacuum, globally it is normal-ordered relative to that vacuum. That is in spite of the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state not being populated by η-spinons.

For present elementary-object formulation, the $N_c = 2S_c$ c fermions and the $M_s = 2S_c$ spinons describe the c hidden $U(1)$ symmetry and spin $SU(2)$ symmetry degrees of freedom, respectively, of the same $2S_c$ sites singly occupied by rotated electrons for all energy eigenstates. Similarly, the $N_c^h = [N_a - 2S_c]$ c fermion holes and the $M_\eta = [N_a - 2S_c]$ spinons describe the c hidden $U(1)$ symmetry and η-spin $SU(2)$ symmetry degrees of freedom, respectively, of the same $[N_a - 2S_c]$ sites unoccupied and doubly occupied by rotated electrons. This is why the number of c fermions equals that of spinons and the number of c fermion holes that of η-spinons, respectively. Thus the number of c

fermions vanishes for the $S_\eta = N_a/2; S_s = 0; 2S_c = 0$ electron and rotated-electron vacuum whereas for it the number of c fermion holes reaches its maximum value, $N^h_c = N_a$. Furthermore, the number of c fermion holes vanishes for the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state whereas for it the number of c fermions reaches its maximum value, $N_c = N_a$.

Thus, in contrast to the traditional descriptions of Refs. [40–43], whose holons and spinons are ground-state excitations, the present formulation accounts for the occupancy configurations of its elementary objects that generate the ground states from the $S_\eta = N_a/2; S_s = 0; 2S_c = 0$ electron and rotated-electron vacuum. In general in this paper we consider ground states with densities in the ranges $n \in [0,1]$ and $m \in [0,n]$ since those are spin and η-spin LWSs and thus Bethe states. Within the present operator formulation, those have $M_s = [2N_{s1} + 2S_s] = N$ spinons, $M_\eta = 2S_\eta = [N_a - N]$ η-spinons, $N_c = N$ c fermions, and $N^h_c = [N_a - N]$ c fermion holes. For such states the number of rotated-electron doubly occupied sites vanishes, so that the number of electrons N equals that of rotated electrons that singly occupy sites. Since such ground states are spin and η-spin LWSs, their $M^{un}_s = 2S_s$ unbound spinons have have spin projection $+1/2$ and their $M^{un}_\eta = 2S_\eta$ unbound η-spinons have have η-spin projection $+1/2$. Within the present formulation, spin-flipping and η-spin-fliping, respectively, such unbound objects generates energy eigenstates outside the BA subspace. The ground-state $M_s = N$ spinons refer to the N spin-1/2 spins of the rotated electrons that singly occupy sites. The ground-state $N_c = N$ c fermions describe the charge degrees of freedom of such rotated electrons. The ground-state $M_\eta = [N_a - N]$ η-spinons describe the η-spin degrees of freedom of the $[N_a - N]$ sites unoccupied by rotated electrons. The ground-state $N^h_c = [N_a - N]$ c fermion holes describe the c hidden $U(1)$ symmetry degrees of freedom of the latter $[N_a - N]$ sites ground-state occupancies.

For the description of Ref. [40], which is normal ordered relative to an initial $S_s = 0$ ground state of arbitrary electronic density n, we denote the holon, antiholon, and spinon number values by N^h_h, N^p_h, and N_s, respectively. Since such holons and antiholons are unrelated to η-spin and rather refer to "holes" and "particles" emerging in the BA c momentum band, we use the upper indices h and p, respectively, to label their numbers. For the description of Ref. [42], which is normal ordered relative to the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state, the holon and spinon number values are denoted by L_h and L_s, respectively. For general excited states of the corresponding initial ground state such numbers may be expressed in terms of the numbers of rotated electrons as defined in the first paper. For the initial ground states of both the descriptions of Refs. [40] and [41, 42], the spin degrees of freedom of the N rotated electrons that singly occupy sites are described by $N_{s1} = N/2$ BA spin strings of length $\nu = 1$. (For the initial absolute ground state of Refs. [41, 42], one has that $N = N_a$.) Within the present formulation, that refers to $N_{s1} = N/2$ two-spinon composite $s1$ fermions, which contain all $2N_{s1} = N$ spinons. For the excited states, the number $2N_{s1}$ of spinons that are part of two-spinon composite $s1$ fermions is such that $2N_{s1} \leq N$. Both for the ground states and excited states, the $2N_{s1} \leq N$ spinons of the present formulation are the spins of rotated electrons created by the generator of such states onto the $S_\eta = N_a/2; S_s = 0; 2S_c = 0$ electron and rotated-electron vacuum.

On the other hand, the $2N_{s1} \leq N$ spinons of the present formulation that remain after the transition from a $S_s = 0$ ground state to its excited states are not accounted for by both ground-state normal-ordered traditional holon and spinon descriptions. Consistent, the number value of traditional spinons equals that of rotated-electron singly occupied sites minus $2\sum_{\nu=1}^{\infty} N_{s\nu}$, which gives $N_s = [2S_c - 2\sum_{\nu=1}^{\infty} N_{s\nu}]$ and $L_s = [2S_c - 2\sum_{\nu=1}^{\infty} N_{s\nu}]$ for the descriptions of Refs. [40] and [41, 42], respectively. Although the numbers N_s and L_s have the same expression, that of the latter applies only to the PS of the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state.

The numbers N^h_h of holons and N^p_h of antiholons of Ref. [40] are in terms of the c band momentum distribution function deviations given by,

$$\begin{aligned} N^h_h &= -\frac{N_a}{2\pi}\int_{-2k_F}^{2k_F} dq\,\delta N_c(q) = 0,1,2,\ldots, \\ N^p_h &= \frac{N_a}{2\pi}\left[\int_{-\pi}^{-2k_F} dq\,\delta N_c(q) + \int_{2k_F}^{\pi} dq\,\delta N_c(q)\right] = 0,1,2,\ldots. \end{aligned} \tag{85}$$

Indeed, as mentioned above the holons and antiholons of that reference correspond to the c pseudofermion holes of bare momentum $q \in [-2k_F, 2k_F]$ and c pseudofermions of bare momentum $q \in [-\pi, -2k_F]$ and $q \in [2k_F, \pi]$, respectively, created under a transition from the ground state to the excited state. The number of spinons of Ref. [40] may be rewritten as $N_s = N^h_{s1} = [M_s - 2\sum_{\nu=1}^{\infty} N_{s\nu}] = [2S_s + \sum_{\nu=2}^{\infty} 2(\nu-1)\,N_{s\nu}]$. The corresponding numbers of spin-down and spin-up spinons are then given by $N_{s,-1/2} = [M^{un}_{s,-1/2} + \sum_{\nu=2}^{\infty}(\nu-1)\,N_{s\nu}]$ and $N_{s,+1/2} = [M^{un}_{s,+1/2} + \sum_{\nu=2}^{\infty}(\nu-1)\,N_{s\nu}]$, respectively.

Moreover, for the description of Refs. [41, 42] the number of holons value equals that of the number of rotated-electron doubly occupied plus unoccupied sites, $L_h = [N_a - 2S_c]$. (Both such numbers values vanish for the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state.) The numbers of traditional holons and spinons of that reference may be rewritten as, $L_h = N^h_c = M_\eta = [2S_\eta + \sum_{\nu=1}^{\infty} 2\nu\,N_{\eta\nu}]$ and $L_s = N^h_{s1} = [M_s - 2\sum_{\nu=1}^{\infty} N_{s\nu}] = [2S_s + \sum_{\nu=2}^{\infty} 2(\nu - 1)\,N_{s\nu}]$, respectively. The corresponding numbers of antiholons and holons as defined in that reference are $L_{h,-1/2} =$

$[M^{un}_{\eta,-1/2}+\sum_{\nu=1}^{\infty}\nu\,N_{\eta\nu}]$ and $L_{h,+1/2} = [M^{un}_{\eta,+1/2}+\sum_{\nu=1}^{\infty}\nu\,N_{\eta\nu}]$, respectively. Here $-1/2$ and $+1/2$ refer to their η-spin projections. Furthermore, the down-spin and up-spin spinon numbers are $L_{s,-1/2} = [M^{un}_{s,-1/2} + \sum_{\nu=2}^{\infty}(\nu-1)\,N_{s\nu}]$ and $L_{s,+1/2} = [M^{un}_{s,+1/2} + \sum_{\nu=2}^{\infty}(\nu-1)\,N_{s\nu}]$, respectively.

Let us consider as an example both the spin-triplet $S_s = 1$ excited states and simplest $S_s = 0$ spin-singlet excited states with one BA spin-string of length $\nu = 2$ of any $S_s = 0$ ground state, which are populated by two traditional spinons. Specifically, within the descriptions of Refs. [40–42] such spin-singlet and three spin-triplet excited states involve two traditional spinons of spin 1/2 in a spin-singlet configuration and in the three spin-triplet configurations, respectively. In contrast, within the present object formulation both such spin-triplet and spin-singlet energy eigenstates are N-spinon states. Each spin-triplet $S_s = 1$ excited state has two unbound spinons, $M^{un}_s = 2S_s = 2$, whose occupancies generate the corresponding three spin-triplet states. In addition, each such a state contains $M^{bo}_s = 2N_{s1} = [N-2]$ bound spinons within $N_{s1} = [N/2-1]$ two-spinon spin-neutral configurations. Furthermore, the spin-singlet $S_s = 0$ excited states have no unbound spinons. They are populated by $M^{bo}_s = [2N_{s1} + 4N_{s2}] = N$ bound spinons within $N_{s1} = [N/2-2]$ two-spinon and $N_{s2} = 1$ four-spinon spin-neutral configurations. That each of the above energy eigenstates has different number values of spinons of either traditional description and of the present-formulation spinons confirms that the former and latter spinons are completely different objects.

In the first paper it was found that out of the $M_s = 2S_c$ spinons, $M^{un}_s = 2S_s$ are unbound spinons, which is consistent with the expressions of their numbers provided in Eq. (3) determining the state spin value S_s. The remaining $M^{bo}_s = [2S_c - 2S_s] = \sum_{\nu=1}^{\infty} 2\nu N_{s\nu}$ bound spinons participate in spin-neutral configurations within 2ν-spinon $s\nu$ fermions. The results of that paper have shown that both the configurations of the N_{s1} $s1$ fermions and corresponding $N^h_{s1} = [2S_s + \sum_{\nu=2}^{\infty} 2(\nu-1)\,N_{s\nu}]$ $s1$ fermion holes have a spin-singlet character. On the other hand, the N_s (and L_s) traditional spinons as defined in Ref. [40] (and Refs. [41, 42]) both have spin 1/2 and carry the BA $s1$ band momentum values q_j carried within the present formulation by the $s1$ fermion holes. Thus straightforward arguments that follow from the spin $SU(2)$ symmetry algebra imply that since a state of spin S_s has $N_s = [2S_s + \sum_{\nu=2}^{\infty} 2(\nu-1)\,N_{s\nu}]$ (and $L_s = [2S_s + \sum_{\nu=2}^{\infty} 2(\nu-1)\,N_{s\nu}]$) traditional spinons, $2S_s$ of those must be in spin-multiplet configurations and the remaining $\sum_{\nu=2}^{\infty} 2(\nu-1)\,N_{s\nu}$ in spin-neutral configurations. This is fully consistent with the above number expressions $N_{s,-1/2} = [M^{un}_{s,-1/2} + \sum_{\nu=2}^{\infty}(\nu-1)\,N_{s\nu}]$ and $N_{s,+1/2} = [M^{un}_{s,+1/2} + \sum_{\nu=2}^{\infty}(\nu-1)\,N_{s\nu}]$ (and $L_{s,-1/2} = [M^{un}_{s,-1/2} + \sum_{\nu=2}^{\infty}(\nu-1)\,N_{s\nu}]$ and $L_{s,+1/2} = [M^{un}_{s,+1/2} + \sum_{\nu=2}^{\infty}(\nu-1)\,N_{s\nu}]$.)

That the N_s (and L_s) traditional spinons have been constructed inherently to being excitations of a $S_s = 0$ ground state (and the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state) justifies that $N_s \leq M_s = [N_s + 2\sum_{\nu=1}^{\infty} N_{s\nu}]$ (and $L_s \leq M_s = [L_s + 2\sum_{\nu=1}^{\infty} N_{s\nu}]$.) Note however that the number of traditional spinons participating in spin-multiplet configurations equals that of unbound spinons of the present formulation, $M^{un}_s = 2S_s$. Thus the difference in the number of spinons refers to those participating in spin-neutral configurations. Specifically, the number values of such spinons are $M^{bo}_s = [2S_c - 2S_s] = \sum_{\nu=1}^{\infty} 2\nu N_{s\nu}$ and $[N_s - 2S_s] = [M^{bo}_s - 2\sum_{\nu=1}^{\infty} N_{s\nu}] = \sum_{\nu=2}^{\infty} 2(\nu-1)\,N_{s\nu}$ (and $[L_s - 2S_s] = M^{bo}_s - 2\sum_{\nu=1}^{\infty} N_{s\nu} = \sum_{\nu=2}^{\infty} 2(\nu-1)\,N_{s\nu}$) for the present formulation and the traditional spinon descriptions, respectively.

On the one hand, the N^h_h holons and N^p_h antiholons of the description of Ref. [40], whose number values are given in Eq. (85), correspond to c pseudofermion holes and c pseudofermions, respectively, of the present operator formulation created under a transition to an excited state. This choice is consistent with within the present operator formulation the unbound η-spinons whose occupancies as given in the expressions of their numbers provided in Eq. (3) determine the excited states η-spin number values S_η and $S^{x_3}_\eta$ neither being scatterers nor scattering centers. Thus the holon and antiholon description of Ref. [40] implicitly accounts for the separation of the η-spin $SU(2)$ symmetry degrees of freedom and c hidden $U(1)$ symmetry degrees of freedom, respectively, of the rotated-electron occupancy configurations. Indeed, the results of the first paper reveal that the BA numbers occupancy configurations corresponding to t hose of the holons and antiholons as defined in Ref. [40] generate the representations of the c hidden $U(1)$ symmetry algebra.

On the other hand, the $L_h = [N_a - 2S_c]$ traditional η-spin-1/2 holons of Refs. [41, 42] are different both from the $M_\eta = [N_a - 2S_c]$ η-spin-1/2 η-spinons and $N^h_c = [N_a - 2S_c]$ spin-less and η-spin-less c fermion holes of the current formulation. This is in spite of all such objects having the same number value. The L_h traditional holons as defined in Ref. [42] both have η-spin 1/2 and carry the BA c band momentum values q_j carried here by the c fermion holes. In the first paper it has been shown that the momentum occupancy configurations of the BA c band generate representations of the c hidden $U(1)$ symmetry algebra. Moreover, the traditional holons η-spin 1/2 is associated with the η-spin $SU(2)$ symmetry. Combining these two properties, reveals that the traditional holons of Refs. [41, 42] have been implicitly constructed inherently to describing both η-spin $SU(2)$ symmetry and c hidden $U(1)$ symmetry degrees of freedom of the set of $[N_a - 2S_c]$ rotated-electron doubly and unoccupied sites. (Although the authors of that reference did not explicitly mention the latter symmetry, as confirmed in the first paper the c hidden $U(1)$ symmetry is accounted for by the BA solution, which they have used in their studies.) The general formulation used here and in the first paper accounts for the separation of the latter two degrees of freedom. (That such a formulation number value $M_\eta = [N_a - 2S_c]$ of $1/2$-η-spin η-spinons exactly equals that $N_c^h = [N_a - 2S_c]$ of c fermion holes was shown in the first paper to merely following from the occupancy configurations of such objects describing the η-spin $SU(2)$ symmetry and c hidden $U(1)$ symmetry degrees of freedom, respectively, of the same set of $[N_a - 2S_c]$ rotated-electron doubly and unoccupied sites.)

It was confirmed in that paper that the number value of η-spinon, spinon, and c fermion occupancy configurations in each subspace of fixed η-spin S_η, spin S_s, and $2S_c$ exactly equals that of the model global η-spin $SU(2)$ symmetry, spin $SU(2)$ symmetry, and c hidden $U(1)$ symmetry algebra representations, respectively. This is not required though for the configurations of the traditional spinons and holons of both the descriptions of Refs. [40] and [41, 42]. Indeed, their holon and spinon occupancy configurations refer to ground-state normal-ordered descriptions that account only for the configuration changes relative to those of the initial ground-state occupancies. Thus they do not account for the extra occupancy configurations associated with the generation of that ground state from the $S_\eta = N_a; S_s = 0; 2S_c = 0$ electron and rotated-electron vacuum.

A quantum problem is defined by a Hamiltonian plus the Hilbert space it acts onto. The elementary object descriptions considered in the first paper, here, in Ref. [40], and in Refs. [41–43] refer to four quantum problems, involving the 1D Hubbard model in different subspaces. It is confirmed below that the subspaces where each description scatterers and scattering centers are defined is determined by the corresponding different choices of scattering states being allowed by the degeneracy of the excited energy eigenstates that span such subspaces. The subspaces under consideration are the following:

– The operator formulation introduced in the first paper and used here, which involves the η-spin-$1/2$ η-spinons, spin-$1/2$ spinons, and c fermions along with the related 2ν-η-spinon composite $\eta\nu$ fermions and 2ν-spinon composite $s\nu$ fermions, refers to the 1D Hubbard model in its full Hilbert space, spanned by all 4^{N_a} energy eigenstates.

– The related β pseudofermion operator description associated with the pseudofermion scattering theory introduced in this paper and the related PDT of Refs. [7, 8] refers to the 1D Hubbard model in a PS. We recall that each PS is spanned by an initial ground state of densities in the ranges $n \in [0, 1]$ and $m \in [0, n]$ and the excited states generated from it whose number deviations are such that $\delta N_\beta/N_a \rightarrow 0$, $\delta M_{\eta,-1/2}/N_a \rightarrow 0$, and $\delta M_{s,-1/2}/N_a \rightarrow 0$ as $N_a \rightarrow \infty$ for all $\beta = c, \eta\nu, s\nu$ pseudofermion branches where $\nu = 1, ..., \infty$ and unbound $-1/2$ η-spinons and unbound $-1/2$ spinons.

– The holon and spinon description of Ref. [40] refers to the 1D Hubbard model in any PS spanned by an initial ground state of arbitrary electronic density n and spin density $m = 0$ and the excited states generated from it whose traditional-holon and traditional-antiholon numbers given in Eq. (85) and traditional-spinon numbers are such that $N_h^h/N_a \rightarrow 0$, $N_h^p/N_a \rightarrow 0$, and $N_s/N_a = [2S_c - 2\sum_{\nu=1}^{\infty} N_{s\nu}]/N_a \rightarrow 0$, respectively, as $N_a \rightarrow \infty$.

– Finally, the traditional holon and spinon description of Refs. [41–43] applies to the half-filled 1D Hubbard model in the specific PS generated from the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state. It is spanned by that ground state and excited states generated from it whose number deviations are such that $L_h/N_a = [1 - 2S_c/N_a] \rightarrow 0$ and $L_s/N_a = [2S_c - 2\sum_{\nu=1}^{\infty} N_{s\nu}]/N_a \rightarrow 0$ as $N_a \rightarrow \infty$.

The PS spanned by the excited energy eigenstates of the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state is that common to the three scattering theories. It has an interesting property: The energy contribution from the η-spin degrees of freedom of energy eigenstates with the same number $L_h = [N_a - 2S_c]$ of holons as defined in Ref. [42] is the same. The different energies of such L_h-holon configurations stem only from the c hidden $U(1)$ symmetry degrees of freedom associated with the c band momentum values q_j occupancy configurations.

The PSs spanned by the excited energy eigenstates of any $S_s = 0$ ground state, including the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state, have a similar interesting property: The energy contribution from the spin configurations that involve the unbound spinons and the $s\nu$ fermions with $\nu > 1$ spinons pairs of the present formulation is the same for energy eigenstates with the same number $N_s = [2S_c - 2\sum_{\nu=1}^{\infty} N_{s\nu}]$ (and $L_s = [2S_c - 2\sum_{\nu=1}^{\infty} N_{s\nu}]$) of traditional spinons as defined in Ref. [40] (and in Ref. [42]). The different energies of such N_s-spinon (and L_s-spinon) configurations stem only from the $s1$ band momentum values q_j occupancy configurations.

B. Relation between scattering theories

Here we compare the three scattering theories for the 1D Hubbard model in the $[N_c^h + N_{s1}^h] = 2$ reduced subspace, which is contained in the PS spanned by the excited states of the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state. In addition, we extend our discussion to that PS, which is the subspace common to the three scattering theories under consideration. Moreover, we compare the theories of the present formulation and of Ref. [40] for the 1D Hubbard model in the PSs spanned by the excited states of $S_s = 0$ ground states with electronic density in the range $n \in [0, 1]$.

In Appendix C it is shown that the dressed phase shifts considered in Refs. [41–43] for the excited states of the $[N^h_c + N^h_{s1}] = 2$ reduced subspace are particular cases of the overall phase-shift functionals $Q_c(q)$ and $Q_{s1}(q)$ of Eq. (15). Similar results are obtained for the phase shifts considered in Ref. [40] associated with the excited states of $S_s = 0$ ground states with electronic density in the range $n \in [0, 1]$.

The dressed phase shifts of both Ref. [40] and Refs. [41, 42] were directly extracted from the BA solution. The latter references used the method of Ref. [53], which is equivalent to that used in the studies of Ref. [40]. In the $[N^h_c + N^h_{s1}] = 2$ reduced subspace, the dressed phase shifts refer to twelve types of excited states whose number and number deviation values are given in Tables II-IV of Appendix C. For the three scattering theories, the scattering centers are elementary objects created under the transitions to the excited states. Within the scattering theories of both Ref. [40] and Refs. [41, 42], only such elementary objects play the role of scatterers. The elementary objects of the present formulation whose occupancy configurations remain unchanged under the transitions from the ground state to the excited states are not accounted for by such ground-state normal-ordered scattering theories.

The form of the original spin-1/2 electron bare S matrix is uniquely defined by the BA solution. (See, for instance, Lecture 1 of Ref. [40].) On the other hand, that in Refs. [41, 42] the scatters and scattering centers associated with the dressed phase shifts are identified with spin-1/2 spinons and η-spin-1/2 holons leads within the reduced subspace to a non-diagonal spinon-spinon 4×4 dressed S matrix block and a non-diagonal holon-holon 4×4 dressed S matrix block of an overall 16×16 dressed S matrix given below. The spinons as defined in Ref. [40] also have spin 1/2, so that within the reduced subspace the corresponding spinon-spinon dressed S matrix block is again a 4×4 non-diagonal matrix. However, the holons (and antiholons for electronic densities $n \neq 1$) of the description of that reference have no internal degrees of freedom. Thus except for its spinon-spinon 4×4 block, the dressed S matrix of Ref. [40] given below is otherwise diagonal.

The number values of elementary objects of the present formulation for all classes of excited states that span the reduced subspace are provided in Table II of Appendix C. The corresponding number values of the elementary objects of the traditional holon and spinon descriptions of both Ref. [40] and Refs. [41, 42] are given in Table III of that appendix. The holon-holon 4×4 dressed S matrix block of Refs. [41, 42] (and spinon-spinon 4×4 dressed S matrix block of both Ref. [40] and Refs. [41, 42]) corresponds to subspaces spanned by four types of $L_h = 2; L_s = 0$ η-spin-triplet and η-spin-singlet excited energy eigenstates (and $L_h = 0; L_s = 2$ and thus $N^h_h = 0; N_s = 2$ spin-triplet and spin-singlet excited states). In this case the two objects created under the ground-state - excited-state transitions are traditional holons (and traditional spinons) and thus it is unimportant which of them is chosen as scatterer and scattering center, since the phase shifts are the same. Thus, at fixed c band hole momentum values q_j and $q_{j'}$ of the two created objects the number of relevant one-particle scattering states equals that of excited energy eigenstates.

On the other hand, for each of the four types of $L_h = 1; L_s = 1$ and thus $N^h_h = 1; N_s = 1$ doublet excited energy eigenstates, one must consider two one-particle scattering states. Indeed, both within the descriptions of Refs. [40] and [41, 42] one traditional holon and one traditional spinon are in this case created under the ground-state - excited-state transitions and thus the one-particle scattering states where that traditional holon and traditional spinon is the scatterer are different: The traditional holon and traditional spinon scattering states are associated with different phase shifts that refer to the c and $s1$ phase shifts given in Eq. (C19) of Appendix C, respectively. Therefore, while at fixed values of the c and/or $s1$ band hole momenta q_j and $q_{j'}$ of the two created objects the reduced subspace is spanned by the twelve excited energy eigenstates whose number and number deviation values are given in the Tables II-IV of Appendix C, the dressed S matrix for the corresponding full scattering problem involves sixteen states and thus has indeed dimension 16×16. However, for each pair of one-particle scattering states associated with the same doublet energy eigenstate ones uses the latter state in the evaluation of the corresponding phase shifts that appear in the entries of that dressed S matrix.

For the scattering theory of Refs. [41, 42] it is of the form,

$$\mathbf{S} = \begin{bmatrix} \mathbf{S}_{SS} & 0 & 0 & 0 \\ 0 & \mathbf{S}_{S\eta} & 0 & 0 \\ 0 & 0 & \mathbf{S}_{\eta S} & 0 \\ 0 & 0 & 0 & \mathbf{S}_{CC} \end{bmatrix} , \tag{86}$$

where $\mathbf{S}_{S\eta}$ and $\mathbf{S}_{\eta S}$ are 4×4 diagonal dressed matrix blocks corresponding to scattering events where the traditional spinons and holons are the scatterers and the traditional holons and spinons the scattering centers, respectively. On the other hand, $\mathbf{S}_{SS}$ and $\mathbf{S}_{CC}$ are the above mentioned two 4×4 dressed S matrix blocks associated with the spinon-spinon and holon-holon scattering, respectively. The latter two matrices are not diagonal.

In the present case of the scattering theory of Refs. [41, 42], we denote the two sets of four states that refer to the four one-particle scattering states by $|+1/2, +1/2; \alpha\rangle$, $|-1/2, -1/2; \alpha\rangle$, $|+1/2, -1/2; \alpha\rangle$, and $|-1/2, +1/2; \alpha\rangle$. Here $\alpha = \eta$ and $\alpha = s$ refer to the holon-holon and spinon-spinon states, respectively. The two η-spin ($\alpha = \eta$) or spin ($\alpha = s$) projections $\pm 1/2$ labeling these states are those of the two involved traditional holons or spinons, respectively.

The 4×4 permutation matrix $\mathbf{P}$ transforms the above four states as,

$$\begin{array}{llll} |+1/2,+1/2;\alpha\rangle \;\Longrightarrow\; |+1/2,+1/2;\alpha\rangle\,; & |+1/2,-1/2;\alpha\rangle \;\Longrightarrow\; |-1/2,+1/2;\alpha\rangle\,, \\ |-1/2,+1/2;\alpha\rangle \;\Longrightarrow\; |+1/2,-1/2;\alpha\rangle\,; & |-1/2,-1/2;\alpha\rangle \;\Longrightarrow\; |-1/2,-1/2;\alpha\rangle\,; \quad \alpha=\eta,\,s\,. \end{array} \tag{87}$$

The two non-diagonal dressed S matrix blocks $\mathbf{S}_{SS}$ and $\mathbf{S}_{CC}$ in Eq. (86) are then of the following form,

$$\mathbf{S}_{\beta\beta} = \frac{1}{2}(S_{\beta T}+S_{\beta S})\,\mathbf{I} + \frac{1}{2}(S_{\beta T}-S_{\beta S})\,\mathbf{P}\,; \quad \beta = C,\,S\,, \tag{88}$$

where $\mathbf{I}$ is the 4×4 unity matrix,

$$S_{\beta\tau} = (-1)^{y_\beta}\,e^{i\delta_{\beta\tau}}\,; \qquad \beta = C,S\,; \qquad \tau = T,S\,; \qquad y_C = 1\,, \quad y_S = 0\,. \tag{89}$$

Here $\delta_{\beta\tau}$ with $\beta = C,\,S$ and $\tau = T,\,S$ denotes the four phase shifts defined by Eqs. (C15), (C16), (C17), and (C18) of Appendix C. On the other hand, the four diagonal entries of the above 4×4 diagonal dressed S matrix blocks $\mathbf{S}_{S\eta}$ and $\mathbf{S}_{\eta S}$ are equal and given by,

$$S_{\beta'\beta''} = (-1)^{z_{\beta'}}\,e^{i\delta_{\beta'\beta''}}\,; \qquad \beta'\beta'' = S\eta,\eta S\,; \qquad z_\eta = 1\,, \quad z_S = 0\,, \tag{90}$$

where $\delta_{\beta'\beta''}$ with $\beta'\beta'' = S\eta$, ηS are the two phase shifts defined in Eq. (C19) of Appendix C.

In the case of the scattering theory of Ref. [40], we find for the reduced subspace at fixed values of the two c and/or $s1$ band hole momenta q_j and $q_{j'}$ a 16×16 dressed S matrix, which is related to that given in Eq. (86) by a unitary transformation as follows,

$$\tilde{\mathbf{S}} = \mathbf{R}^\dagger\mathbf{S}\mathbf{R} = \begin{bmatrix} \tilde{\mathbf{S}}_{SS} & 0 & 0 & 0 \\ 0 & \tilde{\mathbf{S}}_{S\eta} & 0 & 0 \\ 0 & 0 & \tilde{\mathbf{S}}_{\eta S} & 0 \\ 0 & 0 & 0 & \tilde{\mathbf{S}}_{CC} \end{bmatrix}\,; \quad \mathbf{R} = \begin{bmatrix} \mathbf{I} & 0 & 0 & 0 \\ 0 & \mathbf{I} & 0 & 0 \\ 0 & 0 & \mathbf{I} & 0 \\ 0 & 0 & 0 & \mathbf{J} \end{bmatrix}\,; \quad \mathbf{J} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1/\sqrt{2} & -1/\sqrt{2} & 0 \\ 0 & 1/\sqrt{2} & 1/\sqrt{2} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}\,. \tag{91}$$

Here $\mathbf{R}$ denotes the corresponding unitary matrix. The only non-diagonal part of this 16×16 dressed S matrix corresponds to the spinon-spinon 4×4 dressed S matrix block $\tilde{\mathbf{S}}_{SS}$, which remains identical to the spinon-spinon 4×4 dressed S matrix block $\mathbf{S}_{SS}$ in the 16×16 dressed S matrix expression given in Eq. (86). It has the form given in Eqs. (88) and (89) for $\beta = S$. The only change relative to the dressed S matrix of Eq. (86) occurs in the 4×4 matrix block $\tilde{\mathbf{S}}_{CC}$ of the 16×16 matrix provided in Eq. (91), which now is diagonal. Its three first diagonal entries (and the fourth diagonal entry) equal the phase factor S_{CT} (and equals the phase factor S_{CS}) defined in Eq. (89).

For the scattering theory of Ref. [40], we denote again the set of four states that refer to the four one-particle scattering states by $|+1/2,+1/2;s\rangle$, $|-1/2,-1/2;s\rangle$, $|+1/2,-1/2;s\rangle$, and $|-1/2,+1/2;s\rangle$. Those are associated with the spinon-spinon scattering. The two spin projections $\pm 1/2$ labeling these states are those of the two involved traditional spinons. The 4×4 permutation matrix $\mathbf{P}$ that transforms such four states is given in Eq. (87) for $\alpha = s$. The two phase shifts of Eqs. (C17) and (C18) of Appendix C, which are the quantities $\delta_{\beta\tau}$ appearing in Eq. (89) for $\beta = S$ and $\tau = T,\,S$, refer to the half-filling limit of corresponding spinon phase shifts of Ref. [40] for spin-triplet ($\tau = T$) and spin-singlet ($\tau = S$) excited states.

We denote by $|c,c;+1\rangle$, $|c,c;0\rangle$, $|c,c;-1\rangle$, and $|c,c,\eta 1;0\rangle$ (and $|s1,s1;+1\rangle$, $|s1,s1;0\rangle$, $|s1,s1;-1\rangle$, and $|s1,s1,s2;0\rangle$), where the index with values $0,\pm 1$ refers to $S_\eta^{x_3}$ (and $S_s^{x_3}$), the four excited energy eigenstates associated with the four one-pseudofermion scattering states needed to express the four holon-holon states $|+1/2,+1/2;\eta\rangle$, $|+1/2,-1/2;\eta\rangle$, $|-1/2,+1/2;\eta\rangle$, and $|-1/2,-1/2;\eta\rangle$ of the scattering theory of Ref. [42] (and with the four spinon-spinon states $|+1/2,+1/2;s\rangle$, $|+1/2,-1/2;s\rangle$, $|-1/2,+1/2;s\rangle$, and $|-1/2,-1/2;s\rangle$ of both the scattering theories of Refs. [40] and [41, 42].) It is confirmed below that concerning the four holon-holon states of Ref. [41, 42] and the four spinon-spinon states of both Refs. [40] and [41, 42] there is no one-to-one correspondence to the corresponding four states of the pseudofermion scattering theory.

At fixed values of the two excited-state emerging momenta q_j and $q_{j'}$, the three states denoted by $|c,c;S_\eta^{x_3}\rangle$ (and $|s1,s1;S_s^{x_3}\rangle$) corresponding to $S_\eta^{x_3} = 0,\pm 1$ (and $S_s^{x_3} = 0,\pm 1$) are the three η-spin-triplet excited states (and three spin-triplet excited states) considered in Appendix C. (See the corresponding number and number deviation values in the Tables II-IV of that appendix.) Our phase shifts refer to all the N_c excited-state c pseudofermion scatterers, N_c^h c pseudofermion-hole scatterers, N_{s1} $s1$ pseudofermion scatterers, and N_{s1}^h $s1$ pseudofermion-hole scatterers. On the other hand, the phase shifts of the scatterers of both Refs. [40] and [41, 42] refer only to one of the present c (and $s1$) pseudofermion-hole scatterers. Indeed, the two branch indices c,c (and $s1,s1$) of the above states refer to the specific c (and $s1$) pseudofermion-hole scatterer and c (and $s1$) pseudofermion-hole scattering center, respectively, created under the transition from the ground state to the excited state. Furthermore, $|c, c, \eta 1; 0\rangle$ (and $|s1, s1, s2; 0\rangle$) denotes the η-spin-singlet excited state (and spin-singlet excited state) whose three branch indices $c, c, \eta 1$ (and $s1, s1, s2$) refer to the specific c pseudofermion-hole scatterer, c pseudofermion-hole scattering center, and $\eta 1$ pseudofermion scattering center (and $s1$ pseudofermion-hole scatterer, $s1$ pseudofermion-hole scattering center, and $s2$ pseudofermion scattering center) created under such a transition.

We note that the eight one-particle scattering states of both Refs. [40] and [42] associated with ground-state - excited-state transitions where one traditional holon and one traditional spinon are created correspond to only the four doublet excited energy eigenstates whose number and number deviation values are provided in Tables II-IV of Appendix C. In that case the four many-particle states of the three representations associated with these eight one-particle scattering states are the same states.

For the pseudofermion scattering theory we find for the reduced subspace at fixed values of the two c and/or $s1$ band hole momenta q_j and $q_{j'}$ the following fully diagonal 16×16 dressed S matrix,

$$\bar{\mathbf{S}} = \mathbf{U}^{\dagger}\mathbf{S}\mathbf{U} = \begin{bmatrix} \bar{\mathbf{S}}_{SS} & 0 & 0 & 0 \\ 0 & \bar{\mathbf{S}}_{S\eta} & 0 & 0 \\ 0 & 0 & \bar{\mathbf{S}}_{\eta S} & 0 \\ 0 & 0 & 0 & \bar{\mathbf{S}}_{CC} \end{bmatrix} \, ; \quad \mathbf{U} = \begin{bmatrix} \mathbf{J} & 0 & 0 & 0 \\ 0 & \mathbf{I} & 0 & 0 \\ 0 & 0 & \mathbf{I} & 0 \\ 0 & 0 & 0 & \mathbf{J} \end{bmatrix} \, , \tag{92}$$

where $\mathbf{U}$ is a unitary matrix and the 4×4 matrix $\mathbf{J}$ is given in Eq. (91).

The three first diagonal entries (and the fourth diagonal entry) of the two 4×4 diagonal dressed S matrix blocks $\bar{\mathbf{S}}_{\beta\beta}$ such that $\beta = C, S$ of the $\bar{\mathbf{S}}$ expression given in Eq. (92) equal the phase factor $S_{\beta T}$ (and equals the phase factor $S_{\beta S}$) provided in Eq. (89). The four diagonal entries of the other two 4×4 diagonal dressed S matrix blocks $\bar{\mathbf{S}}_{\beta'\beta''}$ such that $\beta'\beta'' = S\eta, \eta S$ of the same expression are equal and given in Eq. (90).

For the general pseudofermion scattering theory the diagonal entries of the 16×16 diagonal dressed matrix $\bar{\mathbf{S}}$ provided in Eq. (92) are sixteen dressed S matrices $S_{\beta}(q_j)$ of dimension one whose general form is given in Eq. (35). They correspond to the c and $s1$ pseudofermion-hole scatterers of the reduced-subspace excited states considered here. We recall that for each excited state the phase shifts in the entries of the dressed S matrices of the scattering theories of Refs. [40, 42] given in Eqs. (86) and (91) and those in the corresponding diagonal dressed S matrix provided in Eq. (92) refer to the specific scatterers created under the transition to the excited state.

A major difference between the pseudofermion scattering theory and the traditional holon and spinon scattering theories of both Ref. [40] and Refs. [41, 42] is though that the scatterers of the present formulation include the macroscopic number of those that pre-exist in the ground state. Specifically, the pseudofermion scattering theory general S matrices $S_{\beta}(q_j)$ of Eq. (35) apply to all the N_{a_β} scatterers of that theory. Most of those re-exist in the ground state. As further discussed below in Section VI C, that the pseudofermion scattering theory accounts for the phase shifts of scatterers other than only those created under the transitions to the excited states is a necessary condition for it accounting as well for the microscopic processes that control the one- and two-electron dynamical and spectral properties of the 1D Hubbard model.

Use of the unitary matrices defined in Eqs. (91) and (92) reveals that the above four holon-holon ($\alpha = \eta$) states of Ref. [42] and four spinon-spinon ($\alpha = s$) states of both the descriptions of Refs. [42] and [40] can be expressed in terms of the excited energy eigenstates that contain the one-pseudofermion scattering states of the present formulation as follows,

$$\begin{aligned} |+1/2, +1/2; \alpha\rangle &= |\beta, \beta; -1\rangle \, ; \quad |+1/2, -1/2; \alpha\rangle = \frac{1}{\sqrt{2}}\Big[|\beta, \beta; 0\rangle - |\beta, \beta, \alpha\nu'; 0\rangle\Big] \, , \\ |-1/2, +1/2; \alpha\rangle &= \frac{1}{\sqrt{2}}\Big[|\beta, \beta; 0\rangle + |\beta, \beta, \alpha\nu'; 0\rangle\Big] \, ; \quad |-1/2, -1/2; \alpha\rangle = |\beta, \beta; +1\rangle \, . \end{aligned} \tag{93}$$

Here $\alpha = \eta$, $\beta = c$, and $\alpha\nu' = \eta 1$ for the holon-holon states of Refs. [41, 42] and $\alpha = s$, $\beta = s1$, and $\alpha\nu' = s2$ for the spinon-spinon states of both Refs. [40] and [41, 42]. Such relations confirm that the states $|+1/2, -1/2; \alpha\rangle$ and $|-1/2, +1/2; \alpha\rangle$ associated with the holon-holon and/or spinon-spinon one-particle scattering states of the traditional holon and spinon descriptions are not eigenstates of the η-spin ($\alpha = \eta$) or spin ($\alpha = s$). Importantly, however, they are energy eigenstates, since they are a superposition of two energy eigenstates with the same energy eigenvalue.

The validity of the scattering theories requires that the excited states associated with the asymptotic one-particle scattering states have a well-defined energy. For the present quantum problem, this requirement is fulfilled provided that the excited states associated with the one-particle scattering states are energy eigenstates of the model. Such a requirement is obeyed by all the excited states associated with the one-particle scattering states of the pseudofermion scattering theory, which are indeed energy eigenstates of the 1D Hubbard model. This is true for all PS excited states of initial ground states with arbitrary values of the electronic density n and spin density m.

For the traditional holon and spinon scattering theory of Ref. [40] that requirement is obeyed for the PS excited states of $S_s = 0$ ground states of arbitrary electronic density n. That such theory is valid only for such excited states follows from only as $m \rightarrow 0$ the energy of the set of excited states with the same set of $s1$ band hole momentum values $\{q_h\}$ where $h = 1, ..., N^h_{s1}$ and $N^h_{s1} = N_s$ but different values for the unbound-spinon numbers M^{un}_s and set of $s\nu$ pseudofermion numbers $\{N_{s\nu}\}$ of the present formulation being the same. This allows each of the scattering states of theory of Ref. [40] being expressed as a superposition of energy eingenstates with the same energy eigenvalue. Hence it is the degeneracy of the above types of excited energy eigenstates that determines the subspace where the holons and spinons as defined in Ref. [40] refer to a valid choice of scattering states.

Finally, for the traditional spinons and holons as defined in Refs. [41, 42] such a property holds only for the PS excited states of the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state. This is is due to at fixed values of the set of c (and $s1$) band hole momentum values $\{q_h\}$ where $h = 1, ..., N^h_c$ and $N^h_c = L_c$ (and $h = 1, ..., N^h_{s1}$ and $N^h_{s1} = L_s$) the energy of that ground-state excited states with the same number L_h of traditional holons (and L_s of traditional spinons) but different values for the unbound-η-spinon numbers M^{un}_η and set of $\eta\nu$ pseudofermion numbers $\{N_{\eta\nu}\}$ (and for the unbound-spinon numbers M^{un}_s and set of $s\nu$ pseudofermion numbers $\{N_{s\nu}\}$) of the present formulation being the same. This holds provided that the PS limiting conditions $L_h/N_a \rightarrow 0$ and $L_s/N_a \rightarrow 0$ as $N_a \rightarrow \infty$ are fulfilled. This property again assures that the scattering states of that traditional holon and spinon description can be expressed as a superposition of energy eingenstates with the same energy eigenvalue. Thus it is again the degeneracy of the above types of excited energy eingenstates that determines the subspace where the holons and spinons as defined in Refs. [41, 42] correspond to a acceptable choice of scattering states.

Consistent, the traditional holon and spinon scattering theory of Ref. [40] (and Refs. [41, 42]) refers only to the 1D Hubbard model at zero magnetization (and at half filling and zero magnetization). More specifically, its applicability refers in the thermodynamic limit to PSs spanned by the excited states of any $S_s = 0$ ground state (the PS spanned by the excited states of the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state.)

In Appendix D the role of the degeneracy of the excited energy eigenstates in the definition of the subspaces where the traditional spinon and holon descriptions of Ref. [40] and Refs. [41, 42] are defined is further discussed. The situation is found to be qualitatively different for finite u values and in the $u \rightarrow \infty$ limit, respectively.

C. Relation to the finite-energy spectral properties

Both the traditional holon and spinon scattering theories of Ref. [40] and Refs. [41, 42], respectively, consider explicitly only the scatterers created under the transitions from the ground state to the excited states. It is confirmed below that within the PDT [7–9] the phase shifts of the macroscopic number of $\beta = c, s1$ pseudofermion scatterers that pre-exist in the initial ground state play a major role in the 1D Hubbard model unusual finite-energy spectral properties.

A one- or two-electron excitation $\mathcal{O}^\dagger|GS\rangle$ is contained in a well-defined direct sum, Eq. (7), of CPHS ensemble subspaces. It is a superposition of excited energy eigenstates. Each of these states is described by a set of deviations $\{\delta N_\beta(q_j)\}$, Eq. (A18) of Appendix A, involving a finite number of scattering centers. One can consider that the order of the direct sum of Eq. (7) of CPHS ensemble subspaces associated with a given one- or two-electron excitation is that of the increasing values of the minimum energy $\omega_0(D_r, S_r)$, Eq. (6), of each CPHS ensemble subspace, relative to that of the initial ground state. For fixed value of the numbers S_r and D_r defined in Eq. (A19) of Appendix A, these energy values are such that,

$$\omega_0(0, S_r) < \omega_0(1, S_r) < \omega_0(2, S_r) < ... \, , \tag{94}$$

and

$$\omega_0(D_r, 0) < \omega_0(D_r, 1) < \omega_0(D_r, 2) < ... \, , \tag{95}$$

respectively. Note that for fixed values of electronic density $n \in [0, 1]$ and spin density $m \in]0, n]$, the ordering of the energies $\omega_0(D_r, S_r)$ is well defined and such that,

$$\omega_0(0, 0) < \min\{\omega_0(0, 1), \omega_0(1, 0)\} < ... \, . \tag{96}$$

An important property is that for a given energy range $\delta E_{l_o, l_\triangle} \in [0, \omega_0]$ the number of CPHS ensemble subspaces of the direct sum of Eq. (7) is finite. Moreover, as each energy $\omega_0(D_r, S_r)$ is reached, an increasing number of channels opens up, which correspond to CPHS ensemble subspaces of increasing energy. By new channels we mean here $\beta = \alpha\nu$ pseudofermion branches of increasing ν value. Thus, for a given energy range $\delta E_{l_o, l_\triangle} \in [0, \omega_0]$ there is a well defined set of β pseudofermion dressed S matrices, whose "in" and "out" asymptote one-pseudofermion scattering states correspond to the set of excited energy eigenstates spanning the direct sum given in Eq. (7) of CPHS ensemble subspaces of energy smaller than or equal to ω_0. The β pseudofermion dressed S matrices belonging to that finite set control the spectral properties for energies in the range under consideration, $\delta E_{l_o,l_\triangle} \in [0, \omega_0]$.

The changes in the pseudofermion occupancy configurations under the ground-state - excited-state transitions inside the scattering-theory PS play a major role in the PDT [7–9]. Within the $N_a \gg 1$ limit of the present formulation the corresponding pseudofermion elementary processes that generate the PS from the ground state can be classified into three types:

(A) Finite-energy and finite-momentum elementary $\beta = c, s1$ (and $\beta = \alpha\nu \neq s1$) pseudofermion processes involving creation or annihilation (and creation) of one or a finite number of pseudofermions at momentum values away and different from the $\beta = c, s1$ Fermi momenta, $\bar{q}^{\iota}_{F\beta}$, where $\iota = \pm 1$ (and at $\beta = \alpha\nu \neq s1$ band momentum values);

(B) Zero-energy and finite-momentum processes that change the number of $\beta = c, s1$ pseudofermions at the $\iota = +1$ right and $\iota = -1$ left $\beta = c, s1$ Fermi points.

(C) Low-energy and small-momentum elementary $\beta = c, s1$ pseudofermion particle-hole processes in the vicinity of the $\iota = +1$ right and $\iota = -1$ left $\beta = c, s1$ Fermi points, relative to the excited-state $\alpha = c, s$ pseudofermion momentum occupancy configurations generated by the above elementary processes (A) and (B).

In the following we provide some useful information that confirms the major role plaid by the phase shifts and dressed S matrix of the pseudofermion scattering theory introduced in this paper in the one- and two-electron spectral function expressions of the PDT near particular (k, ω)-plane points and spectral branch lines.

The energy and momentum spectrum of the excited energy eigenstates generated by the processes (A) and (B),

$$\delta E^{\odot}_{l_o} = E^{\odot}_{l_o} - E_{GS}\, ; \quad \delta P^{\odot}_{l_o} = P^{\odot}_{l_o} - P_{GS}\, , \tag{97}$$

are important pieces of the PDT spectral-function expressions. For simplicity, here we have omitted the states number $l_\triangle$ that within our notation stands for the set of numbers $[2S_c, S_\eta, S_s, M^{un}_{\eta,-1/2}, M^{un}_{s,-1/2}]$. Indeed, the sum over excited states considered in the following is over states with the same $l_\triangle$ values.

For instance, the processes (A) and (B) that generate a $\beta = c, s1$ branch line involve creation of one $\beta = c, s1$ pseudofermion ($c_0 = +1$) or $\beta = c, s1$ pseudofermion hole ($c_0 = -1$) of bare momentum q away from the $\beta = c, s1$ Fermi points. The remaining $\beta' = c, \alpha\nu$ pseudofermions or $\beta' = c, s1$ pseudofermion holes are created by such processes at the corresponding Fermi points ($\beta' = c, s1$) or band limiting momentum values $\pm q_{\alpha\nu}$ ($\beta' = \alpha\nu \neq s1$). The spectra, Eq. (97), of the excited energy eigenstates generated from the ground state by such processes (A) and (B) have the general form,

$$\delta E^{\odot}_{l_o} = \omega_0 + |\varepsilon_\beta(q)|\, ; \quad \delta P^{\odot}_{l_o} = P_0 + c_0\, q\, , \quad \beta = c, s1\, , \quad c_0 = \pm 1\, . \tag{98}$$

Here $\omega_0 = \omega_0(D_r, S_r)$ is the energy given in Eq. (6) and the constant momentum P_0 results from the possible creation of unbound $-1/2$ η-spinons and $\beta' = c, \alpha\nu$ pseudofermions and $\beta' = c, s1$ pseudofermion holes other than the $\beta = c, s1$ pseudofermion or $\beta = c, s1$ pseudofermion hole of bare momentum q.

Another type of spectral feature refers to a point-like singularity. It is generated by processes such that all $\beta = c, \alpha\nu$ pseudofermions and $\beta = c, s1$ pseudofermion holes are created under the transition to the excited state at the corresponding Fermi points $\pm q_{F\beta'}$ ($\beta' = c, s1$) or β' band limiting momentum values $\pm q_{\beta'}$ ($\beta' = \alpha\nu \neq s1$). In that case the spectra, Eq. (97), of the corresponding excited energy eigenstate generated by the processes (A) and (B) have the form $\delta E^{\odot}_{l_o} = \omega_0$ and $\delta P^{\odot}_{l_o} = P_0$.

Within the PDT, a one- or two-electron spectral function $B(k,\, \omega)$ can near the above spectral features be written as a sum of terms. Each of those refers to a small subspace spanned by states with the same elementary-objects numbers. Near the (k, ω)-plane points and spectral branch lines mentioned above, the spectral function $B(k,\, \omega)$ may be replaced by the dominant term of such a set of terms, which here we denote by $B^{\odot}(k,\, \omega)$ [7–9],

$$B(k,\, \omega) \approx B^{\odot}(k,\, \omega) = \sum_{l^{\odot}_o} \Theta\Big(\Omega - \delta\omega_{l_o}\Big)\, \Theta\Big(\delta\omega_{l_o}\Big)\, \Theta\left(|v_{l_o}| - v_{\bar{\beta}}\right) \breve{B}^{\odot}_{l_o}(\delta\omega_{l_o}, v_{l_o})\, . \tag{99}$$

The $\sum_{l^{\odot}_o}$ sum appearing here is over excited energy eigenstates generated by processes (A) and (B) with excitation energy and momentum, Eq. (97), in the ranges $\delta E^{\odot}_{l_o} \in [\omega - \Omega, \omega]$ and $\delta P^{\odot}_{l_o} \in [k - \Omega/v_{l_o}, k]$,

$$\begin{aligned} \delta\omega_{l_o} &= [\omega - \delta E^{\odot}_{l_o}]\, ; \quad \delta k_{l_o} = k - \delta P^{\odot}_{l_o}\, , \\ \delta E_{l_o} &= \delta E^{\odot}_{l_o} + \delta\omega_{l_o} = \omega\, ; \quad P_{l_o} = \delta P^{\odot}_{l_o} + \delta k_{l_o} = k\, , \\ v_{l_o} &= \frac{\delta\omega_{l_o}}{\delta k_{l_o}}\, ; \quad v_{\bar{\beta}} = \min\{v_c, v_{s1}\}\, ; \quad v_\beta = \max\{v_c, v_{s1}\}\, . \end{aligned} \tag{100}$$

Here $\delta E_{l_o} = \omega$ and $\delta P_{l_o} = k$ denote the excitation energy and momentum, respectively, of final energy eigenstates including contributions from the processes (C) and Ω denotes the energy cut-off of such processes. The value of that cut-off defines a small energy range within which the velocity v_{l_o} given in Eq. (100) remains nearly unchanged. Fortunately, Ω does not appear in the k and ω dependences found for the spectral functions by the PDT [7–9].

The general excitation energy spectrum expressed in Eq. (21) in terms of $\beta = c, s1$ pseudofermion occupancy configurations lacks second-order residual interaction terms, in contrast to exactly the same excitation energy spectrum when expressed in terms of $\beta = c, s1$ fermions, as given in Eqs. (18)-(19). Consistent, within the present $N_a \gg 1$ limit the $\beta = c, s1$ pseudofermions are not energy entangled, so that the function $\breve{B}^{\odot}_{l_o}(\delta\omega_{l_o}, v_{l_o})$ appearing in Eq. (99) can be expressed as a convolution of c and $s1$ pseudofermion spectral functions as follows,

$$\begin{aligned}\breve{B}^{\odot}_{l_o}(\delta\omega_{l_o}, v_{l_o}) &= \frac{\mathrm{sgn}(v_{l_o})}{2\pi}\int_0^{\delta\omega_{l_o}} d\omega' \int_{-\mathrm{sgn}(v_{l_o})\delta\omega_{l_o}/v_\beta}^{+\mathrm{sgn}(v_{l_o})\delta\omega_{l_o}/v_\beta} dk'\, B_{Q_{\bar{\beta}}}(\delta\omega_{l_o}/v_{l_o} - k', \delta\omega_{l_o} - \omega') \\ &\times\; B_{Q_\beta}(k', \omega')\,. \end{aligned} \tag{101}$$

Here $\bar{\beta} = c, s1$ and $\beta = s1, c$, respectively, are chosen according to the criterion of Eq. (100) concerning the relative magnitudes of the two $\beta = c, s1$ Fermi-points velocities v_β defined in Eq. (A23) of Appendix A.

Due to the orthogonal catastrophe associated with the shake up effects resulting from the $\beta = c, s1$ band discrete canonical-momentum value shift under the transition to the excited states, a large number of particle-hole processes (C) near the $\beta = c, s1$ pseudofermion Fermi points lead to finite-spectral weight contributions. The processes (C) refer to the linear part of the $\beta = c, s1$ energy dispersions, in the vicinity of the $\beta = c, s1$ Fermi points. Thus they involve the $\beta = c, s1$ Fermi-points velocities, v_β, given in Eq. (A23) of Appendix A. Within the PDT, the number $m_{\beta,\,\iota} = 1, 2, 3, ...$ labels the different types of processes (C) near the α, ι Fermi point, where $\beta = c, s1$ and $\iota = \pm 1$. Its value, $m_{\beta,\,\iota} = 1, 2, 3, ...$, gives the number of elementary processes of momentum $\pm 2\pi/N_a$ involved in the processes (C) under consideration.

The PDT expressions of the c and $s1$ pseudofermion spectral functions in Eq. (101) involve sums that run over the numbers $m_{\beta,\,\iota} = 1, 2, 3, ...$ associated with the processes (C) [7–9],

$$\begin{aligned}B_{Q_\beta}(k', \omega') &= \frac{N_a}{2\pi}\sum_{m_{\beta,\,+1}}\sum_{m_{\beta,\,-1}} A_\beta(m_{\beta,\,+1},\, m_{\beta,\,-1}) \\ &\times\; \delta\Big(\omega' - \frac{2\pi}{N_a}v_\beta\sum_{\iota=\pm 1} m_{\beta,\,\iota}\Big)\,\delta\Big(k' - \frac{2\pi}{N_a}\sum_{\iota=\pm 1}\iota\, m_{\beta,\,\iota}\Big)\,, \quad \beta = c, s1\,. \end{aligned} \tag{102}$$

Here,

$$A_\beta = A^{(0,0)}_\beta\, a_\beta(m_{\beta,\,+1},\, m_{\beta,\,-1})\,, \quad \beta = c, s1\,, \tag{103}$$

where the weight $A^{(0,0)}_\beta$ is associated with the transition from the ground state to an excited state generated by the processes (A) and (B) and the relative weights $a_\beta(m_{\beta,\,+1},\, m_{\beta,\,-1})$ are generated by the additional processes (C).

Within the algebra used by the PDT to derive the spectral weights given in Eq. (103), the effective pseudofermion anticommutators provided in Eq. (24) play a major role. Such anticommutators can be expressed solely in terms of the difference $\bar{q} - \bar{q}'$ of the corresponding β and β' pseudofermions canonical momentum values and the dressed S matrix $S_\beta(q)$ given in Eq. (35) for the β pseudofermion. The latter object plays the role of scatterer in the excited energy eigenstate. The anticommutators under consideration can be written as,

$$\{f^\dagger_{\bar{q},\beta}, f_{\bar{q}',\beta'}\} = \delta_{\beta,\beta'}\frac{1}{N_{a_\beta}}\Big[S_\beta(q)\Big]^{1/2} e^{-i(\bar{q}-\bar{q}')/2}\,\frac{\mathrm{Im}\Big[S_\beta(q)\Big]^{1/2}}{\sin([\bar{q}-\bar{q}']/2)}\,. \tag{104}$$

The related anticommutators between two pseudofermion creation or annihilation operators vanish. The expression given in Eq. (104) reveals that the dressed S matrix, $S_\beta(q_j)$, controls the pseudofermion effective anticommutators. After a suitable algebra that involves such anticommutators, one finds that the weight $A^{(0,0)}_\beta$ on the right-hand side of Eq. (103) reads [7, 8],

$$\begin{aligned} A_{\beta}^{(0,0)} &= \Big(\frac{1}{N_{a_\beta}}\Big)^{2N_\beta^{\odot}} \prod_{j=1}^{N_{a_\beta}} \sin^2\Big(\frac{N_\beta^{\odot}(q_j)[Q_\beta(q_j)-\pi]+\pi}{2}\Big) \prod_{j=1}^{N_{a_\beta}-1} \Big[\sin\Big(\frac{\pi j}{N_{a_\beta}}\Big)\Big]^{2[N_{a_\beta}-j]} \\ &\times \prod_{i=1}^{N_{a_\beta}} \prod_{j=1}^{N_{a_\beta}} \theta(j-i)\, \sin^2\Big(\frac{N_\beta^{\odot}(q_j)N_\beta^{\odot}(q_i)[Q_\beta(q_j)-Q_\beta(q_i)+2\pi(j-i)-\pi N_{a_\beta}]+\pi N_{a_\beta}}{2N_{a_\beta}}\Big) \\ &\times \prod_{i=1}^{N_{a_\beta}} \prod_{j=1}^{N_{a_\beta}} \frac{1}{\sin^2\Big(N_\beta^{\odot}(q_i)N_\beta^{\odot}(q_j)[\frac{\pi(j-i)}{N_{a_\beta}}+\frac{Q_\beta(q_j)}{2N_{a_\beta}}-\frac{\pi}{2}]+\frac{\pi}{2}\Big)}\,, \quad \beta = c, s1\,. \end{aligned} \tag{105}$$

The number of β band discrete momentum values, N_{a_β}, and the β band momentum distribution function, $N_\beta^{\odot}(q_j)$, in this expression are those of the excited state generated by the processes (A) and (B) and $Q_\beta(q_j)$ is the phase-shift functional defined by Eqs. (11), (15), and (16). The deviations in that functional expression are those generated by the corresponding ground-state - excited-state transition.

The form of the weight given in Eq. (105) confirms that within the PDT of Refs. [7, 8], the overall β pseudofermion and hole phase-shift functional $Q_\beta(q_j)/2$ in Eq. (15), which is associated with the dressed S matrix $S_\beta(q_j) = e^{i\,Q_\beta(q_j)}$, Eq. (35), controls the one- and two-electron spectral properties. Within the algebra used to derive such a weight expression, that phase-shift functional arises from of the pseudofermion effective anticommutators, Eq. (104). For the PDT, the general expression of the relative weights $a_\beta(m_{\beta,+1}, m_{\beta,-1})$ appearing in Eq. (103) in the tower of excited energy eigenstates generated by the processes (C) reads [7],

$$a_\beta(m_{\beta,+1}, m_{\beta,-1}) = \Big[\prod_{\iota=\pm1} a_{\beta,\iota}(m_{\beta,\iota})\Big]\Big[1+\mathcal{O}\Big(\frac{\ln N_a}{N_a}\Big)\Big]\,, \quad \beta = c, s1\,. \tag{106}$$

Here the relative weight $a_{\beta,\iota}(m_{\beta,\iota})$ is given by,

$$a_{\beta,\iota}(m_{\beta,\iota}) = \prod_{j=1}^{m_{\beta,\iota}} \frac{(2\Delta_\beta^\iota + j - 1)}{j} = \frac{\Gamma(m_{\beta,\iota}+2\Delta_\beta^\iota)}{\Gamma(m_{\beta,\iota}+1)\,\Gamma(2\Delta_\beta^\iota)}\,, \quad \beta = c, s1\,, \quad \iota = \pm 1\,, \tag{107}$$

where $\Gamma(x)$ is the usual gamma function. It follows from Eq. (107) that,

$$a_{\beta,\iota}(1) = 2\Delta_\beta^\iota\,, \quad \beta = c, s1\,, \quad \iota = \pm 1\,. \tag{108}$$

The four functionals $2\Delta_\beta^\iota$ appearing here play a major role in the PDT. They are directly related to the right ($\iota = +1$) and left ($\iota = -1$) $\beta = c, s1$ canonical-momentum Fermi-point deviations $\delta\bar{q}_{F\beta}^\iota \approx \iota\,\delta\bar{q}_{F\beta}$. Specifically, they are given by $2\Delta_\beta^\iota = [\delta\bar{q}_{F\beta}^\iota/(2\pi/N_a)]^2$. It follows from the expressions provided in Eq. (22) that in units of $2\pi/N_a$ the corresponding four $\beta = c, s1$ and $\iota = \pm1$ canonical-momentum Fermi-point deviations read $\delta\bar{q}_{F\beta}^\iota/(2\pi/N_a) = \iota\,\delta N_{\beta,\iota}^F + Q_\beta^\Phi(\iota q_{F\beta})/2\pi$. They can alternatively be written as $\delta\bar{q}_{F\beta}^\iota/(2\pi/N_a) = \iota\,\delta N_{\beta,\iota}^{0,F} + Q_\beta(\iota q_{F\beta})/2\pi$. Here $\delta N_{\beta,\iota}^{0,F}$ denotes the deviation in the number of $\beta = c, s1$ pseudofermions at the right ($\iota = +1$) and left ($\iota = +1$) $\beta = c, s1$ Fermi points, which results from the ground-state - excited-state transition, without accounting for the possible shift of the BA quantum numbers I_j^β considered in the first paper. The latter can be integers or half-odd integers and refer to the β band bare discrete momentum values, $q_j = [2\pi/N_a]\,I_j^\beta$, in units of $2\pi/N_a$.

The four functionals $2\Delta_\beta^\iota$ can be expressed in terms of the β pseudofermion dressed S matrix $S_\beta(q) = e^{i\,Q_\beta(q)}$ and corresponding β phase shift $Q_\beta(q)/2$ introduced in this paper of the scatterers at the four $\beta = c, s1$ and $\iota = \pm1$ Fermi points, $q = \iota q_{F\beta}$, as follows,

$$\begin{aligned} 2\Delta_\beta^\iota &= \left(\iota\delta N_{\beta,\iota}^{0,F} - i\frac{\log S_\beta(\iota q_{F\beta})}{2\pi}\right)^2\,, \\ &= \left(\iota\delta N_{\beta,\iota}^{0,F} + \frac{Q_\beta(\iota q_{F\beta})}{2\pi}\right)^2\,, \quad \beta = c, s1\,, \quad \iota = \pm 1\,. \end{aligned} \tag{109}$$

Consistent with Eq. (35), here the log branch is $\log S_\beta(\iota q_{F\beta}) = i\,Q_\beta(\iota q_{F\beta})$.

The expression provided in Eq. (108) reveals that the functional given in Eq. (109) is the relative weight associated with the $m_{\beta,\,\iota} = 1$ peak of the α, ι pseudofermion spectral function. Moreover,

$$a_{\beta}(1,\,0) = 2\Delta_{\beta}^{+1}\,;\quad a_{\beta}(0,\,1) = 2\Delta_{\beta}^{-1}\,,\quad \beta = c, s1\,. \tag{110}$$

The function of Eq. (107) has the following asymptotic behavior,

$$a_{\beta,\,\iota}(m_{\beta,\,\iota}) \approx \frac{1}{\Gamma(2\Delta_{\beta}^{\iota})}\Big(m_{\beta,\,\iota}\Big)^{2\Delta_{\beta}^{\iota}-1}\,;\quad 2\Delta_{\beta}^{\iota} \neq 0\,;\quad \beta = c, s1\,;\quad \iota = \pm 1\,. \tag{111}$$

The $\beta = c, s1$ spectral function weight $A_{\beta}^{(0,0)}$, Eq. (105), determines mostly the absolute weight of the spectral feature, as k and ω tend to the corresponding (k,ω)-plane singular branch line or point. That feature k and ω dependence in the vicinity of such a line or point is rather controlled by the relative weights $a_{\beta}(m_{\beta,\,+1},\, m_{\beta,\,-1})$ in the general weight expression $A_{\beta} = A_{\beta}^{(0,0)}\, a_{\beta}(m_{\beta,\,+1},\, m_{\beta,\,-1})$, Eq. (103), which are given in Eqs. (106), (107), (108), (110), and (111). Within the PDT and present formulation $N_a \gg 1$ limit, the dominant processes correspond to very large values $m_{\beta,\,\iota} \gg 1$ in the product $\prod_{j=1}^{m_{\beta,\,\iota}}$ of the relative weight $a_{\beta,\,\iota}(m_{\beta,\,\iota})$ expression provided in Eq. (107). Such large $m_{\beta,\,\iota}$ values imply a large range for the product index value $j = 1, 2, ..., m_{\beta,\,\iota}$. This means that such a relative weight has contributions from processes (C) involving a large number of elementary $\beta = c, s1$ band particle-hole processes of increasing momentum $\iota\,[2\pi/N_a]\,j$ absolute value, $[2\pi/N_a]\,j = 2\pi/N_a, 4\pi/N_a, ..., 2m_{\beta,\,\iota}\pi/N_a$, where $\iota = \pm 1$. A large range for the product $\prod_{j=1}^{m_{\beta,\,\iota}}$ implies as well that the relative weight $a_{\beta,\,\iota}(m_{\beta,\,\iota})$ behaves as $a_{\beta,\,\iota}(m_{\beta,\,\iota}) \approx [1/\Gamma(2\Delta_{\beta}^{\iota})]\,(m_{\beta,\,\iota})^{2\Delta_{\beta}^{\iota}-1}$ for $N_a \gg 1$, as given in Eq. (111).

Importantly, the contributions from the creation by the processes (A) and (B) of elementary objects with no occupancy in the initial ground state, specifically $\beta = \alpha\nu \neq s1$ pseudofermions, unbound $-1/2$ η-spinons, and unbound $-1/2$ spinons, is within the function $\breve{B}_{l_o}^{\odot}(\delta\omega_{l_o}, v_{l_o})$ given in Eq. (101) merely accounted for through their contribution to the excited-state energy and momentum spectra, Eq. (97). Their absence in the initial ground state renders the corresponding spectral-function matrix-element quantum overlap orthogonal-catastrophe free [7, 8]. The contribution to the spectral weights from the tower-states associated with the above $j = 1, 2, ..., m_{\beta,\,\iota}$ processes (C) refers only to the $\beta = c, s1$ branches with finite occupancy in the initial ground state. That so many tower states contribute to the spectral-function matrix elements results from the shake up effects and corresponding $\beta = c, s1$ band discrete canonical-momentum value shifts under the transition to the excited states. Those stem from the initial ground state and the excited state having different values for the set of $\beta = c, s1$ band discrete canonical-momentum values. (Such effects do not occur for the $\beta = \alpha\nu \neq s1$ momentum bands, since they are empty in the initial ground state.)

The pseudofermion description considerably simplifies the study of the exotic quantum-liquid finite-energy spectral properties [7, 8, 22, 23]. Note, however, that within the PDT Lehmann representations, technical limitations exist. Those arise from the difficulty in summing up all contributions to the spectral weights from the overlap of the initial ground state with the whole set of suitable excited energy eigenstates. In general such sums are difficult to perform, since the overall β pseudofermion and hole phase-shift functional provided in Eq. (15) has a different value for each excited state. Fortunately, such sums can be performed for some line shapes corresponding to the vicinity of important spectral-weight features. That includes the $\beta = c, s1$ branch lines and singularities near important (k,ω)-plane points [7, 8].

Indeed, within the present $N_a \gg 1$ limit, following the use of the relative weight $a_{\beta,\,\iota}(m_{\beta,\,\iota})$ expression provided in Eq. (111) in Eqs. (99), (101), (102), and (103), the sum $\sum_{l_o^{\odot}}$ in the spectral-function expression given in Eq. (99) can be performed. This leads to the following spectral-function behavior near a $\beta = c, s1$ branch line whose general shape is defined by the energy and momentum spectra of Eq. (98) [7, 8],

$$\begin{aligned} B(k,\,\omega) &\propto (\omega - \omega_{\beta}(k))^{\zeta(k)}\,;\quad (\omega - \omega_{\beta}(k)) \geq 0\,, \\ \omega_{\beta}(k) &= |\varepsilon_{\beta}(q)|_{q=c_0\,[k-P_0]}\,;\quad \zeta(k) = -1 + \sum_{\beta'=c,s1}\sum_{\iota=\pm 1} 2\Delta_{\beta'}^{\iota}(q)|_{q=c_0\,[k-P_0]}\,. \end{aligned} \tag{112}$$

Within the PDT, this expression is exact for $\beta = c, s1$ branch lines that limit the bottom of the (k,ω)-plane finite spectral-weight region. Hence there is no weight beneath such lines. For the particular case of the one-electron spectral function, this expression is a good approximation for the $\beta = c, s1$ branch lines that have a small amount of spectral weight beneath.

In the expressions provided in Eq. (112) $\varepsilon_{\beta}(q)$ is the $\beta = c, s1$ energy dispersion given in Eq. (A21) of Appendix (A) for the $\beta = c, s1$ branch-line excited states generated by processes (A) and (B) whose spectra is provided in Eq. (98) and the four functionals $2\Delta_{\beta'}^{\iota} = 2\Delta_{\beta'}^{\iota}(q)$ of Eq. (109) depend on the momentum $k = P_0 + c_0\,q$ through their dependence on the bare-momentum q of the $\beta = c, s1$ scattering center created under the transition to such states [7–9]. Indeed, we recall that each ground-state - excited-state transition corresponds to uniquely defined values of the set of deviations $\delta N_{\beta'}(q_{j'})$ in the expression of the β phase shift $Q_\beta(q_j)/2$ defined by Eqs. (11), (15), and (16) and corresponding β dressed S matrix $S_\beta(q_j) = e^{iQ_\beta(q_j)}$ given in Eq. (35). The latter depends on such deviations through each of the elementary S matrices, $S_{\beta,\beta'}$, in the expression provied in Eq. (36). Therefore, in the expression of the above functionals $2\Delta^\iota_{\beta'}$ provided in Eq. (109), the dressed S matrix $S_{\beta'}(\iota q_{F\beta'})$ and corresponding phase shift $Q_{\beta'}(\iota q_{F\beta'})/2$ depend on both the scatterer bare momentum $\iota q_{F\beta'}$ and the momentum q of the $\beta = c, s1$ branch-line scattering center.

On the other hand, near a spectral-function singularity (k, ω)-plane point (P_0, ω_0) associated with a transition to an excited energy eigenstate generated by processes (A) and (B) whose spectra, Eq. (97), have the form $\delta E^{\odot}_{l_o} = \omega_0$ and $\delta P^{\odot}_{l_o} = P_0$, the spectral-function behavior is found to be [7–9],

$$\begin{aligned} B(k,\,\omega) &\propto (\omega-\omega_0)^{\zeta}\,; \quad (\omega-\omega_0) \geq 0\,, \\ \zeta &= -2 + \sum_{\beta'=c,s1}\sum_{\iota=\pm1} 2\Delta^{\iota}_{\beta'}\,; \quad \omega \neq \pm v_\beta\,(k-P_0)\,, \quad \beta = c, s1\,, \\ \zeta &= -1 - 2\Delta^{\mp}_{\beta} + \sum_{\beta'=c,s1}\sum_{\iota=\pm1} 2\Delta^{\iota}_{\beta'}\,; \quad \omega \approx \pm v_\beta\,(k-P_0)\,, \quad \beta = c, s1\,. \end{aligned} \tag{113}$$

Here v_β is the $\beta = c, s1$ band Fermi velocity given in Eq. (A23) of Appendix A. The expressions provided in Eq. (113) apply to the finite-weight region above the singular point (P_0, ω_0).

The general dynamical and spectral function expressions provided in in Eqs. (112) and (113) refer to the vicinity of both low-energy and finite-energy spectral features. It is shown in Ref. [8] that in the limit of low excitation energy such expressions recover those provided by conformal-field theory. As shown in that reference, for low-energy eigenstates of $S_\eta > 0$ and $S_s \geq 0$ ground states the four functionals $2\Delta^\pm_c$ and $2\Delta^\pm_{s1}$ of Eq. (109) are the conformal dimensions of the $c, \pm$ and $s, \pm$ primary fields. Within conformal-field theory, those control the expressions of the low-energy correlation-function exponents. At low-energy the finite-size corrections associated with the energy excitation spectrum can be mapped onto a semi-direct product of two independent Virasoro algebras, each having central charge equal to one [54].

We have the following comments about the general PDT spectral-weight expressions given above in terms of the β phase shift $Q_\beta(q_j)/2$ and thus of the corresponding dressed S matrix $S_\beta(q_j) = e^{iQ_\beta(q_j)}$ of the pseudofermion scattering theory introduced in this paper:

– The important PDT $\beta = c, s1$ spectral function weight $A^{(0,0)}_\beta$ given in Eq. (105) involves products $\prod_{i=1}^{N_{a_\beta}}$ over all $N_{a_\beta} = [N_\beta + N^h_\beta]$ corresponding $\beta = c, s1$ band scatterers of the excited state. That includes both those that pre-existed in the initial ground state and the scatterers created under the transition to the excited state. This confirms that the 1D Hubbard model dynamical and spectral properties are within the pseudofermion scattering theory controlled by microscopic processes involving the scattering of all such $\beta = c, s1$ scatterers with the scattering centers created under the transitions to the ground state. In contrast, both the traditional scattering theories of Ref. [40] and Refs. [41, 42], respectively, consider only the few phase shifts associated with scatterers created under the transition to the excited state. Thus they do not account for most of the microscopic processes that control the model dynamical and spectral properties.

– For electronic densities $n \neq 1$ the four functionals of Eq. (109) control the exponents in the dynamical and spectral function expressions given in Eqs. (112) and (113), which refer to the vicinity of well-defined $\beta = c, s1$ branch lines and points, respectively. For instance, c and $s1$ branch lines as those of the expression provided in Eq. (112) occur in the case of the one-electron spectral functions [9]. Under the processes (A) and (B) that generate the one-electron addition LHB excited states (k) and (n) (and one-electron removal excited states (k) and (n)) of Table V of Appendix C, two scattering centers are created: one c pseudofermion (and one c pseudofermion hole) and one $s1$ pseudofermion hole. For the particular processes (A) and (B) that generate the c branch lines (and $s1$ branch lines), the c scattering center (and $s1$ scattering center) is created at a bare momentum q away from the corresponding c (and $s1$) Fermi points. Thus only the other $s1$ scattering center (and c scattering center) is created at a $s1$ (and c) Fermi point. This reveals that out of the four scatterers with phase shifts $Q_c(\pm q_{Fc})/2$ and $Q_{s1}(\pm q_{Fc})/2$, which appear in the expression of Eq. (109) for the corresponding functionals $2\Delta^\pm_c$ and $2\Delta^\pm_{s1}$, respectively, that control the branch-line momentum-dependent exponent $\zeta(k)$ of Eq. (112), three pre-existed in the initial ground state. On the other hand, the scattering theory of Ref. [40] only provides the phase shift of one of such four scatterers: That of the spinon corresponding to the $s1$ scatterer (and holon or antiholon corresponding to the c scatterer) created under the transition to the ground state at a $s1$ (and c) Fermi point, which also plays the role of scattering center. (The same applies to the phase shifts of Refs. [41, 42], which in addition refer to half filling whereas the PDT expressions are valid for the model metallic phase.)

Thus not only the $\beta = c, s1$ spectral function weight $A_{\beta}^{(0,0)}$ given in Eq. (105) involves the phase shifts of both all $\beta = c, s1$ scatterers that pre-existed in the ground state and those created under the transition to the excited state, as also, in the case of one-electron excited states, out of the four phase shifts that control the branch-line momentum-dependent exponent $\zeta(k)$ given in Eq. (112), three phase shifts refer to $\beta = c, s1$ scatterers that pre-existed in the ground state. Such results confirm the unsuitability of the traditional holon and spinon scattering theories to describe the model dynamical and spectral properties.

After the PDT was introduced, a set of novel methods have been developed to also tackle the finite-energy physics of 1D quantum fluids, beyond the low-energy Luttinger-liquid limit. Some of such methods have been very recently reviewed in Ref. [14]. They combine phenomenology built on the ideas of the Fermi-edge singularity and the Fermi-liquid theory, perturbation theory in the interaction strength, and new ways of treating finite-size properties of integrable models. Their application to specific 1D correlated models often involve the use of suitable numerical techniques, as those used in the studies of Ref. [17] for the 1D Hubbard model. The numerical results for the dependence of the one-electron removal spectral-function exponents on the excitation momentum reported in Section VIII of that reference are for electronic density $n = 0.59$, interaction values $u = U/4t = 0.25, 1.225, 2.5$, and the whole range of momentum in full quantitative agreement with those previously obtained by use of the PDT [9, 23, 24]. (The analytical justification of the equivalence of the PDT expressions for such exponents and the corresponding numerical results of Ref. [17] is an issue that will be addressed elsewhere.)

VII. CONCLUDING REMARKS

In this paper it is found that the 1D Hubbard model natural scatterers and scattering centers that emerge from the operator formulation introduced in the first paper are the spin-less and η-spin-less c pseudofermions, the spin-neutral composite $s\nu$ pseudofermions, and the η-spin-neutral composite $\eta\nu$ pseudofermions. The corresponding $\beta = c, \alpha\nu$ pseudofermion dressed S matrix has dimension *one*. Since for the theory initial ground states only the c and $s1$ pseudofermions have finite occupancies, such scatterers phase shifts and dressed S matrices play a major role in the model physics.

There is a β pseudofermion scattering theory for each initial ground state and corresponding PS. The theory is valid for PSs of initial ground states with arbitrary values of the electronic density n and spin density m. Within each of such large subspaces, the c pseudofermions, $s\nu$ pseudofermions, and $\eta\nu$ pseudofermions emerge from the c fermions, $s\nu$ fermions, and $\eta\nu$ fermions, respectively, considered in the first paper, through a well-defined unitary transformation. Under each ground-state - excited-state transition, the β fermions discrete momentum values are slightly shifted, what leads to the corresponding β pseudofermions discrete canonical momentum values, which are phase-shift dependent. Otherwise, the β fermions and β pseudofermions have exactly the same properties.

As shown in the first paper, the present object operator formulation accounts for the representations of the model global symmetry algebra representations. Furthermore, the corresponding scatterers and scattering centers have been constructed from rotated-electron configurations inherently to profiting from the effects of the conservation of an infinite number of conservation laws associated with the model integrability. Indeed, for such scatterers and scattering centers there is only zero-momentum forward scattering, in contrast to the underlying involved non-perturbative many-electron interactions.

In terms of the β pseudofermion zero-momentum forward scattering, the quantities associated with the spectral and dynamical properties of the model $n \neq 1$ metallic phase are naturally controlled for both low and finite energies and finite values of the on-site repulsion by the corresponding dressed S matrices and phase shifts. Our results reveal that the transport of spin and charge is also controlled by such phase shifts.

The "in" and "out" states of the pseudofermion scattering theory are exact excited energy eigenstates. They can be written as a simple direct product of "in" asymptote and "out" asymptote one-pseudofermion scattering states, respectively. Such a property combined with the also simple form obtained for the β pseudofermion and β pseudofermion hole dressed S matrices is behind the suitability of the present representation to the description of the finite-energy spectral and dynamical properties of the model metallic phase. In this paper it has been confirmed that the pseudofermion scattering theory describes the microscopic processes behind the PDT [7, 8]. Such processes control the exotic transport of spin and charge. Rather than in terms of electronic processes, the microscopic processes that determine the spectral and transport properties of the 1D Hubbard model are simple in terms of β pseudofermion zero-momentum forward scattering.

In this paper we have also clarified the relation of the pseudofermion scattering theory to both the traditional spinon and holon scattering theories of Ref. [40] and Refs. [41–43], respectively. There is no contradiction whatsoever between those and the pseudofermion scattering theory. The degeneracy of the excited energy eigenstates plays a central role in the determination of the subspaces where distinct choices of scattering states and thus of different elementary objects is allowed. The present pseudofermion scattering theory is normal ordered relative to the electron vacuum whereas the traditional holon and spinon descriptions of Ref. [40] and Refs. [41–43] are normal ordered relative to an initial $S_s = 0$ ground state of arbitrary electronic density and the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state, respectively. That the ground-state normal-ordered traditional holon and spinon scattering theories consider the few scatterers that are created under the transitions to the excited states was found to limit their applicability to the study of the model dynamical and spectral properties. Indeed, they lack microscopic processes associated with the phase shifts of the scatterers that pre-exist in the initial ground state, which play an active role in such properties.

Several properties predicted by the 1D Hubbard model were observed in low-dimensional complex materials [25]. The investigations presented in Refs. [9, 22–24] confirm that the PDT describes successfully the unusual finite-energy spectral features observed by angle-resolved photoelectron spectroscopy in quasi-1D organic metals. The results of this paper contribute to the further understanding of the non-perturbative microscopic scattering mechanisms behind these properties.

Acknowledgments

We thank N. Andrei for careful reading of the manuscript and fruitful discussions, S.-J. Gu, H. Q. Lin, A. Moreno, A. Muramatsu, K. Penc, and R. G. Pereira for illuminating discussions and the hospitality and support of the Beijing Computational Science Research Center. J. M. P. C. thanks the hospitality of the Institut für Theoretische Physik III, Universität Stuttgart, and the financial support by the Portuguese FCT under SFRH/BSAB/1177/2011, German transregional collaborative research center SFB/TRR21 and Max Planck Institute for Solid State Research.

Appendix A: Generalization of the thermodynamic BA equations to the 1D Hubbard model in a uniform vector potential

Here some general results useful for the studies of the paper are presented. That includes extension of the first paper to the 1D Hubbard model in a uniform vector potential. Indeed, the thermodynamic BA equations considered in that paper may be extended to the 1D Hubbard model in a uniform vector potential, $A_{x_1}\,\vec{e}_{x_1}$ [51, 55]. In that general case the Hamiltonian $\hat{H}$ of Eqs. (4) and (5) becomes,

$$\begin{aligned}
\hat{H} &= \hat{H}_{symm} - \sum_{\beta=\eta,s} \mu_\alpha\, \hat{S}_\alpha^{x_3} \, , \\
\hat{H}_{symm} &= t\,[\hat{T}(\phi_\uparrow,\phi_\downarrow) + 4u\,\hat{V}_D] \, , \\
\hat{T}(\phi_\uparrow,\phi_\downarrow) &= -\sum_{\sigma=\uparrow,\downarrow}\sum_{j=1}^{N_a}\left[c_{j,\sigma}^\dagger\, c_{j+1,\sigma}\, e^{i\phi_\sigma/N_a} + c_{j+1,\sigma}^\dagger\, c_{j,\sigma}\, e^{-i\phi_\sigma/N_a}\right] \, .
\end{aligned} \tag{A1}$$

Here $\hat{V}_D$ is the electron on-site repulsion operator whose expression is provided in Eq. (5). The uniform vector potential does not alter the form of that operator. On the other hand, it modifies the kinetic energy operator by the usual Peierls phase factor.

Such an extension is needed for the studies of this paper presented in Section V. Otherwise, only the corresponding $\phi_\uparrow = \phi_\downarrow = 0$ thermodynamic BA equations are required. For the case of LWS Bethe states, the thermodynamic BA equations of the Hamiltonian given in Eq. (A1) are slightly changed relative to those provided in the first paper. Using the same notations and definitions as in that paper, they read [51, 55],

$$\begin{aligned}
k_c(q_j) &= q_j + \frac{\phi_\uparrow}{N_a} - \frac{2}{N_a}\sum_{\nu=1}^{\infty}\sum_{j'=1}^{N_{a_{s\nu}}} N_{s\nu}(q_{j'})\arctan\left(\frac{\sin k_c(q_j) - \Lambda_{s\nu}(q_{j'})}{\nu u}\right) \\
&- \frac{2}{N_a}\sum_{\nu=1}^{\infty}\sum_{j'=1}^{N_{a_{\eta\nu}}} N_{\eta\nu}(q_{j'})\arctan\left(\frac{\sin k_c(q_j) - \Lambda_{\eta\nu}(q_{j'})}{\nu u}\right) \; ; \quad j = 1, ..., N_a \, ,
\end{aligned} \tag{A2}$$

and

$$\begin{aligned}
k_{\alpha\nu}(q_j) &= q_j - 2\nu\left(\frac{\phi_{\uparrow} - e^{i\pi\delta_{\alpha,\eta}}\,\phi_{\downarrow}}{2N_a}\right) - \frac{2e^{i\pi\delta_{\alpha,\eta}}}{N_a}\sum_{j'=1}^{N_a} N_c(q_{j'})\arctan\left(\frac{\Lambda_{\alpha\nu}(q_j) - \sin k_c(q_{j'})}{\nu u}\right) \\
&+ \frac{1}{N_a}\sum_{\nu'=1}^{\infty}\sum_{j'=1}^{N_{a_{\alpha\nu'}}} N_{\alpha\nu'}(q_{j'})\Theta_{\nu,\,\nu'}\left(\frac{\Lambda_{\alpha\nu}(q_j) - \Lambda_{\alpha\nu'}(q_{j'})}{u}\right)\,, \\
\nu &= 1,...,\infty\,, \quad j = 1,...,N_{a_{\alpha\nu}}\,, \quad \alpha = \eta, s\,, \qquad \text{(A3)}
\end{aligned}$$

where

$$k_{\alpha\nu}(q_j) = \delta_{\alpha,\eta}\,2\,\mathrm{Re}\,\{\arcsin(\Lambda_{\eta\nu}(q_j) + i\nu u)\}\,, \quad \nu = 1,...,\infty\,, \quad j = 1,...,N_{a_{\alpha\nu}}\,, \quad \alpha = \eta, s\,, \qquad \text{(A4)}$$

is the $\alpha\nu$ rapidity-momentum functional, which vanishes for $\alpha = s$, and the function $\Theta_{\nu,\nu'}(x)$ is given in Eq. (B16) of Appendix B.

Alike in the first paper, our BA representation refers to LWSs of both the spin and η-spin $SU(2)$ algebras. For such Bethe states the numbers,

$$M^{un}_{\eta,-1/2} = S_\eta - \frac{1}{2}(N_a - N) = 0, 1, ..., 2S_\eta\,; \qquad M^{un}_{s,-1/2} = S_s - \frac{1}{2}(N_{\uparrow} - N_{\downarrow}) = 0, 1, ..., 2S_s\,, \qquad \text{(A5)}$$

of η-spin projection $-1/2$ unbound η-spinons and spin projection $-1/2$ unbound spinons, respectively, vanish. Moreover, we use the same notation, $|\Psi_{l_o,l_\triangle,u}\rangle$, for the energy eigenstates as in the first paper. Within such a notation, the Bethe states are written as $|\Psi_{l_o,l^0_\triangle,u}\rangle$. A non-LWS energy eigenstate $|\Psi_{l_o,l_\triangle,u}\rangle$ is generated from the corresponding Bethe state $|\Psi_{l_o,l^0_\triangle,u}\rangle$ as follows,

$$|\Psi_{l_o,l_\triangle,u}\rangle = \prod_{\alpha=\eta,s}\left[\frac{1}{\sqrt{\mathcal{C}_\alpha}}(\hat{S}^{+}_{\alpha})^{M^{un}_{s,-1/2}}\right]|\Psi_{l_o,l^0_\triangle,u}\rangle\,. \qquad \text{(A6)}$$

Here,

$$\begin{aligned}
\mathcal{C}_\alpha &= \langle\Psi_{l_o,l^0_\triangle,u}|(\hat{S}^{-}_{\alpha})^{M^{un}_{s,-1/2}}(\hat{S}^{+}_{\alpha})^{M^{un}_{s,-1/2}}|\Psi_{l_o,l^0_\triangle,u}\rangle \\
&= [M^{un}_{\alpha,-1/2}!]\prod_{j'=1}^{M^{un}_{\alpha,-1/2}}[\,M^{un}_{\alpha} + 1 - j'\,]\,, \quad M^{un}_{\alpha,-1/2} = 1, ..., M^{un}_{\alpha}\ \ , \alpha = \eta, s\,, \qquad \text{(A7)}
\end{aligned}$$

are normalization constants, $\hat{S}^{+}_{\alpha}$ and $\hat{S}^{-}_{\alpha}$ are the η-spin ($\alpha = \eta$) and spin ($\alpha = s$) off-diagonal generators considered in the first paper, $M^{un}_{\alpha} = 2S_\alpha$ denotes the total number of unbound η-spinons ($\alpha = \eta$) and unbound spinons ($\alpha = s$), and $l_\triangle$ and $l^0_\triangle$ stand for the set of numbers $[2S_c, S_\eta, S_s, M^{un}_{\eta,-1/2}, M^{un}_{s,-1/2}]$ and $[2S_c, S_\eta, S_s, 0, 0]$, respectively. Within our notation, $l^0_\triangle$ refers to limiting values of the general index $l_\triangle$ such that $M^{un}_{\eta,-1/2} = M^{un}_{s,-1/2} = 0$, which are those of the Bethe states. The triangle index $\triangle$ in $l_\triangle$ and $l^0_\triangle$ symbolizes the three symmetries: The c hidden $U(1)$ symmetry associated with the number $2S_c$ and the two $SU(2)$ symmetries associated with the numbers S_η, n_η and S_s, n_s, respectively, in $[2S_c, S_\eta, S_s, n_\eta, n_s]$. Moreover, l_o stands for all remaining quantum numbers beyond $l_\triangle$ needed to uniquely define an energy eigenstate $|\Psi_{l_o,l_\triangle,u}\rangle$.

The BA solution of the Hubbard model in a uniform vector potential, Eq. (A1), provides as well the energy eigenvalues $E_{l_o,l^0_\triangle}(\phi_{\uparrow}, \phi_{\downarrow})$ of the Bethe states $|\Psi_{l_o,l^0_\triangle,u}\rangle$, which are given by,

$$\begin{aligned}
E_{l_o,l^0_\triangle} &= E_{symm} + \mu\, M^{un}_{\eta,+1/2} + \mu_B\, H\, M^{un}_{s,+1/2}\,, \\
E_{symm} &= -2t\sum_{j=1}^{N_a}[N_c(q_j)\ (\cos k(q_j) + u) - u/2] \\
&+ 4t\sum_{\nu=1}^{\infty}\sum_{j=1}^{N_{a_{\eta\nu}}} N_{\eta\nu}(q_j)\left[\mathrm{Re}\left\{\sqrt{1 - (\Lambda_{\eta\nu}(q_j) + i\nu\, u)^2}\right\} - \nu\, u\right]\,. \qquad \text{(A8)}
\end{aligned}$$

For the non-LWSs the unbound η-spinon number $M^{un}_{\eta,+1/2}$ and unbound spinon number $M^{un}_{s,+1/2}$ appearing in the first expression given here are replaced by $[M^{un}_{\eta,+1/2} - M^{un}_{\eta,-1/2}]$ and $[M^{un}_{s,+1/2} - M^{un}_{s,-1/2}]$, respectively. The dependence on

$\phi_\uparrow$ and $\phi_\downarrow$ of this general energy spectrum occurs through that of the rapidity functions appearing in its expression. The latter $\phi_\uparrow$ and $\phi_\downarrow$ dependence is uniquely defined by Eqs. (A2) and (A3).

Within the $N_a \rightarrow \infty$ limit that the thermodynamic BA equations, Eqs. (A2)-(A4), refer to, it is often convenient to replace the β fermion discrete momentum values q_j, such that $q_{j+1} - q_j = 2\pi/N_a$, by corresponding continuous momentum variables q. In the absence of a vector potential, $\phi_\uparrow = \phi_\downarrow = 0$ in Eqs. (A2)-(A4), those belong to domains $q \in [-q_\beta, +q_\beta]$, whose limiting absolute values q_β read [5],

$$\begin{aligned} q_c &= \pi \, , \\ q_{\alpha\nu} &= \frac{\pi}{N_a}(N_{a_{\alpha\nu}} - 1) \ \text{ for } N_{a_{\alpha\nu}} \text{ odd} \, , \\ &= \frac{\pi}{N_a} N_{a_{\alpha\nu}} \ \text{ for } N_{a_{\alpha\nu}} \text{ even} \, . \end{aligned} \tag{A9}$$

The number N_{a_c} of c band discrete momentum values and that of $\alpha\nu$ band discrete momentum values $N_{a_{\alpha\nu}}$ appearing here are found in the first paper to read,

$$\begin{aligned} N_{a_c} = [N_c + N_c^h] = N_a \, , \\ N_{a_{\alpha\nu}} \ &= \ [N_{\alpha\nu} + N_{\alpha\nu}^h] \, . \end{aligned} \tag{A10}$$

where $N_c = 2S_c$ and $N_{\alpha\nu}$ denote the number of occupied c and $\alpha\nu$ band discrete momentum values, respectively. Those of unoccupied values are given by $N_c^h = 2S_c^h = [N_a - 2S_c]$ and,

$$N_{\alpha\nu}^h = [2S_\alpha + 2\sum_{\nu'=\nu+1}^{\infty} (\nu' - \nu) N_{\alpha\nu'}] = [M_\alpha^{un} + 2\sum_{\nu'=\nu+1}^{\infty} (\nu' - \nu) N_{\alpha\nu'}] \, , \ \ \alpha = \eta, s \, , \tag{A11}$$

respectively. For the $\beta = \alpha\nu$ branches the β discrete-momentum values distribution is symmetrical and bound by the momentum values $\pm q_{\alpha\nu}$ with $q_{\alpha\nu}$ given in Eq. (A9). On the other hand, if one accounts for corrections of order $1/N_a$, one finds that for the c band the q range is given by $q \in [q_c^-, q_c^+]$ where the limiting values $q_c^\pm$ are provided in Eqs. (B.15)-(B.17) of Ref. [33].

For the ground states, the limiting momentum values of Eq. (A9) have simple expressions. For the densities in the ranges considered in this section, they read,

$$q_c = \pi \, ; \quad\ q_{s1} = k_{F\uparrow} \, ; \quad\ q_{s\nu} = [k_{F\uparrow} - k_{F\downarrow}] \, , \quad \nu > 1 \, ; \quad\ q_{\eta\nu} = [\pi - 2k_F] \, . \tag{A12}$$

Here we have again ignored the corrections of order $1/N_a$.

Concerning β fermion occupancies, ground states are described by compact c and $s1$ fermion finite occupancies for momenta in the ranges $q \in [-q_{F\beta}, +q_{F\beta}]$ where $\beta = c, s1$. On the other hand, the $\alpha\nu \neq s1$ branches have vanishing ground-state occupancy. For the above ground states with densities in the ranges $n \in [0, 1]$ and $m \in [0, n]$, the c and $s1$ Fermi momenta are given by,

$$q_{Fc} = 2k_F \, ; \quad\ q_{Fs1} = k_{F\downarrow} \, . \tag{A13}$$

Here we have ignored corrections of order $1/N_a$, which are provided in Eqs. (C.4)-(C.11) of Ref. [33].

Alike in the first paper, we denote by $k_c^0(q_j)$ and $\Lambda_\beta^0(q_j)$ the specific rapidity-function solutions of the thermodynamic integral equations given in Eqs. (A2)-(A4) for $\phi_\uparrow = \phi_\downarrow = 0$ that refer to a ground state. Those play an important role both in the PDT of Refs. [7, 8] and pseudofermion scattering theory introduced in this paper. Upon suitable manipulations of such thermodynamic BA equations, the ground-state functions $k_c^0(q_j)$ and $\Lambda_\beta^0(q_j)$ may be defined in terms of their inverse functions $q_j = q_j(\Lambda_{\beta,j}^0)$ as,

$$\begin{aligned} q_j &= F_\beta^0(\Lambda_{\beta,}^0(q_j)) + (-1)^{\delta_{\beta,\eta\nu}} \int_{-Q}^{Q} dk \, \bar{\Phi}_{c,\beta}\left(\frac{\sin k}{u}, \frac{\Lambda_\beta^0(q_j)}{u}\right) \, , \quad j = 1, ..., N_{a_\beta} \, , \\ F_c^0(\Lambda_c^0(q_j)) &= \arcsin(\Lambda_c^0(q_j)) = k_c^0(q_j) \, , \\ F_{\alpha\nu}^0(\Lambda_{\alpha\nu}^0(q_j)) &= \delta_{\alpha,\eta} \, 2\text{Re}\left[\arcsin(\Lambda_{\eta\nu}^0(q_j) + i\nu \, u)\right] \, . \end{aligned} \tag{A14}$$

Here $N_{a_\beta} = N_a$ for $\beta = c$. The number $N_{a_{\alpha\nu}}$ is for the $\beta = \alpha\nu$ branches given in Eqs. (A10) and (A11). The functions $\bar{\Phi}_{c,\beta}(r, r')$ also appearing here are defined in Eqs. (B7)- (B9) of Appendix B. Moreover, the parameter Q in the above equation and the related parameters B, r_c^0, and r_{s1}^0 appearing in other quantities considered in this paper may be expressed in terms of the ground-state rapidity functions $k_c^0(q)$ and $\Lambda_{s1}^0(q)$ at the corresponding c and $s1$ Fermi points of Eq. (A13), respectively, as follows,

$$\begin{aligned} Q &\equiv k_c^0(2k_F)\,; \qquad B \equiv \Lambda_{s1}^0(k_{F\downarrow})\,, \\ r_c^0 &= \frac{\sin Q}{u}\,; \qquad r_s^0 = \frac{B}{u}\,. \end{aligned} \tag{A15}$$

For the $n=1$ and $m=0$ absolute ground state the $\beta \neq c, s1$ bands do not exist and the equations given in Eq. (A14) have an analytical solution for the $\beta = c, s1$ branches. For that ground state the rapidity functions $k_c^0(q_j)$ (where $\Lambda_c^0(q_j) = \sin k_c^0(q_j)$) and $\Lambda_{s1}^0(q_j)$ are defined in terms of their inverse rapidity functions as follows,

$$\begin{aligned} q_j &= k_c^0(q_j) + 2\int_0^\infty d\omega \frac{\sin(\omega\,\sin k_c^0(q_j))}{\omega\,(1+e^{2\omega\,u})}\,J_0(\omega)\,, \quad j = 1, ..., N_a\,, \\ q_j &= \int_0^\infty d\omega \frac{\sin(\omega\,\Lambda_{s1}^0(q_j))}{\omega\,\cosh(\omega\,u)}\,J_0(\omega)\,, \quad j = 1, ..., N_{a_{s1}}\,. \end{aligned} \tag{A16}$$

Here $J_0(\omega)$ is a Bessel function.

To first order in the β fermion momentum-distribution function deviations, the general energy spectrum of the Hamiltonian (4) is of the form [5],

$$\begin{aligned} \delta E_{l_o,l_\triangle} &= \sum_\beta \sum_{j=1}^{N_{a_\beta}} \varepsilon_\beta(q_j)\delta N_\beta(q_j) + 2\mu\,M_{\eta,-1/2}^{un} + 2\mu_B\,H\,M_{s,-1/2}^{un}\,, \\ &= \sum_{\beta=c,s1} \sum_{j=1}^{N_{a_\beta}} \varepsilon_\beta(q_j)\delta N_\beta(q_j) + \sum_{\beta\nu\neq s1} \sum_{j=1}^{N_{a_{\alpha\nu}}} \varepsilon_{\alpha\nu}^0(q_j)\delta N_{\alpha\nu}(q_j) \\ &+ \; 2|\mu|\,M_{\eta,-1/2} + 2\mu_B\,|H|\,[\delta M_{s,-1/2} - \delta N_{s1}]\,. \end{aligned} \tag{A17}$$

The β fermion momentum-distribution function deviations appearing here are given by,

$$\delta N_\beta(q_j) = N_\beta(q_j) - N_\beta^0(q_j)\,, \quad j = 1, ..., N_{a_\beta}\,, \quad \beta = c, \alpha\nu\,, \tag{A18}$$

where $N_\beta(q_j)$ and $N_\beta^0(q_j)$ are the corresponding excited-state and ground-state β fermion momentum-distribution functions, respectively.

In the expressions given in Eq. (A17), $M_{\eta,\pm 1/2}^{un}$ and $M_{s,\pm 1/2}^{un}$ are the numbers of η-spin projection $\pm 1/2$ η-spinons and spin projection $\pm 1/2$ spinons, respectively, given in Eq. (3). (The numbers $M_{\eta,-1/2}^{un}$ and $M_{s,-1/2}^{un}$ are those of Eq. (A5).) Related important conserving numbers for the finite-energy physics are that of rotated-electron doubly occupied sites, D_r, and spin-down rotated-electron singly occupied sites whose rotated electrons are not associated with the $s1$ fermions, S_r. Such conserving numbers read,

$$\begin{aligned} D_r \equiv M_{\eta,-1/2} &= [M_{\eta,-1/2}^{un} + M_\eta^{bo}/2] = [M_{\eta,-1/2}^{un} + \sum_{\nu=1}^\infty \nu N_{\eta\nu}]\,, \\ S_r &\equiv [M_{s,-1/2} - N_{s1}] = [M_{s,-1/2}^{un} + M_s^{bo}/2 - N_{s1}] = [M_{s,-1/2}^{un} + \sum_{\nu=2}^\infty \nu N_{s\nu}]\,. \end{aligned} \tag{A19}$$

The energy dispersions $\varepsilon_\beta^0(q_j)$ also appearing in Eq. (A17) can be expressed in terms of the functions $\bar{\Phi}_{c,\beta}(r,r')$ defined by Eqs. (B7)- (B9) of Appendix B and of the initial ground-state rapidity functions $\Lambda_\beta^0(q_j)$ (with $\Lambda_c^0(q_j) \equiv \sin k_c^0(q_j)$ for $\beta = c$) defined by their inverse functions in Eq. (A14) as follows,

$$\begin{aligned} \varepsilon_\beta^0(q_j) &= E_\beta^0(q_j) + 2t\int_{-Q}^{Q} dk\,\bar{\Phi}_{c,\beta}\left(\frac{\sin k}{u}, \frac{4t\Lambda_\beta^0(q_j)}{U}\right)\sin k\,, \quad j = 1, ..., N_{a_\beta}\,, \\ E_c^0(q_j) &= -\frac{U}{2} - 2t\cos k_c^0(q_j)\,, \\ E_{\alpha\nu}^0(q_j) &= \delta_{\alpha,\eta}\left\{-\nu\,U + 4t\mathrm{Re}\left[\sqrt{1 - (\Lambda_{\eta\nu}^0(q_j) + i\nu\,u)^2}\right]\right\}\,. \end{aligned} \tag{A20}$$

The related energy dispersions $\varepsilon_{\beta}(q_j)$ appearing in Eq. (A17) for the $\beta = c, s1$ branches can be defined for all branches. They are given by [5],

$$
\begin{aligned}
\varepsilon_c(q_j) &= \varepsilon_c^0(q_j) - \varepsilon_c^0(q_{Fc}) = \varepsilon_c^0(q_j) + (1-\delta_{n,1})\,|\mu| + \delta_{n,1}\,\mu^0 - \mu_B\,|H|\,, \\
\varepsilon_{s1}(q_j) &= \varepsilon_{s1}^0(q_j) - \varepsilon_{s1}^0(q_{Fs1}) = \varepsilon_{s1}^0(q_j) + 2\mu_B\,|H|\,, \\
\varepsilon_{s\nu}(q_j) &= \varepsilon_{s\nu}^0(q_j) + 2\nu\mu_B\,|H|\,, \\
\varepsilon_{\eta\nu}(q_j) &= \varepsilon_{\eta\nu}^0(q_j) + (1-\delta_{n,1})\,2\nu|\mu| + \delta_{n,1}\,2\nu\mu^0\,,
\end{aligned}
\tag{A21}
$$

where $2\mu^0$ is the $n = 1$ Mott-Hubbard gap whose limiting behaviors are given in Eq. (38). The zero-energy level of the energy bands in Eq. (A17) is such that,

$$
\begin{aligned}
\varepsilon_c(\pm 2k_F) &= \varepsilon_{s1}(\pm k_{F\downarrow}) = 0\,, \\
\varepsilon_{\eta\nu}^0(\pm[\pi - 2k_F]) &= \varepsilon_{s\nu}^0(\pm[k_{F\uparrow} - k_{F\downarrow}]) = 0\,.
\end{aligned}
\tag{A22}
$$

Hence for the $\alpha\nu \neq s1$ bands it refers to the limiting momenta of Eq. (A12) whereas for the c and $s1$ bands it corresponds to the $\beta = c, s1$ Fermi momenta given in Eq. (A13).

The corresponding β fermion group velocities read,

$$
\begin{aligned}
v_{\beta}(q) &= \frac{\partial\varepsilon_{\beta}(q)}{\partial q} = \frac{\partial\varepsilon_{\beta}^0(q)}{\partial q}\,, \quad \beta = c, \eta\nu, s\nu\,, \quad \nu = 1, ..., \infty\,, \\
v_{\beta} &\equiv v_{\beta}(q_{F\beta})\,, \quad \beta = c, s1\,.
\end{aligned}
\tag{A23}
$$

Appendix B: Two-pseudofermion phase shifts

Here we first derive the integral equations that define the two-pseudofermion phase shifts $\pi\,\bar{\Phi}_{\beta,\beta'}(r, r')$ on the right-hand side of Eq. (14) for densities whose ranges obey the inequalities $0 < n < 1$ and $0 < m < n$. To achieve that goal, we solve the thermodynamic BA equations given Eqs. (A2)-(A4) of Appendix A for $\phi_{\uparrow} = \phi_{\downarrow} = 0$ up to first order in the deviations $\delta N_{\beta}(q_j)$ provided in Eq. (A18) of that appendix. This is accomplished on using in them rapidity functionals of form provided in Eq. (8) with the canonical momentum $\bar{q}_j = \bar{q}(q_j)$ defined by Eqs. (10) and (11). Such a procedure leads to the expression given in Eq. (14) with the two-pseudofermion phase shift $\pi\,\bar{\Phi}_{\beta,\beta'}(r, r')$ uniquely defined by the integral equations provided below.

A first group of two-pseudofermion phase shifts obey integral equations by their own. These equations read,

$$
\pi\,\bar{\Phi}_{s1,c}(r, r') = -\arctan(r - r') + \int_{-r_s^0}^{r_s^0} dr''\,G(r, r'')\,\pi\,\bar{\Phi}_{s1,c}(r'', r')\,,
\tag{B1}
$$

$$
\pi\,\bar{\Phi}_{s1,\eta\nu}(r, r') = -\frac{1}{\pi}\int_{-r_c^0}^{r_c^0} dr''\frac{\arctan\Bigl(\frac{r''-r'}{\nu}\Bigr)}{1 + (r - r'')^2} + \int_{-r_s^0}^{r_s^0} dr''\,G(r, r'')\,\pi\,\bar{\Phi}_{s1,\eta\nu}(r'', r')\,,
\tag{B2}
$$

and

$$
\begin{aligned}
\pi\,\bar{\Phi}_{s1,s\nu}(r, r') &= \delta_{1,\nu}\,\arctan\Bigl(\frac{r - r'}{2}\Bigr) + (1 - \delta_{1,\nu})\Bigl\{\arctan\Bigl(\frac{r - r'}{\nu - 1}\Bigr) + \arctan\Bigl(\frac{r - r'}{\nu + 1}\Bigr)\Bigr\} \\
&- \frac{1}{\pi}\int_{-r_c^0}^{r_c^0} dr''\frac{\arctan\Bigl(\frac{r''-r'}{\nu}\Bigr)}{1 + (r - r'')^2} + \int_{-r_s^0}^{r_s^0} dr''\,G(r, r'')\,\pi\,\bar{\Phi}_{s1,s1}(r'', r')\,.
\end{aligned}
\tag{B3}
$$

The parameters r_c^0 and r_s^0 appearing in such equations are given in Eq. (A15) of Appendix A and the kernel $G(r, r')$ reads,

$$
G(r, r') = -\frac{1}{2\pi}\left[\frac{1}{1 + ((r - r')/2)^2}\right]\left[1 - \frac{1}{2}\left(t(r) + t(r') + \frac{l(r) - l(r')}{r - r'}\right)\right]\,.
\tag{B4}
$$

Here

$$
t(r) = \frac{1}{\pi}\left[\arctan(r + r_c^0) - \arctan(r - r_c^0)\right]\,,
\tag{B5}
$$

and

$$l(r) = \frac{1}{\pi}\left[\ln(1+(r+r_c^0)^2) - \ln(1+(r-r_c^0)^2)\right] . \tag{B6}$$

The kernel defined in Eqs. (B4)-(B6) was first introduced in Ref. [48] within the low-energy two-component c and $s \equiv s1$ pseudoparticle theory studied in that reference.

A second group of two-pseudofermion phase shifts are expressed in terms of the basic functions given in Eqs. (B1)-(B3) as follows,

$$\pi\,\bar{\Phi}_{c,c}\,(r,r') = \frac{1}{\pi}\int_{-r_s^0}^{r_s^0} dr''\frac{\pi\,\bar{\Phi}_{s1,c}\,(r'',r')}{1+(r-r'')^2}\,, \tag{B7}$$

$$\pi\,\bar{\Phi}_{c,\eta\nu}\,(r,r') = -\arctan\Big(\frac{r-r'}{\nu}\Big) + \frac{1}{\pi}\int_{-r_s^0}^{r_s^0} dr''\frac{\pi\,\bar{\Phi}_{s1,\eta\nu}\,(r'',r')}{1+(r-r'')^2}\,, \tag{B8}$$

and

$$\pi\,\bar{\Phi}_{c,s\nu}\,(r,r') = -\arctan\Big(\frac{r-r'}{\nu}\Big) + \frac{1}{\pi}\int_{-r_s^0}^{r_s^0} dr''\frac{\pi\,\bar{\Phi}_{s1,s\nu}\,(r'',r')}{1+(r-r'')^2}\,. \tag{B9}$$

Finally, the remaining two-pseudofermion phase shifts can be expressed either in terms of the functions provided in Eqs. (B7)-(B9) only,

$$\pi\,\bar{\Phi}_{\eta\nu,c}\,(r,r') = \arctan\Big(\frac{r-r'}{\nu}\Big) - \frac{1}{\pi}\int_{-r_c^0}^{+r_c^0} dr''\frac{\pi\,\bar{\Phi}_{c,c}\,(r'',r')}{\nu[1+(\frac{r-r''}{\nu})^2]}\,, \tag{B10}$$

$$\pi\,\bar{\Phi}_{\eta\nu,\eta\nu'}\,(r,r') = \frac{1}{2}\,\Theta_{\nu,\nu'}(r-r') - \frac{1}{\pi}\int_{-r_c^0}^{+r_c^0} dr''\frac{\pi\,\bar{\Phi}_{c,\eta\nu'}\,(r'',r')}{\nu[1+(\frac{r-r''}{\nu})^2]}\,, \tag{B11}$$

$$\pi\,\bar{\Phi}_{\eta\nu,s\nu'}\,(r,r') = -\frac{1}{\pi}\int_{-r_c^0}^{+r_c^0} dr''\frac{\pi\,\bar{\Phi}_{c,s\nu'}\,(r'',r')}{\nu[1+(\frac{r-r''}{\nu})^2]}\,, \tag{B12}$$

or both in terms of the basic functions given in Eqs. (B1)-(B3) and of the phase shifts of Eqs. (B7)-(B9),

$$\pi\,\bar{\Phi}_{s\nu,c}\,(r,r') = -\arctan\Big(\frac{r-r'}{\nu}\Big)+\frac{1}{\pi}\int_{-r_c^0}^{r_c^0} dr''\frac{\pi\,\bar{\Phi}_{c,c}\,(r'',r')}{\nu[1+(\frac{r-r''}{\nu})^2]}-\int_{-r_s^0}^{r_s^0} dr''\pi\,\bar{\Phi}_{s1,c}\,(r'',r')\,\frac{\Theta^{[1]}_{\nu,1}(r-r'')}{2\pi}\,; \qquad \nu>1\,, \tag{B13}$$

$$\pi\,\bar{\Phi}_{s\nu,\eta\nu'}\,(r,r') = \frac{1}{\pi}\int_{-r_c^0}^{r_c^0} dr''\frac{\pi\,\bar{\Phi}_{c,\eta\nu'}\,(r'',r')}{\nu[1+(\frac{r-r''}{\nu})^2]} - \int_{-r_s^0}^{r_s^0} dr''\pi\,\bar{\Phi}_{s1,\eta\nu'}\,(r'',r')\,\frac{\Theta^{[1]}_{\nu,1}(r-r'')}{2\pi}\,; \qquad \nu>1\,, \tag{B14}$$

$$\pi\,\bar{\Phi}_{s\nu,s\nu'}\,(r,r') = \frac{1}{2}\,\Theta_{\nu,\nu'}(r-r') + \frac{1}{\pi}\int_{-r_c^0}^{r_c^0} dr''\frac{\pi\,\bar{\Phi}_{c,s\nu'}\,(r'',r')}{\nu[1+(\frac{r-r''}{\nu})^2]} - \int_{-r_s^0}^{r_s^0} dr''\pi\,\bar{\Phi}_{s1,s\nu'}\,(r'',r')\,\frac{\Theta^{[1]}_{\nu,1}(r-r'')}{2\pi}\,. \tag{B15}$$

In the above two-pseudofermion phase shift expressions, $\Theta_{\nu,\nu'}(x)$ is the function,

$$\begin{aligned}\Theta_{\nu,\nu'}(x) \;&=\; \delta_{\nu,\nu'}\Big\{2\arctan\Big(\frac{x}{2\nu}\Big)+\sum_{l=1}^{\nu-1}4\arctan\Big(\frac{x}{2l}\Big)\Big\} + (1-\delta_{\nu,\nu'})\Big\{2\arctan\Big(\frac{x}{|\,\nu-\nu'|}\Big)\\ &+\; 2\arctan\Big(\frac{x}{\nu+\nu'}\Big)+\sum_{l=1}^{\frac{\nu+\nu'-|\,\nu-\nu'|}{2}-1} 4\arctan\Big(\frac{x}{|\,\nu-\nu'|+2l}\Big)\Big\}\,,\end{aligned} \tag{B16}$$

and $\Theta^{[1]}_{\nu,\nu'}(x)$ is its derivative,

$$\begin{aligned}\Theta^{[1]}_{\nu,\nu'}(x) &= \frac{d\Theta_{\nu,\nu'}(x)}{dx} = \delta_{\nu,\nu'}\Big\{\frac{1}{\nu[1+(\frac{x}{2\nu})^2]} + \sum_{l=1}^{\nu-1}\frac{2}{l[1+(\frac{x}{2l})^2]}\Big\} + (1-\delta_{\nu,\nu'})\Big\{\frac{2}{|\nu-\nu'|[1+(\frac{x}{|\nu-\nu'|})^2]} \\ &+ \frac{2}{(\nu+\nu')[1+(\frac{x}{\nu+\nu'})^2]} + \sum_{l=1}^{\frac{\nu+\nu'-|\nu-\nu'|}{2}-1}\frac{4}{(|\nu-\nu'|+2l)[1+(\frac{x}{|\nu-\nu'|+2l})^2]}\Big\}\,. \qquad \text{(B17)}\end{aligned}$$

As discussed in Section IV B, the general two-pseudofermion phase-shift $\pi\,\Phi_{\beta,\beta'}(q_j,q_{j'})$ expression provided in Eq. (14) applies to initial ground states with densities in the ranges defined by the inequalities $0 < n < 1$ and $0 < m < n$. On the other hand, for $n = 1$ (and $m = 0$) ground states and $\beta = \eta\nu$ and/or $\beta' = \eta\nu'$ (and $\beta = s\nu \neq s1$ and/or $\beta' = s\nu' \neq s1$) the rapidity function $\Lambda^0_{\beta}(q_j)$ and/or $\Lambda^0_{\beta'}(q_{j'})$ in the argument of the phase-shift $\pi\,\bar{\Phi}_{\beta,\beta'}(r,r')$ on the right-hand side of Eq. (14) must be replaced by the corresponding excited-state function $\Lambda_{\beta}(q_j)$ and/or $\Lambda_{\beta'}(q_{j'})$, respectively. Fortunately, however, the $m \to 0$ (and $n \to 1$) limit of the above equations defining the two-pseudofermion phase-shifts $\pi\,\bar{\Phi}_{\beta,\beta'}(r,r')$ defines as well such phase shifts for an initial $m = 0$ (and $n = 1$) ground state.

We start by using such a property in the case of an initial $m = 0$ ground state with electronic density in the range $n \in [0,1]$. We consider the $m \to 0$ spin-density limit of the above expressions, which provides the correct two-pseudofermion phase-shift values. The rapidity two-pseudofermion phase shifts $\pi\,\bar{\Phi}_{\beta,\beta'}(r,r')$ are defined by the integral equations, Eqs. (B1)-(B15). By Fourier transforming such equations after accounting for that $B = \infty$ and thus $r^0_s = 4t\,B/U = \infty$ for u finite, we arrive to the following equations valid for $m \to 0$ and u finite,

$$\pi\,\bar{\Phi}_{c,c}(r,r') = -B(r-r') + \int_{-r^0_c}^{+r^0_c} dr''\,A(r-r'')\,\pi\,\bar{\Phi}_{c,c}(r'',r')\,, \qquad \text{(B18)}$$

$$\pi\,\bar{\Phi}_{c,s1}(r,r') = -\frac{1}{2}\arctan\Big(\sinh\Big(\frac{\pi}{2}(r-r')\Big)\Big) + \int_{-r^0_c}^{+r^0_c} dr''\,A(r-r'')\,\pi\,\bar{\Phi}_{c,s1}(r'',r')\,, \qquad \text{(B19)}$$

$$\begin{aligned}\pi\,\bar{\Phi}_{s1,c}(r,r') &= -\frac{1}{2}\arctan\Big(\sinh\Big(\frac{\pi}{2}(r-r')\Big)\Big) + \frac{1}{4}\int_{-r^0_c}^{+r^0_c} dr''\,\frac{\pi\,\bar{\Phi}_{c,c}(r'',r')}{\cosh\Big(\frac{\pi}{2}(r-r'')\Big)}\,; \qquad r \neq \pm\infty \\ &= -\frac{\mathrm{sgn}(r)\pi}{2\sqrt{2}}\,; \qquad r = \pm\infty\,, \qquad \text{(B20)}\end{aligned}$$

$$\begin{aligned}\pi\,\bar{\Phi}_{s1,s1}(r,r') &= B(r-r') + \frac{1}{4}\int_{-r^0_c}^{+r^0_c} dr''\,\frac{\pi\,\bar{\Phi}_{c,s1}(r'',r')}{\cosh\Big(\frac{\pi}{2}(r-r'')\Big)}\,; \quad r \neq \pm\infty \\ &= \frac{\mathrm{sgn}(r)\pi}{2\sqrt{2}}\,; \qquad r = \pm\infty\,, \qquad r' \neq r \\ &= [\mathrm{sgn}(r)]\Big(\frac{3}{2\sqrt{2}} - 1\Big)\pi\,; \qquad r = r' = \pm\infty\,, \qquad \text{(B21)}\end{aligned}$$

$$\pi\,\bar{\Phi}_{c,\eta\nu}(r,r') = -\arctan\Big(\frac{r-r'}{\nu}\Big) + \int_{-r^0_c}^{+r^0_c} dr''\,A(r-r'')\,\pi\,\bar{\Phi}_{c,\eta\nu}(r'',r')\,, \qquad \text{(B22)}$$

$$\pi\,\bar{\Phi}_{c,s\nu}(r,r') = 0\,; \qquad \nu > 1\,, \qquad \text{(B23)}$$

$$\pi\,\bar{\Phi}_{s1,\eta\nu}(r,r') = \frac{1}{4}\int_{-r^0_c}^{+r^0_c} dr''\,\frac{\pi\,\bar{\Phi}_{c,\eta\nu}(r'',r')}{\cosh\Big(\frac{\pi}{2}(r-r'')\Big)}\,, \qquad \text{(B24)}$$

$$\begin{aligned}\pi\,\bar{\Phi}_{s1,s\nu}(r,r') &= \arctan\Big(\frac{r-r'}{\nu-1}\Big)\,; \qquad r\neq\pm\infty\,; \qquad \nu>1\,,\\ &= \pm\frac{\pi}{\sqrt{2}}\,; \qquad r=\pm\infty\,; \qquad \nu>1\,,\end{aligned}\tag{B25}$$

$$\pi\,\bar{\Phi}_{s\nu,c}\,(r,r') = \pi\,\bar{\Phi}_{s\nu,\eta\nu'}\,(r,r') = 0\,; \qquad \nu>1\,,\tag{B26}$$

$$\pi\,\bar{\Phi}_{s\nu,s1}\,(r,r') = \arctan\Big(\frac{r-r'}{\nu-1}\Big)\,; \qquad \nu>1\,,\tag{B27}$$

$$\pi\,\bar{\Phi}_{s\nu,s\nu'}\,(r,r') = \frac{1}{2}\,\Theta_{\nu,\nu'}(r-r') - \arctan\Big(\frac{r-r'}{\nu+\nu'-2}\Big) - \arctan\Big(\frac{r-r'}{\nu+\nu'}\Big)\,, \qquad \nu,\nu'>1\,.\tag{B28}$$

The two-pseudofermion phase shifts $\pi\,\bar{\Phi}_{\eta\nu,c}\,(r,r')$, $\pi\,\bar{\Phi}_{\eta\nu,\eta\nu'}\,(r,r')$, and $\pi\,\bar{\Phi}_{\eta\nu,s\nu'}\,(r,r')$ remain being given by Eqs. (B10)-(B12).

In the above expressions the function $\Theta_{\nu,\nu'}(x)$ is defined in Eq. (B16),

$$B(r) = \int_0^\infty d\omega\frac{\sin(\omega\,r)}{\omega(1+e^{2\omega})} = \frac{i}{2}\ln\frac{\Gamma\Big(\frac{1}{2}+i\frac{r}{4}\Big)\,\Gamma\Big(1-i\frac{r}{4}\Big)}{\Gamma\Big(\frac{1}{2}-i\frac{r}{4}\Big)\,\Gamma\Big(1+i\frac{r}{4}\Big)}\,,\tag{B29}$$

and

$$A(r) = \frac{1}{\pi}\frac{dB(r)}{dr} = \frac{1}{\pi}\int_0^\infty d\omega\frac{\cos(\omega\,r)}{1+e^{2\omega}}\,,\tag{B30}$$

where $\Gamma(x)$ is the usual Γ function.

We note that the four expressions provided in Eq. (B18)-(B21) with $s1=s$, $r=x$, and $r_c^0=x_0$ are equivalent to expressions given in Eqs. (A9)-(A12) of Ref. [48]. (For the phase shifts of Eqs. (B20) and (B21) this equality refers to values of r such that $r\neq\infty$.) The rapidity phase-shift expressions defined here for $m\rightarrow 0$ correspond to some of the two-pseudofermion phase shifts plotted in units of π in Figs. 1-6.

The phase-shift expressions given in Eqs. (B23) and (B25)-(B28) apply to excited states generated by transitions from a $m=0$ and $n=1$ ground state. Closed-form analytical expressions for the remaining phase shifts of such excited states are then readily obtained by considering $r_c^0=0$ in the integral equations provided in Eqs. (B19)-(B22) and (B24). This leads to,

$$\pi\,\bar{\Phi}_{c,c}(r,r') = -B(r-r') = -\frac{i}{2}\ln\frac{\Gamma\Big(\frac{1}{2}+i\frac{(r-r')}{4}\Big)\,\Gamma\Big(1-i\frac{(r-r')}{4}\Big)}{\Gamma\Big(\frac{1}{2}-i\frac{(r-r')}{4}\Big)\,\Gamma\Big(1+i\frac{(r-r')}{4}\Big)}\,,\tag{B31}$$

$$\pi\,\bar{\Phi}_{c,s1}(r,r') = -\frac{1}{2}\arctan\Big(\sinh\Big(\frac{\pi}{2}(r-r')\Big)\Big) = \frac{i}{2}\ln\left(-i\,\frac{1+ie^{\frac{\pi}{2}(r-r')}}{1-ie^{\frac{\pi}{2}(r-r')}}\right)\,,\tag{B32}$$

$$\begin{aligned}\pi\,\bar{\Phi}_{s1,c}(r,r') &= -\frac{1}{2}\arctan\Big(\sinh\Big(\frac{\pi}{2}(r-r')\Big)\Big) = \frac{i}{2}\ln\left(-i\,\frac{1+ie^{\frac{\pi}{2}(r-r')}}{1-ie^{\frac{\pi}{2}(r-r')}}\right)\,; \qquad r\neq\pm\infty\\ &= -\frac{\mathrm{sgn}(r)\pi}{2\sqrt{2}}\,; \qquad r=\pm\infty\,,\end{aligned}\tag{B33}$$

$$\begin{aligned}\pi\,\bar{\Phi}_{s1,s1}(r,r') &= B(r-r') = \frac{i}{2}\ln\frac{\Gamma\Big(\frac{1}{2}+i\frac{(r-r')}{4}\Big)\,\Gamma\Big(1-i\frac{(r-r')}{4}\Big)}{\Gamma\Big(\frac{1}{2}-i\frac{(r-r')}{4}\Big)\,\Gamma\Big(1+i\frac{(r-r')}{4}\Big)}\,; \quad r\neq\pm\infty\\ &= \frac{\mathrm{sgn}(r)\pi}{2\sqrt{2}}\,; \qquad r=\pm\infty\,, \qquad r'\neq r\\ &= [\mathrm{sgn}(r)]\Big(\frac{3}{2\sqrt{2}}-1\Big)\pi\,; \qquad r=r'=\pm\infty\,,\end{aligned}\tag{B34}$$

$$\pi\,\bar{\Phi}_{c,\eta\nu}(r,r') = -\mathrm{arc\,tan}\Big(\frac{r-r'}{\nu}\Big)\,, \qquad \pi\,\bar{\Phi}_{s1,\eta\nu}(r,r') = \pi\,\bar{\Phi}_{c,s\nu'}(r,r') = 0\,, \qquad \nu' > 1\,, \tag{B35}$$

$$\begin{aligned}\pi\,\bar{\Phi}_{s1,s\nu}(r,r') &= \mathrm{arc\,tan}\Big(\frac{r-r'}{\nu-1}\Big)\,; \quad r \neq \pm\infty\,; \quad \nu > 1\,, \\ &= \pm\frac{\pi}{\sqrt{2}}\,; \quad r = \pm\infty\,; \quad \nu > 1\,,\end{aligned} \tag{B36}$$

$$\pi\,\bar{\Phi}_{\eta\nu,c}\,(r,r') = \mathrm{arctan}\Big(\frac{r-r'}{\nu}\Big)\,; \qquad \pi\,\bar{\Phi}_{\eta\nu,\eta\nu'}\,(r,r') = \frac{1}{2}\,\Theta_{\nu,\nu'}(r-r')\,; \qquad \pi\,\bar{\Phi}_{\eta\nu,s\nu'}\,(r,r') = 0\,, \qquad \nu > 0\,. \tag{B37}$$

Some of such rapidity two-pseudofermion phase-shift expressions are used in the derivation of the equalities of Eq. (55).

The two-pseudofermion phase shifts $\pi\,\Phi_{c,c}(q,q')$, $\pi\,\Phi_{c,s1}(q\,q')$, $\pi\,\Phi_{s1,c}(q,q')$, $\pi\,\Phi_{s1,s1}(q,q')$, $\pi\,\Phi_{c,\eta 1}(q,q')$, and $\pi\,\Phi_{s1,\eta 1}(q,q')$ plotted in Figs. 1-6 in units of π for $n = 0.59$, $m = 0$, and several U/t values are defined in Eq. (14). The $u \rightarrow 0$ phase shifts plotted in such figures have analytical expressions that we provide in the following. The form of the general two-pseudofermion phase-shift expression given in Eq. (14) reveals that the evaluation of such $u \rightarrow 0$ analytical expressions involves as well the derivation of the ground-state rapidity functions $\Lambda^0_{\beta}(q)$. The latter functions are defined in terms of their inverse functions in Eq. (A14). For a $m = 0$ initial ground state with an electronic density in the range defined by the inequality $0 \leq n \leq 1$ we find from use of the latter equation the following closed-form expressions for the ground-state functions $k^0_c(q)$, $\Lambda^0_c(q)$, $\Lambda^0_{\eta\nu}(q)$, and $\Lambda^0_{s1}(q)$, which refer to the $u \rightarrow 0$ limit,

$$\begin{aligned}k^0_c(q) &= \frac{q}{2}\,; \quad |q| \leq 2k_F\,, \\ &= \mathrm{sgn}(q)\,[|q| - k_F]\,; \quad 2k_F \leq |q| < \pi\,, \\ &= \mathrm{sgn}(q)\,\pi\,; \quad |q| = \pi\,,\end{aligned} \tag{B38}$$

$$\begin{aligned}\Lambda^0_c(q) &= \sin\Big(\frac{q}{2}\Big)\,; \quad |q| \leq 2k_F\,, \\ &= \mathrm{sgn}(q)\,\sin\Big((|q| - k_F)\Big)\,; \quad 2k_F \leq |q| < \pi\,, \\ &= 0\,; \quad |q| = \pi\,,\end{aligned} \tag{B39}$$

$$\begin{aligned}\Lambda^0_{\eta\nu}(q) &= \mathrm{sgn}(q)\,\sin\Big(\frac{(|q| + \pi n)}{2}\Big)\,; \quad 0 < |q| < (\pi - 2k_F)\,, \\ &= 0\,; \quad q = 0\,, \\ &= \pm\infty\,; \quad q = \pm(\pi - 2k_F)\,,\end{aligned} \tag{B40}$$

and

$$\begin{aligned}\Lambda^0_{s1}(q) &= \sin(q)\,; \quad |q| < k_F\,, \\ &= \pm\infty\,; \quad q = \pm k_F\,,\end{aligned} \tag{B41}$$

respectively.

Next, we use Eqs. (B38)-(B41) in the integral equations, Eqs. (B1)-(B15), which define the two-pseudofermion phase shifts. By manipulation of these equations, we find the expressions for the above bare-momentum two-pseudofermion phase shifts given in the following for the $u \rightarrow 0$ limit. (For the phase shifts $\pi\,\Phi_{s1,\alpha\nu}(q,q')$ we provide the values for $u = 0$ and $u \rightarrow 0$, when different.) The expressions read,

$$\pi\,\Phi_{c,c}(q,q') = -\mathrm{sgn}\Big(\sin k^0_c(q) - \sin k^0_c(q')\Big)\,\frac{\pi}{C_c(q)} + \delta_{|q|,2k_F}\,\delta_{q,q'}\,[\mathrm{sgn}(q)]\Big(\frac{3}{2\sqrt{2}} - 1\Big)\pi\,, \tag{B42}$$

$$\pi\,\Phi_{c,s1}(q,q') = -\mathrm{sgn}\Big(\sin k^0_c(q) - c_{s1}(q')\,\sin(q')\Big)\,\frac{\pi}{C_c(q)}\,, \tag{B43}$$

$$\begin{aligned}\pi\,\Phi_{s1,c}(q,q') &= -\text{sgn}\Big(\sin(q)-\sin k_c^0(q')\Big)\frac{\pi}{2}\,; \quad q\neq\pm k_F\\ &= -\frac{\text{sgn}(q)\pi}{2\sqrt{2}}\,; \quad q=\pm k_F\,; \quad u\rightarrow 0\\ &= 0\,; \quad\quad q=\pm k_F\,; \quad u=0\,,\end{aligned} \tag{B44}$$

$$\begin{aligned}\pi\,\Phi_{s1,s1}(q,q') &= 0\,; \quad\quad q\neq\pm k_F\\ &= \frac{\text{sgn}(q)\pi}{2\sqrt{2}}\Big[1+\delta_{q,q'}2(1-\sqrt{2})\Big]\,; \quad\quad q=\pm k_F\,; \quad\quad u\rightarrow 0\\ &= 0\,; \quad q=\pm k_F\,; \quad u=0\,,\end{aligned} \tag{B45}$$

$$\pi\,\Phi_{c,\eta1}(q,q') = -\text{sgn}\Big(\sin k_c^0(q)-c_{\eta1}(q')\sin k_{\eta1}^0(q')\Big)\frac{2\pi}{C_c(q)}\,, \tag{B46}$$

and

$$\pi\,\Phi_{s1,\eta1}(q,q') = -\theta(k_F-|q|)\,\text{sgn}\Big(\sin q-\sin k_{\eta1}^0(q')\Big)\frac{\pi}{2}\,. \tag{B47}$$

Here the sign function is such that $\text{sgn}(0)=0$ and the $\theta(x)$ function reads $\theta(x)=1$ for $x>0$ and $\theta(x)=0$ for $x\leq 0$, alike that of Eq. (29). Hence, $\pi\,\Phi_{s1,\eta1}(\pm k_F,q')=0$. Moreover, in the above equations, $k_c^0(q)=\lim_{u\rightarrow 0}k_c^0(q)$ where the $u\rightarrow 0$ value of $k_c^0(q)$ is given in Eq. (B38),

$$\begin{aligned}k_{\eta1}^0(q) &= \frac{q}{2}+\text{sgn}(q)\,k_F\,; \quad 0<|q|\leq[\pi-2k_F]\\ &= 0\,; \quad q=0\,,\end{aligned} \tag{B48}$$

$$C_c(q) = 2\Big[\theta(2k_F-|q|)+\sqrt{2}\,\delta_{|q|,2k_F}+2\,\theta(\pi-|q|)\theta(|q|-2k_F)+\delta_{|q|,\pi}\Big]\,, \tag{B49}$$

and

$$\begin{aligned}c_{s1}(q) &= 1\,, \quad |q|<k_F\,; \quad c_{s1}(q)=\infty\,, \quad q=\pm k_F\\ c_{\eta1}(q) &= 1\,, \quad |q|<[\pi-2k_F]\,; \quad c_{\eta1}(q)=\infty\,, \quad q=\pm[\pi-2k_F]\,.\end{aligned} \tag{B50}$$

Appendix C: Relation to the dressed phase shifts of other choices of elementary objects

To our knowledge, the first dressed phase shifts of elementary objects derived for the 1D Hubbard model are those of the c and s pseudoparticles of Refs. [47, 48]. They are defined by the integral equations (20)-(23) of Ref. [47] and (32)-(38) of Ref. [48]. Within the notations of this paper, such phase shifts refer to excited states with number values $N_\beta=0$ for the $\beta\neq c,s1$ branches and $M_{\alpha,-1/2}^{un}=0$ for $\alpha=\eta,s$, whose initial ground states have densities in the ranges $n\in[0,1[$ and $m\in]0,n]$. The c and s pseudoparticles of such references correspond to the c and $s1$ pseudofermions, respectively. The phase shifts first introduced in Refs. [47, 48] are a particular case of those defined by the general integral equations (B1)-(B15) of Appendix B.

The next dressed phase shifts introduced for the 1D Hubbard model are those associated with the holon and spinon scattering theory of Ref. [40]. Shortly afterwards, an alternative holon and spinon scattering theory was introduced in Refs. [41, 42]. While the phase shifts of Refs. [47, 48] refer to a particular case of those studied in this paper, the main goal of this appendix is to provide further information on the relation between the pseudofermion scattering theory and the traditional holon and spinon scattering theories of Refs. [40] and [41, 42], respectively. The traditional spin-1/2 spinons of the descriptions of Refs. [40] and [41, 42] are similar. On the other hand, the holons and antiholons of Ref. [40] are different from the also named holons and antiholons of Ref. [41, 42]. Most traditional holon and spinon descriptions used in the literature follow either the holon and antiholon definitions of Ref. [40] or Refs. [41, 42].

We start by comparing the occupancy configurations of the scattering theories elementary objects that generate the $[N_c^h+N_{s1}^h]=2$ reduced-subspace excited states where the dressed S matrix of Refs. [41, 42] is defined. It is a subspace of the $S_\eta=0;S_s=0;2S_c=N_a$ absolute ground state PS. We then show that for the excited states of the $S_\eta=0;S_s=0;2S_c=N_a$ absolute ground state to which the scattering theory of Refs. [41, 42] applies, the

	N_c^h	N_{s1}^h	δN_c	δN_{s1}	$N_{\eta 1}$	N_{s2}	$M^{un}_{\eta,-1/2}$	$M^{un}_{\eta,+1/2}$	$M^{un}_{s,-1/2}$	$M^{un}_{s,+1/2}$	$M_{\eta,-1/2}$	$M_{\eta,+1/2}$	$\delta M_{s,-1/2}$	$\delta M_{s,+1/2}$
η-spin triplet	2	0	-2	-1	0	0	0	2	0	0	0	2	-1	-1
η-spin triplet	2	0	-2	-1	0	0	1	1	0	0	1	1	-1	-1
η-spin triplet	2	0	-2	-1	0	0	2	0	0	0	2	0	-1	-1
η-spin singlet	2	0	-2	-1	1	0	0	0	0	0	1	1	-1	-1
spin triplet	0	2	0	-1	0	0	0	0	0	2	0	0	-1	1
spin triplet	0	2	0	-1	0	0	0	0	1	1	0	0	0	0
spin triplet	0	2	0	-1	0	0	0	0	2	0	0	0	1	-1
spin singlet	0	2	0	-2	0	1	0	0	0	0	0	0	0	0
doublet	1	1	-1	-1	0	0	1	0	1	0	1	0	0	-1
doublet	1	1	-1	-1	0	0	0	1	1	0	0	1	0	-1
doublet	1	1	-1	-1	0	0	1	0	0	1	1	0	-1	0
doublet	1	1	-1	-1	0	0	0	1	0	1	0	1	-1	0

TABLE II: The number and number deviation values of the present formulation elementary objects for the twelve types of $[N^h_{s1} + N^h_c] = 2$ energy eigenstates that span the reduced subspace.

phase shifts provided in Eqs. (5.19)-(5.21) of that reference correspond to particular cases of the overall phase shift functionals $Q_c(q)$ and $Q_{s1}(q)$ defined by Eq. (15). Next we consider the form of such phase shift functionals for some one- and two-electron excited states of ground states with densities in the ranges $n \in [0, 1]$ and $m \in [0, n]$. Their relation to the phase shifts of the holons and spinons of Ref. [40] is discussed in the case of one- and two-electron excited states of ground states with electronic density in the range $n \in [0, 1]$ and zero spin densisty, $m = 0$.

Finally, the technical problems that prevent the extension of the traditional spinon and holon scattering theory of Ref. [40] (and Refs. [41, 42]) to PSs spanned by excited states of initial ground states with spin density $m \neq 0$ (and densities $n \neq 1$ and $m \neq 0$) are discussed.

1. The reduced subspace of the initial $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state PS

The dressed S matrix of Refs. [41, 42] is defined in a reduced subspace of the PS spanned by the excited states of the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state. That PS is that common to the three scattering theories under consideration. The reduced subspace under consideration is spanned by that ground state and its excited states with numbers of holes in the c and $s1$ bands such that $[N^h_c + N^h_{s1}] = 2$. For the traditional descriptions of Refs. [40] and [41–43] this is equivalent to $[N^h_h + N_s] = 2$ and $[L_h + L_s] = 2$, respectively.

Here we describe the transitions from the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state to such excited states within the present operator formulation and the traditional holon and spinon descriptions of Refs. [40] and [41–43]. In such a reduced subspace the spinons of Refs. [40] and [41–43] are exactly the same. Moreover, the N^h_h holons of Ref. [40] correspond to the N^h_c c pseudofermion holes created under the transition to the excited states. Therefore, for simplicity our analysis of the above transitions refers to the present operator formulation and the traditional holon and spinon description of Refs. [41–43].

The number and number deviation values of the present formulation elementary objects for the twelve types of $[N^h_c + N^h_{s1}] = 2$ excited energy eigenstates that span the reduced subspace are provided in Table II. The corresponding number values of the elementary objects of the traditional holon and spinon descriptions of both Ref. [40] and Refs. [41, 42] are given in Table III. Moreover, in Table IV the values of $\sigma =\uparrow, \downarrow$ electron-number deviations, scatter-less phase shifts Q^0_c and Q^0_{s1} of Eq. (16), and η-spin and spin and corresponding projections are also provided for such excited states. For the initial $S_\eta = 0; S_s = 0; 2S_c = N_a$ and $[N^h_c + N^h_{s1}] = [L_h + L_s] = 0$ absolute ground state both the c and $s1$ momentum bands are full. The two hole momenta of the twelve types of $[N^h_c + N^h_{s1}] = [L_h + L_s] = 2$ excited energy eigenstates whose number and number deviation values are given in Tables II-IV emerge in the c and/or $s1$ band under the ground-state - excited-state transitions. In the $L_h = 2; L_s = 0$ excited states the two hole momenta q_j and $q_{j'}$ belong to the c band. In the $L_h = 0; L_s = 2$ states the two hole momenta q_j and $q_{j'}$ belong to the $s1$ band. In the $L_h = 1; L_s = 1$ states the hole momenta q_j and $q_{j'}$ belong to the c and $s1$ band, respectively. At fixed values of the two hole momenta such twelve types of $[L_h + L_s] = 2$ energy eigenstates are directly related to the two $SU(2)$ symmetry algebras in the $SO(4) = [SU(2) \otimes SU(2)]/Z_2$ symmetry of the 1D Hubbard model global $[SO(4) \otimes U(1)]/Z_2 = SO(3) \otimes SO(3) \otimes U(1)$ symmetry.

Specifically, at fixed values of the two c band hole momenta q_j and $q_{j'}$ the four types of $L_h = 2; L_s = 0$ energy eigenstates refer to the three η-spin-triplet excited energy eigenstates and the η-spin-singlet excited energy eigenstate. Within the present formulation, two η-spin-1/2 unbound η-spinons emerge under the ground-state transition to the three η-spin-triplet excited energy eigenstates. The three states are generated by η-spin flipping the two unbound

Excited states	N_h^p	N_h^h	$L_{h,-1/2}$	$L_{h,+1/2}$	$N_{s,-1/2} = L_{s,-1/2}$	$N_{s,+1/2} = L_{s,+1/2}$
η-spin triplet	0	2	0	2	0	0
η-spin triplet	0	2	1	1	0	0
η-spin triplet	0	2	2	0	0	0
η-spin singlet	0	2	1	1	0	0
spin triplet	0	0	0	0	0	2
spin triplet	0	0	0	0	1	1
spin triplet	0	0	0	0	2	0
spin singlet	0	0	0	0	1	1
doublet	0	1	1	0	1	0
doublet	0	1	0	1	1	0
doublet	0	1	1	0	0	1
doublet	0	1	0	1	0	1

TABLE III: The number values of traditional antiholons N_h^p, holons N_h^h, and $\pm 1/2$ spinons $N_{s,\pm 1/2}$ of Ref. [40] and traditional $\pm 1/2$ holons $L_{h,\pm 1/2}$ and $\pm 1/2$ spinons $L_{s,\pm 1/2}$ of Refs. [41, 42] for the same excited states as Table II. Note that $L_h = [L_{h,+1/2} + L_{h,-1/2}]$, $N_s = [N_{s,+1/2} + N_{s,-1/2}]$, and $L_s = [L_{s,+1/2} + L_{s,-1/2}]$.

η-spinons. The same applies to the two η-spin 1/2 holons emerging within the description of Refs. [41–43]. For the present formulation, under the transition from the ground state to the η-spin-singlet excited energy eigenstate two anti-bound η-spinons emerge within the η-spin-singlet configuration of one two-η-spinon composite $\eta 1$ pseudofermion. Within the description of Refs. [41–43], two holons emerge in a η-spin-singlet configuration. Although the studies of such references have not explicitly considered the c hidden $U(1)$ symmetry in $[SO(4)\otimes U(1)]/Z_2 = SO(3)\otimes SO(3)\otimes U(1)$, it was shown in the first paper that the BA solution used in the studies of these author does. Moreover, it is shown in that paper that the c band momentum occupancy configurations associated with changing the two hole momenta q_j and $q_{j'}$ generate representations of that hidden $U(1)$ symmetry algebra. On the other hand, the above four types of energy eigenstates corresponding to fixed values of the two hole momenta q_j and $q_{j'}$ refer to representations of the η-spin $SU(2)$ symmetry algebra. The two c pseudofermion holes emerging in the c band at the momenta q_j and $q_{j'}$ follow from the annihilation of two ground-state c pseudofermions.

On the other hand, at fixed values of the two $s1$ band hole momenta q_j and $q_{j'}$ the four types of $L_h = 0; L_s = 2$ energy eigenstates refer to the three spin-triplet and one spin-singlet excited energy eigenstates. Such four types of energy eigenstates correspond to spin $SU(2)$ symmetry algebra representations. Within the present formulation, a ground-state spin-neutral spinon pair of one composite $s1$ pseudofermion is broken under the transition from the absolute ground state to any of the three spin-triplet excited states. This gives rise to the two spin-1/2 unbound spinons of such states. The three spin-triplet states are generated within the present formulation by spin flipping the two emerging spin-1/2 unbound spinons. In addition, two spin-neutral $s1$ pseudofermion holes emerge in the $s1$ band at the above momenta q_j and $q_{j'}$. One of such holes follows from the annihilation of the $s1$ pseudofermion and the other results from the emergence of an extra discrete momentum value in the exotic $s1$ band. Within the description of Refs. [41–43], spin flipping the two emerging spin-1/2 traditional spinons also generates the three spin-triplet excited states. For the present formulation, under the transition from the ground state to the spin-singlet excited energy eigenstate four bound spin-1/2 spinons from two ground-state spin-singlet two-spinon composite $s1$ pseudofermions reorganize into a single spin-singlet four-spinon composite $s2$ pseudofermion. In the case of that excitation the two spin-neutral $s1$ pseudofermion holes emerging in the $s1$ band at the above momenta q_j and $q_{j'}$ follow from the annihilation of the two $s1$ pseudofermions. Within the description of Refs. [41–43] two traditional spinons emerge in a spin-singlet configuration. It is shown in the first paper that the $s1$ band momentum occupancy configurations associated with changing the two hole momenta q_j and $q_{j'}$ generate extra spin-singlet $SU(2)$ symmetry algebra representations.

At fixed values of both the c band hole momentum q_j and $s1$ band hole momentum $q_{j'}$, the remaining four types of $L_h = 1; L_s = 1$ energy eigenstates refer to both the η-spin and spin degrees of freedom. Indeed, they correspond to the choices $\sigma_\eta = +1/2; \sigma_s = +1/2$, $\sigma_\eta = -1/2; \sigma_s = +1/2$, $\sigma_\eta = +1/2; \sigma_s = -1/2$, and $\sigma_\eta = -1/2; \sigma_s = -1/2$ where σ_η and σ_s denote here the η-spin projection and spin projection of the created traditional holon and spinon, respectively. Such η-spin projections and spin projections are exactly the same as those of the unbound η-spinon and unbound spinon, respectively, created under the ground-state - excited state transition within the present formulation. Moreover, the c band momentum occupancy configurations associated with changing the hole momentum q_j generate representations of the c hidden $U(1)$ symmetry algebra. The $s1$ band momentum occupancy configurations associated with changing the hole momentum $q_{j'}$ generate extra spin $SU(2)$ symmetry algebra representations. The present four types of $L_h = 1; L_s = 1$ energy eigenstates are one-electron states. For further information on the c and $s1$ band processes involved within the present operator formulation in the ground-state transitions to such excited states, see Section V of the first paper.

Excited state	$\delta N_{\uparrow}$	$\delta N_{\downarrow}$	Q_c^0	Q_{s1}^0	S_η	S_η^{x3}	S_s	S_s^{x3}
η-spin triplet	-1	-1	$\pm\pi$	$\pm\pi$	1	1	0	0
η-spin triplet	0	0	$\pm\pi$	$\pm\pi$	1	0	0	0
η-spin triplet	1	1	$\pm\pi$	$\pm\pi$	1	-1	0	0
η-spin singlet	0	0	0	$\pm\pi$	0	0	0	0
spin triplet	1	-1	$\pm\pi$	$\pm\pi$	0	0	1	1
spin triplet	0	0	$\pm\pi$	$\pm\pi$	0	0	1	0
spin triplet	-1	1	$\pm\pi$	$\pm\pi$	0	0	1	-1
spin singlet	0	0	$\pm\pi$	0	0	0	0	0
doublet	0	1	$\pm\pi$	0	1/2	1/2	1/2	1/2
doublet	-1	0	$\pm\pi$	0	1/2	-1/2	1/2	1/2
doublet	1	0	$\pm\pi$	0	1/2	1/2	1/2	-1/2
doublet	0	-1	$\pm\pi$	0	1/2	-1/2	1/2	-1/2

TABLE IV: The values of the spin-projection $\sigma =\uparrow,\downarrow$ electronic number deviations δN_σ, scatter-less phase shifts Q_c^0 and Q_{s1}^0 of Eq. (16), and S_η, S_η^{x3}, S_s, and S_s^{x3} for each class of $[N_{s1}^h + N_c^h] = 2$ excited states of Tables II and III.

2. Relation to the half-filled reduced-subspace holon and spinon dressed phase shifts as defined in Ref. [42]

The phase shifts considered in Ref. [42] are associated with transitions from the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state to the twelve types of $[N_{s1}^h + N_c^h] = 2$ excited states. In addition, they refer only to the two scatterers created under the transitions to the excited states. According to the results of Section IV B, for such excited states the $\beta = \alpha\nu \neq s1$ pseudofermions are invariant under both the electron - rotated-electron and fermion - pseudofermion unitary transformations. As a result, such objects are not scatterers. Hence for such excited states the only scatterers of the pseudofermion scattering theory are the c pseudofermions and corresponding holes and $s1$ pseudofermions and corresponding holes.

The phase shifts generated by the transitions to the twelve types of $[N_{s1}^h + N_c^h] = 2$ excitated states considered in Section 2 of Ref. [42] were obtained from the BA solution and are shown here to correspond to a particular limit of those studied in this paper. For the asymptote coordinate choice of Eq. (44), $Q_\beta(q)$ is the overall β pseudofermion (or β pseudofermion-hole) phase shift. From analysis of Eqs. (11) and (15), within that choice $2\pi\,\Phi_{\beta,\beta'}(q_j, q_{j'})$ is an elementary two-pseudofermion phase shift. The studies of this paper consider other asymptote coordinates, usually used in standard quantum non-relativistic scattering theory, such that $x \in [-L/2, +L/2]$ and thus $\delta_\beta(q) = Q_\beta(q)/2$ is the overall β pseudofermion or β pseudofermion-hole phase shift and $\pi\,\Phi_{\beta,\beta'}(q_j, q_{j'})$ is an elementary two-pseudofermion phase shift. However, the uniquely defined quantity is the corresponding dressed S matrix, as defined in this paper.

The $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state occupancies are such that the c and $s1$ bands are full. Thus the $\beta = c, s1$ Fermi-momentum values given in Eq. (A13) read $q_{Fc} = q_c = \pi$ and $q_{Fs1} = q_{s1} = \pi/2$. We start by providing the bare-momentum distribution-function deviations of the corresponding twelve classes of excited states whose number and number deviation values are given in Tables II-IV.

The three classes of η-spin-triplet excited energy eigenstates have the same bare-momentum distribution function deviations given by,

$$\begin{aligned}
\delta N_c(q) &= -\frac{2\pi}{N_a}\sum_{h=1}^{2}\delta(q-q_h) - \frac{\pi}{N_a}\,\delta(q+\pi) + \frac{\pi}{N_a}\,\delta(q-\pi)\,, \quad |q_h| \leq \pi\,, \\
\delta N_{s1}(q) &= -\frac{\pi}{N_a}\,\delta(q+\pi/2) - \frac{\pi}{N_a}\,\delta(q-\pi/2)\,.
\end{aligned} \tag{C1}$$

The deviation $\delta N_{s1}(q)$ given here also applies to the η-spin-singlet excited states. On the other hand, for the c and $\eta 1$ branches the deviations read as follows for such states,

$$\delta N_c(q) = -\frac{2\pi}{N_a}\sum_{h=1}^{2}\delta(q-q_h)\,; \qquad \delta N_{\eta 1}(q) = \frac{2\pi}{N_a}\delta(q)\,. \tag{C2}$$

The three classes of spin-triplet excitations have again the same bare-momentum distribution function deviations given by,

$$\begin{aligned} \delta N_c(q) &= -\frac{\pi}{N_a}\,\delta(q+\pi) + \frac{\pi}{N_a}\,\delta(q-\pi)\,, \\ \delta N_{s1}(q) &= -\frac{2\pi}{N_a}\sum_{h=1}^{2}\delta(q-{q'}_h) + \frac{\pi}{N_a}\,\delta(q+\pi/2) + \frac{\pi}{N_a}\,\delta(q-\pi/2)\,, \quad |{q'}_h| \leq \pi/2\,. \end{aligned} \tag{C3}$$

The deviation $\delta N_c(q)$ given here also applies to the spin-singlet excited states. For the $s1$ and $s2$ branches the deviations of such states read,

$$\delta N_{s1}(q) = -\frac{2\pi}{N_a}\sum_{h=1}^{2}\delta(q-{q'}_h)\,, \quad |q_h| \leq \pi/2\,; \qquad \delta N_{s2}(q) = \frac{2\pi}{N_a}\delta(q)\,. \tag{C4}$$

The four classes of doublet excited states have the same bare-momentum distribution function deviations given by,

$$\begin{aligned} \delta N_c(q) &= -\frac{2\pi}{N_a}\delta(q-q_1) - \frac{\pi}{N_a}\,\delta(q+\pi) + \frac{\pi}{N_a}\,\delta(q-\pi)\,, \quad |q_1| \leq \pi\,, \\ \delta N_{s1}(q) &= -\frac{2\pi}{N_a}\delta(q-{q'}_1)\,, \quad |{q'}_1| \leq \pi/2\,. \end{aligned} \tag{C5}$$

Although concerning the comparison with the phase shifts of Ref. [42] for the η-spin-triplet and η-spin-singlet excited states (and spin-triplet and spin-singlet excited states) only the phase-shift functional $Q_c(q)$ (and $Q_{s1}(q)$) is needed, in the following we provide the specific forms of both $Q_c(q)$ and $Q_{s1}(q)$ for all excited states.

The use of Eqs. (C1)-(C5) in Eqs. (11) and (15) for the overall phase shift leads to,

$$Q_c(q) = \pi - \sum_{h=1}^{2} 2\pi\,\Phi_{c,c}(q,q_h) + \pi\,\Phi_{c,c}(q,\pi) - \pi\,\Phi_{c,c}(q,-\pi) - \pi\,\Phi_{c,s1}(q,\pi/2) - \pi\,\Phi_{c,s1}(q,-\pi/2)\,, \tag{C6}$$

and

$$Q_{s1}(q) = \pi - \sum_{h=1}^{2} 2\pi\,\Phi_{s1,c}(q,q_h) + \pi\,\Phi_{s1,c}(q,\pi) - \pi\,\Phi_{s1,c}(q,-\pi) - \pi\,\Phi_{s1,s1}(q,\pi/2) - \pi\,\Phi_{s1,s1}(q,-\pi/2)\,, \tag{C7}$$

for the three classes of η-spin-triplet states,

$$Q_c(q) = -\sum_{h=1}^{2} 2\pi\,\Phi_{c,c}(q,q_h) - \pi\,\Phi_{c,s1}(q,\pi/2) - \pi\,\Phi_{c,s1}(q,-\pi/2) + 2\pi\,\Phi_{c,\eta 1}(q,0)\,, \tag{C8}$$

and

$$Q_{s1}(q) = \pi - \sum_{h=1}^{2} 2\pi\,\Phi_{s1,c}(q,q_h) - \pi\,\Phi_{s1,s1}(q,\pi/2) - \pi\,\Phi_{s1,s1}(q,-\pi/2) + 2\pi\,\Phi_{s1,\eta 1}(q,0)\,, \tag{C9}$$

for the η-spin-singlet states,

$$Q_c(q) = \pi - \sum_{h=1}^{2} 2\pi\,\Phi_{c,s1}(q,{q'}_h) + \pi\,\Phi_{c,c}(q,\pi) - \pi\,\Phi_{c,c}(q,-\pi) + \pi\,\Phi_{c,s1}(q,\pi/2) + \pi\,\Phi_{c,s1}(q,-\pi/2)\,, \tag{C10}$$

and

$$Q_{s1}(q) = \pi - \sum_{h=1}^{2} 2\pi\,\Phi_{s1,s1}(q,{q'}_h) + \pi\,\Phi_{s1,c}(q,\pi) - \pi\,\Phi_{s1,c}(q,-\pi) + \pi\,\Phi_{s1,s1}(q,\pi/2) + \pi\,\Phi_{s1,s1}(q,-\pi/2)\,, \tag{C11}$$

for the three classes of spin-triplet states,

$$Q_c(q) = \pi - \sum_{h=1}^{2} 2\pi\,\Phi_{c,s1}(q,{q'}_h) + \pi\,\Phi_{c,c}(q,\pi) - \pi\,\Phi_{c,c}(q,-\pi) + 2\pi\,\Phi_{c,s2}(q,0)\,, \tag{C12}$$

and

$$Q_{s1}(q) = -\sum_{h=1}^{2} 2\pi\,\Phi_{s1,s1}(q,{q'}_h) + \pi\,\Phi_{s1,c}(q,\pi) - \pi\,\Phi_{s1,c}(q,-\pi) + 2\pi\,\Phi_{s1,s2}(q,0)\,, \tag{C13}$$

for the spin-singlet states and,

$$\begin{aligned} Q_c(q) &= \pi - 2\pi\,\Phi_{c,c}(q,q_1) - 2\pi\,\Phi_{c,s1}(q,{q'}_1) + \pi\,\Phi_{c,c}(q,\pi) - \pi\,\Phi_{c,c}(q,-\pi)\,, \\ Q_{s1}(q) &= -2\pi\,\Phi_{s1,c}(q,q_1) - 2\pi\,\Phi_{s1,s1}(q,{q'}_1) + \pi\,\Phi_{s1,c}(q,\pi) - \pi\,\Phi_{s1,c}(q,-\pi)\,, \end{aligned} \tag{C14}$$

for the four classes of doublet states.

We recall that the expressions of the phase-shift functionals $Q_\beta(q_j)$ where $\beta = c, s1$ given above for the reduced-subspace excited states refer to all corresponding $N_{a_\beta} = [N_\beta + N_\beta^h]$ pseudofermion and pseudofermion-hole scatterers. Indeed, the present formulation accounts for the scatterers that both are created under the transition from the ground state to the excited state and pre-exited in the ground state.

On the other hand, the scattering theory of Ref. [42] is normal ordered relative to the initial $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state, so that only the elementary objects that emerge under such a transition are accounted for. For both theories the elementary objects that emerge under such transitions play the role of scattering centers. Thus the phase shifts of the scattering theory of Ref. [42] refer only to the specific β band momentum values q in the argument of the phase-shift functionals $Q_\beta(q)$ that refer to scattering centers. Obviously, in the phase shift $Q_\beta(q)$ of a given scatterer only the contribution from the interaction with the scattering center other than itself is finite.

For instance, using of the explicit expressions of the two-pseudofermion phase shifts in those given in Eqs. (C6)-(C14) for $Q_c(q)$ and $Q_{s1}(q)$ and considering the values of the latter overall phase shifts at the specific momentum values $q = q_1$ and $q = {q'}_1$, respectively, of the elementary objects created under the transitions to the excited states we find,

$$Q_c(q_1) = 2\pi B\Big(\frac{4t(\sin k_1 - \sin k_2)}{U}\Big) = \delta_{CT}\,, \tag{C15}$$

for the η-spin-triplet states,

$$Q_c(q_1) = -2\arctan\Big(\frac{2t(\sin k_1 - \sin k_2)}{U}\Big) + 2\pi B\Big(\frac{4t(\sin k_1 - \sin k_2)}{U}\Big) = \delta_{CS} - \pi\,, \tag{C16}$$

for the η-spin-singlet states,

$$Q_{s1}({q'}_1) = -2\pi B\Big(\frac{4t({\Lambda'}_1 - {\Lambda'}_2)}{U}\Big) = \delta_{ST} - \pi\,, \tag{C17}$$

for the spin-triplet states,

$$Q_{s1}({q'}_1) = 2\arctan\Big(\frac{2t({\Lambda'}_1 - {\Lambda'}_2)}{U}\Big) - 2\pi B\Big(\frac{4t({\Lambda'}_1 - {\Lambda'}_2)}{U}\Big) = \delta_{SS}\,, \tag{C18}$$

for the spin-singlet states and,

$$\begin{aligned} Q_c(q_1) &= \arctan\Big(\sinh\Big(\pi\Big[\frac{2t(\sin k_1 - {\Lambda'}_1)}{U}\Big]\Big)\Big) = \delta_{\eta S}\,, \\ Q_{s1}({q'}_1) &= \arctan\Big(\sinh\Big(\pi\Big[\frac{2t({\Lambda'}_1 - \sin k_1)}{U}\Big]\Big)\Big) = \delta_{S\eta}\,, \end{aligned} \tag{C19}$$

for the doublet states. Here $k_1 = k_c^0(q_1)$, $k_2 = k_c^0(q_2)$, $\Lambda_c^0(q) = \sin k_c^0(q)$, ${\Lambda'}_1 = \Lambda_{s1}^0({q'}_1)$, ${\Lambda'}_2 = \Lambda_{s1}^0({q'}_2)$, the rapidity functions $k_c^0(q)$ and $\Lambda_{s1}^0(q)$ are defined in terms of their inverse functions in Eq. (A16) of Appendix A, and the function $B(r)$ is provided in Eq. (B29) of Appendix B.

To derive the $m \to 0$ and $n \to 1$ phase-shift expressions given in Eqs. (C15)-(C19) we used Eqs. (B31)-(B34) of Appendix B. For the phase shift of the η-spin-singlet (and spin-singlet) states we also used the solution $\Lambda_{\eta 1}(0) = [\Lambda_c^0(q_1) + \Lambda_c^0(q_2)]/2$ (and $\Lambda_{s2}(0) = [\Lambda_{s1}^0({q'}_1) + \Lambda_{s1}^0({q'}_2)]/2$) for the first (and second) equality of Eq. (55) with $\nu = 1$ (and $\nu' = 2$) in the first (and second) phase shift of Eq. (57).

By inspection of the above phase-shift expressions one indeed confirms that $Q_c(q_1)$ as provided in Eqs. (C15), (C16), and (C19) equals δ_{CT}, $\delta_{CS} - \pi$, and $\delta_{\eta S}$, respectively, with the corresponding phase shifts given in Ref. [42]. The phase shifts δ_{CT} and δ_{CS} are provided in Eq. (5.19) and $\delta_{\eta S}$ in Eq. (5.21) of that reference. Moreover, the phase shift

$Q_{s1}(q'_1)$ given in Eqs. (C17), (C18), and (C19) equals $\delta_{ST}-\pi$, δ_{SS}, and $\delta_{S\eta}$, respectively. In this case the phase shifts δ_{ST} and δ_{SS} are given in Eq. (5.20) and $\delta_{S\eta}$ in Eq. (5.21) of Ref. [42]. (For the phase shifts of the doublet excited states the confirmation of the above equalities involves the use of the relation $\arctan(\sinh(\pi x)) = [2\arctan(\exp(\pi x)) - \pi/2]$ for the branch for which these functions vary between $-\pi/2$ and $+\pi/2$.)

Note that the above phase shifts δ_{CS} and δ_{ST} read $Q_c(q_1)+\pi$ and $Q_{s1}(q'_1)+\pi$ whereas, according to the phase-shift definition of Eq. (44), the corresponding c pseudofermion-hole phase shifts are given by $Q_c(q_1)$ and $Q_{s1}(q'_1)$, respectively. The studies of Ref. [42] used the method of Ref. [53] to evaluate the above phase shifts. That method provides the phase shifts up to an overall constant term. In contrast, our method provides the full corresponding pseudofermion-hole phase shift value.

Hence we have just confirmed that in the reduced subspace spanned by the twelve types of $[N^h_{s1}+N^h_c]=2$ excited states of the $S_\eta = 0; S_s = 0; 2S_c = N_a$ and $[N^h_{s1}+N^h_c]=0$ absolute ground state the dressed phase shifts extracted in Ref. [42] from the BA solution, which in that reference were identified with the holon scatterer and spinon scatterer as defined in it, equal the c pseudofermion-hole and $s1$ pseudofermion-hole phase shifts, respectively. (Except for π in the case of some phase shifts.) Furthermore, we have shown that the phase shifts of Refs. [40, 42] are particular cases of the c and $s1$ pseudofermion overall phase-shift functionals $Q_\beta(q)$ defined by Eq. (15).

3. Phase shifts of simple one- and two-electron excited states of ground states with arbitrary densities

Here we provide the values of the phase shift functionals $Q_c(q)$ and $Q_{s1}(q)$ of Eq. (15) for some classes of one- and two-electron excited states of an initial ground state with densities in the ranges $n\in[0,1]$ and $m\in[0,n]$. Our choice of state classes is such that for $m\rightarrow 0$ the phase shifts considered in Ref. [40] are particular cases of such c or $s1$ phase shift functionals. (The phase shifts of Ref. [40] refer to electronic densities $n\in[0,1]$ and zero spin density.)

Within the rotated-electron operational description introduced in the first paper, the lower-Hubbard-band (LHB) and upper-Hubbard-band (UHB) are well-defined concepts. Specifically, the LHB is generated by transitions from a $n<1$ (and $n>1$) ground state to excited energy eigenstates with no rotated-electron doubly occupied sites, so that $M_{\eta,-1/2}=0$ (and no rotated-electron unoccupied sites, so that $M_{\eta,+1/2}=0$). For an initial $n=1$ ground state such transitions do not exist. The UHB is generated by transitions from $n\leq 1$ (and $n\geq 1$) ground states to excited energy eigenstates with $M_{\eta,-1/2}>0$ rotated-electron doubly occupied sites (and $M_{\eta,+1/2}>0$ rotated-electron unoccupied sites). For one-electron (and two-electron) excited states of $n\leq 1$ ground states, the spectral weight generated by transitions to excited energy eigenstates with two (and three) rotated-electron doubly occupied sites is very small. The same applies to the spectral weight generated by transitions from $n\geq 1$ ground states to one-electron (and two-electron) excited energy eigenstates with two (and three) rotated-electron unoccupied sites.

Moreover, for finite spin densities $m>0$ (and $m<0$) one considers the upper-spin-band (USB). It is generated by transitions from $m>0$ (and $m<0$) ground states to excited energy eigenstates with $[M_{s,-1/2}-N_{s1}]>0$ (and with $[M_{s,+1/2}-N_{s1}]>0$). Indeed, for finite spin density the energy spectrum of these excited states is gapped. For instance, for $m>0$ the minimum magnitude of such a gap is ω_0, Eq. (6), for $D_r=0$ and S_r given in Eq. (A19) of Appendix A.

Our analysis involves excited states of ground states with densities in the ranges $n\in[0,1]$ and $m\in[0,n]$. We call LHB states the excited states for which $M_{\eta,-1/2}=0$ and $[M_{s,-1/2}-N_{s1}]=0$, UHB states those for which $M_{\eta,-1/2}=1,2$ and $[M_{s,-1/2}-N_{s1}]=0$, and USB states those for which $M_{\eta,-1/2}=0$ and $[M_{s,-1/2}-N_{s1}]=1,2$. We recall that $M_{\eta,-1/2}=1$ corresponds to either $N_{\eta 1}=1$ and $M^{un}_{\eta,-1/2}=0$ or $N_{\eta 1}=0$ and $M^{un}_{\eta,-1/2}=1$. Furthermore, $[M_{s,-1/2}-N_{s1}]=1$ implies that either $N_{s2}=1$ and $M^{un}_{s,-1/2}=0$ or $N_{s2}=0$ and $M^{un}_{s,-1/2}=1$. The $M_{\eta,-1/2}=2$ (and $[M_{s,-1/2}-N_{s1}]=2$) excited states considered below have $M^{un}_{\eta,-1/2}=2$ (and $[M^{un}_{s,-1/2}-N_{s1}]=2$). The use of the PDT reveals that for initial ground states with densities in the ranges $n\in[0,1]$ and $m\in[0,n]$ only such classes of states contribute significantly to the one- and two-electron spectral weights.

Specifically, the deviation and number values relative to the numbers of such initial ground states of the corresponding twenty six classes (a)-(z) of excited states that are behind nearly all one- and two-electron spectral weights are given in Table V. As indicated in that table legend, the states classifications LHB, USB, and UHB refer to electronic densities and spin densities obeying the inequalities $n<1$ and $m>0$, respectively. On the other hand, in the spin-density $m\rightarrow 0$ limit the otherwise USB gapped excited states (h), (i), (j), (q), (u), (w), and (x) become for electronic densities $n<1$ LHB gapless states. Moreover, relative to the zero-energy level of an initial $(\mu+\mu^0)>0$ and $n=1$ ground state, the LHB excited states (b), (s), and (v) become gapped UHB states and the USB excited states (t), (u), and (x) become UHB + USB states. Finally, relative to the zero-energy level of an initial $\mu=\mu^0$ and $n=1$ ground state the gapped UHB excited states (d), (k), (m), and (r) become LHB gapless states. The gapless charge (and spin) gapless excited states (a) (and (f)) are generated by a single particle - hole process in the c (and $s1$) momentum band. They do not exist at $n=1$ (and $m=0$) because the c (and $s1$) momentum band is full for the

	Excited state	$\delta N_{\uparrow}$	$\delta N_{\downarrow}$	δS_{η}	$\delta S_{\eta}^{x_3}$	δS_s	$\delta S_s^{x_3}$	δN_c^h	δN_{s1}^h	δN_c	δN_{s1}	$N_{\eta 1}$	N_{s2}	$M_{\eta,-1/2}^{un}$	$\delta M_{\eta,+1/2}^{un}$	$M_{s,-1/2}^{un}$	$\delta M_{s,+1/2}^{un}$
(a)	c-$c(h)$-LHB (1)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
(b)	η-spinon-LHB	1	1	1	1	0	0	2	0	-2	-1	0	0	0	2	0	0
(c)	η-spinon-UHB (*)	0	0	1	0	0	0	2	0	-2	-1	0	0	1	1	0	0
(d)	η-spinon-UHB (*)	-1	-1	1	-1	0	0	2	0	-2	-1	0	0	2	0	0	0
(e)	$\eta 1$-UHB	0	0	0	0	0	0	2	0	-2	-1	1	0	0	0	0	0
(f)	$s1$-$s1(h)$-LHB (0)	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0
(g)	spinon-LHB	-1	1	0	0	1	1	0	2	0	-1	0	0	0	0	0	2
(h)	spinon-USB (*)	0	0	0	0	1	0	0	2	0	-1	0	0	0	0	1	1
(i)	spinon-USB (*)	1	-1	0	0	1	-1	0	2	0	-1	0	0	0	0	2	0
(j)	$s2$-USB	0	0	0	0	0	0	0	2	0	-2	0	1	0	0	0	0
(k)	1-elec-LHB (1)	1	0	-1/2	1/2	1/2	-1/2	-1	1	1	0	0	0	0	-1	0	1
(l)	1-elec-UHB (1)	1	0	-1/2	1/2	1/2	-1/2	1	1	-1	-1	1	0	0	-1	0	1
(m)	1-elec-UHB (*)	1	0	1/2	1/2	1/2	-1/2	1	1	-1	-1	0	0	1	0	0	1
(n)	1-elec-LHB (0,1)	0	1	-1/2	1/2	-1/2	1/2	-1	-1	1	1	0	0	0	-1	0	-1
(o)	1-elec-USB (0,1)	0	1	-1/2	1/2	-1/2	1/2	-1	1	1	-1	0	1	0	-1	0	-1
(p)	1-elec-UHB (0,1)	0	1	-1/2	1/2	-1/2	1/2	1	-1	-1	0	1	0	0	-1	0	-1
(q)	1-elec-USB (*,1)	0	1	-1/2	1/2	1/2	1/2	-1	1	1	0	0	0	0	-1	1	0
(r)	1-elec-UHB (*,0)	0	1	1/2	1/2	-1/2	1/2	1	-1	-1	0	0	0	1	0	0	-1
(s)	1-elec-LHB (0)	-1	0	1/2	-1/2	-1/2	1/2	1	-1	-1	0	0	0	0	1	0	-1
(t)	1-elec-USB (0)	-1	0	1/2	-1/2	-1/2	1/2	1	1	-1	-2	0	1	0	1	0	-1
(u)	1-elec-USB (*)	-1	0	1/2	-1/2	1/2	1/2	1	1	-1	-1	0	0	0	1	1	0
(v)	1-elec-LHB	0	-1	1/2	-1/2	1/2	-1/2	1	1	-1	-1	0	0	0	1	0	1
(w)	1-elec-USB (1)	1	0	-1/2	1/2	1/2	-1/2	-1	3	1	-2	0	1	0	-1	0	1
(x)	1-elec-USB	0	-1	1/2	-1/2	1/2	-1/2	1	3	-1	-3	0	1	0	1	0	1
(y)	1-elec-UHB (0)	-1	0	1/2	-1/2	-1/2	1/2	3	-1	-3	-1	1	0	0	1	0	-1
(z)	1-elec-UHB	0	-1	1/2	-1/2	1/2	-1/2	3	1	-3	-2	1	0	0	1	0	1

TABLE V: The number deviations and numbers of twenty six types (a)-(z) of one- and two-electron excited states of a ground state with densities in the ranges $n \in [0,1]$ and $m \in [0,n]$. Excited states marked with (1) have $\delta S_{\eta} < 0$ (and marked with (0) have $\delta S_s < 0$) and do not exist for an initial $n = 1$ (and $m = 0$) ground state. The excited states marked with (*) are outside the LWS BA solution subspace. The states classifications LHB, USB, and UHB refer to electronic densities and spin densities obeying the inequalities $n < 1$ and $m > 0$, respectively.

initial ground state.

Concerning the relative spectral weights generated by transitions to the excited states of Table V, we briefly report results for the one-electron weight for initial ground states with zero spin density. For instance, in the $m \to 0$ limit about 94% of the one-electron LHB addition spectral weight results for electronic densities $n < 1$ from transitions to the classes of excited states (k) and (n) [56]. They correspond to creation of one spin-up and one spin-down electron, respectively. In that limit such processes generate exactly the same weight. In the limit of zero spin density, most of the remaining one-electron LHB addition spectral weight stems from transitions to the excited states (w) and a corresponding excited-state branch generated by addition of one $s2$ pseudofermion to the excited states (q) [56].

In what the one-electron UHB addition is concerned there are in the $m \to 0$ limit two cases. When the initial ground state has electronic density $n = 1$ nearly the whole one-electron UHB addition spectral weight stems from transitions to the excited states (m) and (r) [56]. Such transitions involve creation of one spin-up and one spin-down electron, respectively. In turn, when the initial ground state has electronic density $n < 1$ nearly the whole one-electron UHB addition spectral weight stems from transitions to the excited state (l) and a corresponding excited-state branch generated by addition of one $\eta 1$ pseudofermion to the excited states (q) [56].

On the other hand, in the $m \to 0$ limit, about 97% of the one-electron removal spectral weight is generated by transitions to the classes of excited states (u) and (v) [56]. Such transitions lead to annihilation of one spin-up and one spin-down electron, respectively. Most of the remaining one-electron removal spectral weight stems from transitions to the excited states (x) and a corresponding excited-state branch generated by addition of one $s2$ pseudofermion to the excited states (u) [56].

For simplicity, we consider the specific form of the general phase-shift functional of Eq. (15) of the $\beta = c, s1$ branches for only eight of the twenty six excited-state classes of Table V: Those of the excited states (a), (e), (g)-(j), (k), and (v) of such a table. Our exited-state classes choice criterion is that in the $m \to 0$ limit some of their $\beta = c, s1$ phase shifts correspond to those studied in Ref. [40] for the electronic density range $n \in [0,1]$ and zero spin density, $m = 0$.

We start by providing the β band momentum distribution function deviations, Eq. (12), of such eight classes of excited states of ground states with density ranges $n \in [0,1]$ and $m \in [0,n]$. Those are needed for evaluating the expressions of the general $\beta = c, s1$ phase-shift functionals of Eq. (15) specific to such excited states. The corresponding $m \rightarrow 0$ expressions are obtained by replacing $k_{F\downarrow}$ by k_F in the phase-shift functionals expressions given in the following. For each class of excited states, we provide the β band momentum distribution function deviations of the β branches with finite occupancy in such states. All have finite occupancy in the c and $s1$ bands. Some of such states have one $s2$ pseudofermion or one $\eta 1$ pseudofermion.

For the class of gapless LHB charge excited states (a) such bare-momentum distribution function deviations are given by,

$$\begin{aligned}
\delta N_c(q) &= \frac{2\pi}{N_a}\,\delta(q-q_1) - \frac{2\pi}{N_a}\,\delta(q-q_1')\,, \\
q_1 &\in [-\pi, -2k_F] \text{ and } q_1 \in [2k_F, \pi]\,, \quad q_1' \in [-2k_F, 2k_F]\,, \\
\delta N_{s1}(q) &= 0\,.
\end{aligned} \tag{C20}$$

For the class of gapped charge excited states (e) they read,

$$\begin{aligned}
\delta N_c(q) &= -\frac{2\pi}{N_a}\sum_{h=1}^{2}\delta(q-q_h)\,, \quad q_h \in [-2k_F, 2k_F]\,, \\
\delta N_{s1}(q) &= -\frac{\pi}{N_a}\,\delta(q+k_{F\downarrow}) - \frac{\pi}{N_a}\,\delta(q-k_{F\downarrow})\,, \\
\delta N_{\eta 1}(q) &= \frac{2\pi}{N_a}\delta(q_1')\,, \quad q_1' \in [-(\pi-2k_F), (\pi-2k_F)]\,.
\end{aligned} \tag{C21}$$

The three classes of LHB spin excited states (g) and USB spin excited states (h) and (i) have the same momentum distribution function deviations, given by,

$$\begin{aligned}
\delta N_c(q) &= -\frac{\pi}{N_a}\,\delta(q+2k_F) + \frac{\pi}{N_a}\,\delta(q-2k_F)\,, \\
\delta N_{s1}(q) &= -\frac{2\pi}{N_a}\sum_{h=1}^{2}\delta(q-{q'}_h) + \frac{\pi}{N_a}\,\delta(q+k_{F\downarrow}) + \frac{\pi}{N_a}\,\delta(q-k_{F\downarrow})\,, \quad {q'}_h \in [-k_{F\downarrow}, k_{F\downarrow}]\,.
\end{aligned} \tag{C22}$$

The deviation $\delta N_c(q)$ provided here also applies to the USB spin excited states (j). For the $s1$ and $s2$ branches the momentum distribution function deviations of the latter excited states read,

$$\delta N_{s1}(q) = -\frac{2\pi}{N_a}\sum_{h=1}^{2}\delta(q-{q'}_h)\,, \quad {q'}_h \in [-k_{F\downarrow}, k_{F\downarrow}]\,; \quad \delta N_{s2}(q) = \frac{2\pi}{N_a}\delta(q_1'')\,, \quad q_1'' \in [-(k_{F\uparrow}-k_{F\downarrow}), (k_{F\uparrow}-k_{F\downarrow})]\,. \tag{C23}$$

For the LHB addition of one up-spin electron excited states (k) the momentum distribution function deviations are given by,

$$\begin{aligned}
\delta N_c(q) &= \frac{2\pi}{N_a}\,\delta(q-q_1)\,, \quad q_1 \in [-\pi, -2k_F] \text{ and } q_1 \in [2k_F, \pi]\,, \\
\delta N_{s1}(q) &= -\frac{2\pi}{N_a}\,\delta(q-{q'}_1) + \frac{\pi}{N_a}\,\delta(q+k_{F\downarrow}) + \frac{\pi}{N_a}\,\delta(q-k_{F\downarrow})\,, \quad {q'}_1 \in [-k_{F\downarrow}, +k_{F\downarrow}]\,,
\end{aligned} \tag{C24}$$

Finally, for the class of LHB excited states (v) involving the removal of one down-spin electron the momentum distribution function deviations read,

$$\begin{aligned}
\delta N_c(q) &= -\frac{2\pi}{N_a}\,\delta(q-q_1) - \frac{\pi}{L}\,\delta(q\mp 2k_F) + \frac{\pi}{N_a}\,\delta(q\pm 2k_F)\,, \quad q_1 \in [-2k_F, +2k_F] \\
\delta N_{s1}(q) &= -\frac{2\pi}{N_a}\,\delta(q-{q'}_1)\,, \quad {q'}_1 \in [-k_{F\downarrow}, +k_{F\downarrow}]\,.
\end{aligned} \tag{C25}$$

The classes of excited states (a), (e), (g)-(j), (k), and (v) have expressions for the general phase-shift functional of Eq. (15) for the $\beta = c, s1$ branches that are uniquely defined by the corresponding momentum distribution function deviations provided in Eqs. (C20)-(C25). For the class of LHB gapless charge excited states (a) with momentum distribution function deviations provided in Eq. (C20) that functional is for such two branches given by,

$$\begin{aligned} Q_c(q) &= 2\pi\,\Phi_{c,c}(q,q_1) - 2\pi\,\Phi_{c,c}(q,q_1')\,, \\ Q_{s1}(q) &= 2\pi\,\Phi_{s1,c}(q,q_1) - 2\pi\,\Phi_{s1,c}(q,q_1')\,. \end{aligned} \tag{C26}$$

For the class of UHB gapped charge excited states (e) whose momentum distribution function deviations are given in Eq. (C21), they read,

$$\begin{aligned} Q_c(q) &= -\sum_{h=1}^{2} 2\pi\,\Phi_{c,c}(q,q_h) - \pi\,\Phi_{c,s1}(q,k_{F\downarrow}) - \pi\,\Phi_{c,s1}(q,-k_{F\downarrow}) + 2\pi\,\Phi_{c,\eta1}(q,q_1')\,, \\ Q_{s1}(q) &= \pi - \sum_{h=1}^{2} 2\pi\,\Phi_{s1,c}(q,q_h) - \pi\,\Phi_{s1,s1}(q,k_{F\downarrow}) - \pi\,\Phi_{s1,s1}(q,-k_{F\downarrow}) + 2\pi\,\Phi_{s1,\eta1}(q,q_1')\,. \end{aligned} \tag{C27}$$

For the three classes of LHB spin excited states (g) and USB spin excited states (h) and (i) whose momentum distribution function deviations are provided in Eq. (C22) the expressions are,

$$\begin{aligned} Q_c(q) &= \pi - \sum_{h=1}^{2} 2\pi\,\Phi_{c,s1}(q,{q'}_h) + \pi\,\Phi_{c,c}(q,2k_F) - \pi\,\Phi_{c,c}(q,-2k_F) + \pi\,\Phi_{c,s1}(q,k_{F\downarrow}) + \pi\,\Phi_{c,s1}(q,-k_{F\downarrow})\,, \\ Q_{s1}(q) &= \pi - \sum_{h=1}^{2} 2\pi\,\Phi_{s1,s1}(q,{q'}_h) + \pi\,\Phi_{s1,c}(q,2k_F) - \pi\,\Phi_{s1,c}(q,-2k_F) + \pi\,\Phi_{c,s1}(q,k_{F\downarrow}) + \pi\,\Phi_{s1,s1}(q,-k_{F\downarrow}) \end{aligned} \tag{C28}$$

For the USB spin excited states (j) whose momentum distribution function deviations are given both in Eqs. (C22) and (C23) the phase-shift functional of Eq. (15) reads for the $\beta = c, s1$ branches,

$$\begin{aligned} Q_c(q) &= \pi - \sum_{h=1}^{2} 2\pi\,\Phi_{c,s1}(q,{q'}_h) + \pi\,\Phi_{c,c}(q,2k_F) - \pi\,\Phi_{c,c}(q,-2k_F) + 2\pi\,\Phi_{c,s2}(q,q_1'')\,, \\ Q_{s1}(q) &= -\sum_{h=1}^{2} 2\pi\,\Phi_{s1,s1}(q,{q'}_h) + \pi\,\Phi_{s1,c}(q,2k_F) - \pi\,\Phi_{s1,c}(q,-2k_F) + 2\pi\,\Phi_{s1,s2}(q,q_1'')\,. \end{aligned} \tag{C29}$$

For the addition of one up-spin electron LHB excited states (k) with momentum distribution function deviations provided in Eq. (C24) the result is,

$$\begin{aligned} Q_c(q) &= 2\pi\,\Phi_{c,c}(q,q_1) - 2\pi\,\Phi_{c,s1}(q,{q'}_1) + \pi\,\Phi_{c,s1}(q,-k_{F\downarrow}) + \pi\,\Phi_{c,s1}(q,+k_{F\downarrow})\,, \\ Q_{s1}(q) &= \pi + 2\pi\,\Phi_{s1,c}(q,q_1) - 2\pi\,\Phi_{s1,s1}(q,{q'}_1) + \pi\,\Phi_{s1,s1}(q,-k_{F\downarrow}) + \pi\,\Phi_{s1,s1}(q,+k_{F\downarrow})\,. \end{aligned} \tag{C30}$$

Finally, for the class of LHB excited states (v) involving the removal of one down-spin electron whose momentum distribution function deviations are given in Eq. (C25) the branches $\beta = c, s1$ the phase-shift functionals are given by,

$$\begin{aligned} Q_c(q) &= \pi - 2\pi\,\Phi_{c,c}(q,q_1) - \pi\,\Phi_{c,c}(q,\mp 2k_F) + \pi\,\Phi_{c,c}(q,\pm 2k_F) - 2\pi\,\Phi_{c,s1}(q,{q'}_1)\,, \\ Q_{s1}(q) &= -2\pi\,\Phi_{s1,c}(q,q_1) - \pi\,\Phi_{s1,c}(q,\mp 2k_F) + \pi\,\Phi_{s1,c}(q,\pm 2k_F) - 2\pi\,\Phi_{s1,s1}(q,{q'}_1)\,. \end{aligned} \tag{C31}$$

The expressions of the phase-shift functionals $Q_\beta(q)$ where $\beta = c, s1$ given above for the excited states (a), (e), (g)-(j), (k), and (v) of Table V also apply to the corresponding discrete momentum values, upon replacing the continuum momentum variable q by q_j. The β band has a set $\{q_j\}$ of discrete momentum values, $j = 1, ..., N_{a_\beta}$, such that $q_{j+1} - q_j = 2\pi/N_a$. Such phase shifts are those of the excited-state N_β β pseudofermion scatterers that carry N_β of the $N_{a_\beta} = [N_\beta + N^h_\beta]$ β band discrete momentum values q_j and N^h_β β pseudofermion-hole scatterers that carry the remaining N^h_β discrete momentum values q_j of the β band. Such phase shifts are due to the transition from the ground state to the excited state under consideration. Hence the phase shift $Q_\beta(q_j)$ refers to all excited-state $N_{a_\beta} = [N_\beta + N^h_\beta]$ scatterers. As discussed in Section VI C, that the present formulation accounts for the scatterers that both are created under the transition from the ground state to the excited state and pre-exit in the ground state is for the present metallic phase a necessary condition for its applicability to the study of the model dynamical and spectral properties.

On the other hand and alike for the scattering theory of Ref. [42], for that of Ref. [40] the few scatterers emerge under the transition from the ground state to the excited state. Indeed, such theories are normal-ordered relative to the initial ground state. For instance, the holon and antiholon phase shifts of Ref. [40] due to the interaction with the antiholon and the holon, respectively, are the phase shift $Q_c(q)$ at $q = q_1$ and $q = q_1'$, respectively, given in Eq. (C26) as $m \rightarrow 0$ for the class of gapless charge excited states (a),

$$Q_c(q_1) = -2\pi\,\Phi_{c,c}(q_1, q_1')\,; \qquad Q_c(q_1') = 2\pi\,\Phi_{c,c}(q_1', q_1)\,, \tag{C32}$$

where we have accounted for once more that both $2\pi\,\Phi_{c,c}(q_1, q_1) = 0$ and $2\pi\,\Phi_{c,c}(q_1', q_1') = 0$. Moreover, the holon - holon phase shift of Ref. [40] is the phase shift $Q_c(q)$ at $q = q_1$ provided in Eq. (C27) as $m \rightarrow 0$ for the class of gapped charge excited states (e),

$$Q_c(q_1) = -2\pi\,\Phi_{c,c}(q_1, q_2) - \pi\,\Phi_{c,s1}(q_1, k_F) - \pi\,\Phi_{c,s1}(q_1, -k_F) + 2\pi\,\Phi_{c,\eta 1}(q, q_1')\,, \tag{C33}$$

where again we used that $2\pi\,\Phi_{c,c}(q_1, q_1) = 0$. (A phase shift with exactly the same information is obtained from that given here by introducing the changes $q_1 \rightarrow q_2$ and $q_2 \rightarrow q_1$.) The phase shift for the spinons of Ref. [40] in the spin-triplet excited states is the phase shift $Q_{s1}(q)$ at $q = {q'}_1$ given in Eq. (C28) as $m \rightarrow 0$ for the three classes of spin excitations (g), (h), and (i),

$$Q_{s1}({q'}_1) = \pi - 2\pi\,\Phi_{s1,s1}({q'}_1, {q'}_2) + \pi\,\Phi_{s1,c}({q'}_1, 2k_F) - \pi\,\Phi_{s1,c}({q'}_1, -2k_F) + \pi\,\Phi_{s1,s1}({q'}_1, k_F) + \pi\,\Phi_{s1,s1}({q'}_1, -k_F)\,, \tag{C34}$$

where we have accounted for that $2\pi\,\Phi_{s1,s1}({q'}_1, {q'}_1) = 0$. The scattering dressed phase shift for the spinons of Ref. [40] in the spin-singlet excited states is the phase shift $Q_{s1}(q)$ at $q = {q'}_1$ given in Eq. (C29) as $m \rightarrow 0$ for the class of spin excitations (j),

$$Q_{s1}({q'}_1) = -2\pi\,\Phi_{s1,s1}({q'}_1, {q'}_2) + \pi\,\Phi_{s1,c}({q'}_1, 2k_F) - \pi\,\Phi_{s1,c}({q'}_1, -2k_F) + 2\pi\,\Phi_{s1,s2}({q'}_1, 0)\,, \tag{C35}$$

where again we used that $2\pi\,\Phi_{s1,s1}({q'}_1, {q'}_1) = 0$ and that for $m \rightarrow 0$ the $s2$ band momentum range $q_1'' \in [-(k_{F\uparrow} - k_{F\downarrow}), (k_{F\uparrow} - k_{F\downarrow})]$ of Eq. (C23) shrinks to the a single value, $q_1'' = 0$, for the excited states. (Again phase shifts with exactly the same information as those of Eqs. (C34) and (C35) are obtained from their expressions provided in such equations by introducing the changes ${q'}_1 \rightarrow {q'}_2$ and ${q'}_2 \rightarrow {q'}_1$.) The phase shifts of Ref. [40] for the antiholon and the spinon due to the interaction with the spinon and the antiholon, respectively, are the phase shifts $Q_c(q)$ at $q = q_1$ and $Q_{s1}(q)$ at $q = {q'}_1$, respectively, given in Eq. (C30) as $m \rightarrow 0$ for the class of excited states (k) involving the addition of one up-spin electron,

$$\begin{aligned} Q_c(q_1) &= -2\pi\,\Phi_{c,s1}(q_1, {q'}_1) + \pi\,\Phi_{c,s1}(q_1, -k_F) + \pi\,\Phi_{c,s1}(q_1, +k_F)\,, \\ Q_{s1}({q'}_1) &= \pi + 2\pi\,\Phi_{s1,c}({q'}_1, q_1) + \pi\,\Phi_{s1,s1}({q'}_1, -k_F) + \pi\,\Phi_{s1,s1}({q'}_1, +k_F)\,, \end{aligned} \tag{C36}$$

where we have accounted for that $2\pi\,\Phi_{c,c}(q_1, q_1) = 0$ and $2\pi\,\Phi_{s1,s1}({q'}_1, {q'}_1) = 0$. The phase shifts of Ref. [40] for the holon and the spinon due to the interaction with the spinon and the holon, respectively, are the phase shifts $Q_c(q)$ at $q = q_1$ and $Q_{s1}(q)$ at $q = {q'}_1$, respectively, given in Eq. (C31) as $m \rightarrow 0$ for the class of excited states (v) involving the removal of one down-spin electron,

$$\begin{aligned} Q_c(q_1) &= -\pi\,\Phi_{c,c}(q_1, \mp 2k_F) + \pi\,\Phi_{c,c}(q_1, \pm 2k_F) - 2\pi\,\Phi_{c,s1}(q_1, {q'}_1)\,, \\ Q_{s1}({q'}_1) &= -2\pi\,\Phi_{s1,c}({q'}_1, q_1) - \pi\,\Phi_{s1,c}({q'}_1, \mp 2k_F) + \pi\,\Phi_{s1,c}({q'}_1, \pm 2k_F)\,, \end{aligned} \tag{C37}$$

where we have accounted for that $2\pi\,\Phi_{c,c}(q_1, q_1) = 0$ and $2\pi\,\Phi_{s1,s1}({q'}_1, {q'}_1) = 0$.

In order to confirm that the c and $s1$ phase shifts given in Eqs. (C32)-(C37) indeed correspond to the phase shifts of Ref. [40] indicated above, we have used procedures similar to those presented in detail in Section C 2, concerning the relation of the two-pseudofermion phase shifts to the $n = 1$ and $m = 0$ phase shifts of Ref. [42]. Such procedures involve manipulations of the general relation given in Eq. (14) and integral equations for the two-pseudofermion phase shifts $\pi\,\bar{\Phi}_{\beta,\beta'}\,(r, r')$ on the right-hand side of that equation provided in Eqs. (B18)-(B30) of Appendix B for electronic density $n \in [0, 1]$ and spin density $m \rightarrow 0$. For the $\beta = c, s1$ and $\beta' = c, s1$ BA branches that includes the expression of the phase shifts $\pi\,\bar{\Phi}_{\beta,\beta'}(r, r') = \pi\,\bar{\Phi}_{\beta,\beta'}(\Lambda/u, \Lambda'/u)$ in terms of the variables Λ such that,

$$\begin{aligned} k &= \arcsin\Lambda = \arcsin(r\,u) \in [-\pi, \pi]\,, \quad \beta = c\,, \\ \Lambda &= r\,u \in [-\infty, \infty]\,, \quad \beta = s1\,. \end{aligned} \tag{C38}$$

Here k and Λ are the continuum variables associated with the ground-state set of c momentum-rapidity values $k_j = k_c^0(q_j)$ where $j = 1, ..., N_a$ and discrete $s1$ rapidity values $\Lambda_j = \Lambda_{s1}^0(q_j)$, where $j = 1, ..., N_\uparrow$ for $m \in (0, n)$

and $j = 1, ..., N$ for $m = 0$, respectively. Such sets of discrete values are uniquely defined by the rapidity functions $\Lambda^0_\beta(q_j)$ appearing in Eq. (14), where $\Lambda^0_c(q_j) = \sin k_j = \sin k^0_c(q_j)$ for $\beta = c$. They are defined in terms of their inverse functions in Eq. (A14) of Appendix A. For ground states with densities in the ranges $n \in (0, 1)$ and $m \in (0, n)$, N and $[N_a - N]$ discrete c momentum-rapidity values k_j are occupied and unoccupied and $N_\downarrow$ and $[N_\uparrow - N_\downarrow]$ discrete $s1$ rapidity values Λ_j are occupied and unoccupied, respectively. For a zero-spin-density ground state all N $s1$ rapidity values Λ_j are occupied.

In terms of the corresponding c rapidity-momentum continuum variable k one has for the ground state an occupied range $k \in (-Q, Q)$ and unoccupied ranges $k \in (-\pi, -Q)$ and $k \in (Q, \pi)$. For the $s1$ rapidity continuum variable Λ the ground-state occupied range is $\Lambda \in (-B, B)$ and unoccupied ranges are $\Lambda \in (-\infty, -B)$ and $\Lambda \in (B, \infty)$. Here Q and B are the parameters defined in Eq. (A15) of Appendix A, with $B \to \infty$ for the zero-spin-density limit considered in Ref. [40].

Appendix D: Role of the excited energy eigenstates degeneracy in the definition of the traditional spinon and holon descriptions subspaces

In this appendix we provide some further information on the the role of the degeneracy of the excited energy eigenstates in the definition of the subspaces where the traditional spinon and holon descriptions of Ref. [40] and Refs. [41, 42] are defined. For simplicity, we consider in general only the case of the traditional spinon and holon scattering theory of Refs. [41, 42]. The arguments concerning the spinon sector of the scattering theory of Ref. [40] are exactly the same.

For any scattering theory, both the one-particle states belonging to the "in" and "out" state are the "in" and "out" asymptote scattering states. Such asymptote scattering states must have a well-defined energy. In the context of the 1D Hubbard model scattering theories associated with the BA dressed phase shifts and dressed S matrices, such states must be energy eigenstates. If one extends the traditional holons and spinons as defined in Refs. [41–43] to PSs other than that of the initial $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state, some of the excited states associated with the corresponding one-particle scattering states become a superposition of energy eigenstates of different energy. Thus such excited states have not a well-defined energy, so that the corresponding scattering theory is not physically acceptable. The occurrence or lack of excited energy eigenstates degeneracy then plays the main role in the definition of the subspaces where it is possible to introduce multiple choices or a single choice, respectively, of scattering states and corresponding elementary objects that play the role of scatterers and scattering centers.

If alike in Refs. [41, 42] for the specific PS of the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state, for PSs of ground states with densities in the ranges $n \in [0, 1]$ and $m \in [0, n]$ one identifies the holes in the BA c and $s1$ momentum bands with η-spin-1/2 holons and spin-1/2 spinons, respectivey, those are not mere excitations of such ground states. Indeed, for such densities the ground state itself has a finite number of BA c and $s1$ band momentum holes, $N^h_c = [N_a - N]$ and $N^h_{s1} = [N_\uparrow - N_\downarrow]$, respectively. Hence the initial ground state itself would be populated by a finite number, $L^0_h = [N_a - N]$ and $L^0_s = [N_\uparrow - N_\downarrow]$, of traditional holons and spinons, respectively. The relevant parameters for the elementary objects description then are the corresponding number deviations, $\delta L_h = [L_h - L^0_h]$ and $\delta L_s = [L_s - L^0_s]$, which refer to the excited states of the initial ground state.

Let us assume that for the excited energy eigenstates that span the PSs of initial ground states with arbitrary densities n and m the BA c and $s1$ band momentum holes indeed refer to traditional η-spin-1/2 holons and traditional spin-1/2 spinons, respectively. For each PS, one can again consider a smaller reduced subspace contained in it, which is spanned by twelve types of $[\delta L_h + \delta L_s] = 2$ excited energy eigenstates. For the initial ground states under consideration there are many more classes of excited states, as for instance those whose number and number deviations are given in Table V of Appendix C. For simplicity, we limit our considerations to $[\delta L_h + \delta L_s] = 2$ excited energy eigenstates, which are those that in the $n \to 1; m \to 0$ limit evolve into the excited energy eigenstates that span the reduced-subspace of the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state. Within the states of Table V of Appendix C, the $[\delta L_h + \delta L_s] = 2$ excited energy eigenstates are the η-spinon and $\eta 1$ states (b)-(e), spinon and $s2$ states (g)-(j), and one-electron states (m) and (v). Consistent with the definition of the L_h and L_s numbers, for such excited states one has that $\delta L_h = \delta N^h_c$ and $\delta L_s = 2N_{s2}$ or $\delta L_s = [\delta M^{un}_{s,-1/2} + \delta M^{un}_{s,+1/2}]$ provided that both $\delta M^{un}_{s,-1/2} \geq 0$ and $\delta M^{un}_{s,+1/2} \geq 0$.

Alike in Section VI B for the excited states of the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state, we denote the corresponding two sets of four holon-holon states ($\alpha = \eta$) and four spinon-spinon states ($\alpha = s$) by $|+1/2, +1/2; \alpha\rangle$, $|-1/2, -1/2; \alpha\rangle$, $|+1/2, -1/2; \alpha\rangle$, and $|-1/2, +1/2; \alpha\rangle$. The two η-spin ($\alpha = \eta$) and spin ($\alpha = s$) projections $\pm 1/2$ labeling these states are those of the two traditional holons and spinons, respectively, created under the transition to the excited state. Furthermore, we denote the four types of excited η-spinon and $\eta 1$ energy eigenstates (b)-(e) (and the four types of excited spinon and $s2$ energy eigenstates (g)-(j)) of Table V of Appendix C, which are needed to express the four holon-holon states (and four spinon-spinon states), by $|c, c; +1\rangle$, $|c, c; 0\rangle$, $|c, c; -1\rangle$, and $|c, c, \eta 1; 0\rangle$ (and $|s1, s1; +1\rangle$, $|s1, s1; 0\rangle$, $|s1, s1; -1\rangle$, and $|s1, s1, s2; 0\rangle$.) The index with values $0, \pm 1$ of these states refers now to their value of the deviation $\delta S_\eta^{x_3}$ (and $\delta S_s^{x_3}$).

The main difference relative to the corresponding excited states of the $S_\eta = 0; S_s = 0; 2S_c = N_a$ absolute ground state stems from the states (e), $|c, c, \eta 1; 0\rangle$, and (j), $|s1, s1, s2; 0\rangle$, of Table V of Appendix C. For the corresponding excited energy eigenstates of the absolute ground state, one has that the $\eta 1$ and $s2$ momentum bands of the two-η-spinon composite $\eta 1$ pseudofermion and four-spinon composite $s2$ pseudofermion of such states, respectively, has a single discrete momentum value, which is $q = 0$. Furthermore, the corresponding energy $\varepsilon_{\eta 1}^0(0) = 0$ and and $\varepsilon_{s2}^0(0) = 0$ given in Eqs. (A20) and (A21) of Appendix A vanishes at $n = 1$ at $m = 0$, respectively. Hence the two energy eigenstates $|c, c; 0\rangle$ and $|c, c, \eta 1; 0\rangle$ (and $|s1, s1; 0\rangle$ and $|s1, s1, s2; 0\rangle$) with the same c (and $s1$) band hole momentum values q_j and $q_{j'}$ involved in the expressions of Eq. (93) have exactly the same energy. Specifically, their energy spectrum is of the form,

$$\begin{aligned}
\delta E_{c,c;0} &= \delta E_{c,c,\eta 1;0} = 2\mu^0 - \varepsilon_c(q) - \varepsilon_c(q') \, , \quad q, q' \in [-\pi, \pi] \, , \quad n = 1 \, , m = 0 \, , \\
\delta E_{s1,s1;0} &= \delta E_{s1,s1,\eta 1;0} = -\varepsilon_{s1}(q) - \varepsilon_{s1}(q') \, , \quad q, q' \in [-\pi/2, \pi/2] \, , \quad n = 1 \, , m = 0 \, ,
\end{aligned} \tag{D1}$$

where $2\mu^0$ denotes the Mott-Hubbard gap, which has the limiting behaviors given in Eq. (38), and the c and $s1$ energy dispersions are defined in Eqs. (A20) and (A21) of Appendix A. It follows that the corresponding holon-holon and spinon-spinon scattering states,

$$\begin{aligned}
| \pm 1/2, \mp 1/2; \eta\rangle &= \frac{1}{\sqrt{2}} \Big[|c, c; 0\rangle \mp |c, c, \eta 1; 0\rangle \Big] \, , \\
| \pm 1/2, \mp 1/2; s\rangle &= \frac{1}{\sqrt{2}} \Big[|s1, s1; 0\rangle \mp |s1, s1, s2; 0\rangle \Big] \, ,
\end{aligned} \tag{D2}$$

respectively, are energy eigenstates of the 1D Hubbard model whose excitation energies are those given in Eq. (D1).

On the other hand, besides the fixed values of the two c (and $s1$) band hole momenta q_j and $q_{j'}$, for initial ground states with electronic density $n \neq 1$ (and spin density $m \neq 0$) the excited states (e), $|c, c, \eta 1; 0\rangle$, (and (j), $|s1, s1, s2; 0\rangle$,) of Table V of Appendix C involve a third momentum $q_{j''}$: That of the $\eta 1$ pseudofermion (and $s2$ pseudofermion) created under the transition to the excited state. Indeed, for electronic densities in the range $n \in [0, 1]$ (and spin densities in the range $m \in [0, n]$) the ground-state $\eta 1$ (and $s2$) momentum band is empty, yet has a finite momentum range width, $q \in [-(\pi - 2k_F), (\pi - 2k_F)]$ (and $q \in [-(k_{F\uparrow} - k_{F\downarrow}), (k_{F\uparrow} - k_{F\downarrow})]$.) The energy spectra of the excited energy eigenstates $|c, c; 0\rangle$ and $|c, c, \eta 1; 0\rangle$ now read,

$$\begin{aligned}
\delta E_{c,c;0} &= 2|\mu| - \varepsilon_c(q) - \varepsilon_c(q') \, , \\
\delta E_{c,c,\eta 1;0} &= 2|\mu| - \varepsilon_c(q) - \varepsilon_c(q') + \varepsilon_{\eta 1}^0(q'') \, , \\
q, q' &\in [-2k_F, 2k_F] \, , \; q'' \in [-(\pi - 2k_F), (\pi - 2k_F)] \, .
\end{aligned} \tag{D3}$$

Corresponding similar expressions hold for the energy spectra of the excited energy eigenstates $|s1, s1; 0\rangle$ and $|s1, s1, s2; 0\rangle$,

$$\begin{aligned}
\delta E_{s1,s1;0} &= 2\mu_B |H| - \varepsilon_{s1}(q) - \varepsilon_{s1}(q') \, , \\
\delta E_{s1,s1,\eta 1;0} &= 2\mu_B |H| - \varepsilon_{s1}(q) - \varepsilon_{s1}(q') + \varepsilon_{s2}^0(q'') \, , \\
q, q' &\in [-k_{F\downarrow}, k_{F\downarrow}] \, , \; q'' \in [-(k_{F\uparrow} - k_{F\downarrow}), (k_{F\uparrow} - k_{F\downarrow})] \, .
\end{aligned} \tag{D4}$$

That for electronic densities $n < 1$ (and finite spin densities $m > 0$) the $\eta 1$ (and $s2$) pseudofermion energy is such that the excited energy eigenstates $|c, c; 0\rangle$ and $|c, c, \eta 1; 0\rangle$ (and $|s1, s1; 0\rangle$ and $|s1, s1, s2; 0\rangle$) in the expressions provided in Eq. (D2) for the holon-holon (and spinon-spinon) states have for $q_{j''} \neq \pm(\pi - 2k_F)$ (and $q_{j''} \neq \pm(k_{F\uparrow} - k_{F\downarrow})$) different energy implies that they are not energy eigenstates. Since they have not a well defined energy, such states cannot play the role of asymptotic scattering states.

The exception refers to the $\eta 1$ (and $s2$) band limiting momentum values $q_{j''} = \pm(\pi - 2k_F)$ (and $q_{j''} = \pm(k_{F\uparrow} - k_{F\downarrow})$) at which $\varepsilon_{\eta 1}^0(\pm(\pi - 2k_F)) = 0$ (and $\varepsilon_{s2}^0(\pm(k_{F\uparrow} - k_{F\downarrow})) = 0$.) For all other $q_{j''}$ values in the range $[-(\pi - 2k_F), (\pi - 2k_F)]$ of Eq. (D3) (and $[-(k_{F\uparrow} - k_{F\downarrow}), (k_{F\uparrow} - k_{F\downarrow})]$ of Eq. (D4)), the magnitude of the $\eta 1$ pseudofermion energy $\varepsilon_{\eta 1}(q_{j'''})$ (and $s2$ pseudofermion energy $\varepsilon_{s2}(q_{j''})$) appearing in the energy spectrum $\delta E_{c,c,\eta 1;0}$ given in Eq. (D3) (and energy spectrum $\delta E_{s1,s1,s2;0}$ given in Eq. (D4)) is such that the states $|c, c; 0\rangle$ and $|c, c, \eta 1; 0\rangle$ (and the states $|s1, s1; 0\rangle$ and $|s1, s1, s2; 0\rangle$) have different energies.

In the $n \to 1$ limit the energy of the holon-holon states given in Eq. (D2) becomes well defined. Indeed, in that limit the $\eta 1$ band momentum range $q_1'' \in [-(\pi - 2k_F), (\pi - 2k_F)]$ of Eq. (D3), which is that of the $\eta 1$ momentum distribution function deviation in Eq. (C21) of Appendix C for the class of gapped charge excited states (e) of Table V of that appendix, $|c, c, \eta 1; 0\rangle$, shrinks to $q_1'' = 0$. This restores the energy degeneracy of the states (c) and (e) of

Table V, $|c,c;0\rangle$ and $|c,c,\eta 1;0\rangle$, respectively. A similar behavior occurs in the $m\rightarrow 0$ limit concerning the energy of the spinon-spinon states of Eq. (D2), which becomes well defined. In that limit the $s2$ band momentum range $q''_1\in[-(k_{F\uparrow}-k_{F\downarrow}),(k_{F\uparrow}-k_{F\downarrow})]$ in Eq. (D4) and in the $s2$ momentum distribution function deviation of Eq. (C23) of Appendix C for the class of the USB spin excited states (j) of Table V of that appendix, $|s1,s1,s2;0\rangle$, also shrinks to $q''_1=0$. This restores the energy degeneracy of the states (h) and (j) of Table V, $|s1,s1;0\rangle$ and $|s1,s1,s2;0\rangle$, respectively. Specifically, in the $n\rightarrow 1;m\rightarrow 0$ limit the energy spectra provided in Eqs. (D3) and (D4) become those given in Eq. (D1).

Thus only when the excited energy eigenstates degeneracy is brought about by the $\eta 1$ (and $s2$) band momentum range of the excited states (e) (and (j)) of Table V of Appendix C shrinking to zero momentum, have the holon-holon states (and spinon-spinon states) given in Eq. (D2) a well-defined energy. Only in that case such holon-holon states (and spinon-spinon states) become asymptotic scattering states of the theory of Refs. [41, 42]. The same arguments apply to the spin-spin states of the scattering theory of Ref. [40]. The only difference is that such a theory is valid for excited states of ground states with zero spin density and electronic densities in the range $n\in[0,1]$, so that the $s1$ band momentum range $q,q'\in[-\pi/2,\pi/2]$ in Eq. (D1) for the momentum values of the two emerging holes changes to $q,q'\in[-k_F,k_F]$.

Similar degeneracies occur for $u>0$ in the $N_a\rightarrow\infty$ limit for excited energy eigenstates whose occupancies of the $\eta\nu$ bands (and $s\nu$ bands of $s\nu$ pseudofermions with $\nu>1$ bound spinon pairs) are such that $\sum_{\nu=1}^{\infty}N_{\eta\nu}/N_a\rightarrow 0$ (and $\sum_{\nu=2}^{\infty}N_{s\nu}/N_a\rightarrow 0$) as $N_a\rightarrow\infty$. The traditional holon and spinon scattering theories of Ref. [40] and Refs. [41, 42] are valid for such excited states provided that the corresponding initial ground state has densities $n=1;m=0$ and $n\in[0,1];m=0$, respectively.

Finally, we consider the $u\rightarrow\infty$ limit, in which the full degeneracy of both η-spin and spin configurations brings about a qualitatively different situation. Indeed, that degeneracy renders the holon and spinon scattering theories as defined in Ref. [40] and Refs. [41–43] well defined in the subspaces spanned by excited states of ground states with both arbitrary electronic density n and spin density m. We recall that for the pseudofermion scattering theory this holds for all $u>0$ values. For the other scattering theories this $u\rightarrow\infty$ behavior follows from the energy bandwidths of the energy dispersions $\varepsilon^0_{\alpha\nu}(q)$ given in Eqs. (A20) and (A21) of Appendix A vanishing for all $\alpha=\eta,s$ and $\nu=1,...,\infty$ branches as $u\rightarrow\infty$. Thus in that very special limit the pseudofermion scattering theory and the traditional holon and spinon scattering theories of Refs. [40–42] are alternative distinct theories for the excited states of ground states with arbitrary electronic density and spin density values. On the other hand, for finite u values only the pseudofermion scattering theory introduced in this paper is valid for such larger set of excited states.

---

[1] Elliott H. Lieb, F. Y. Wu, Phys. Rev. Lett. 20 (1968) 1445;
Elliott H. Lieb, F. Y. Wu, Physica A 321 (2003) 1.
[2] Minoru Takahashi, Progr. Theor. Phys 47 (1972) 69.
[3] F. Woynarovich, J. Phys. C 15 (1982) 85;
F. Woynarovich, J. Phys. C 15 (1982) 97.
[4] P. B. Ramos, M. J. Martins, J. Phys. A 30 (1997) L195;
M. J. Martins, P. B. Ramos, Nucl. Phys. B 522 (1998) 413.
[5] J. M. P. Carmelo, arXiv:1211.5391.
[6] J. Voit, Rep. Prog. Phys. 58 (1995) 977 and references therein.
[7] J. M. P. Carmelo, K. Penc, D. Bozi, Nucl. Phys. B 725 (2005) 421;
J. M. P. Carmelo, K. Penc, D. Bozi, Nucl. Phys. B 737 (2006) 351, Erratum;
J. M. P. Carmelo, K. Penc, Eur. Phys. J. B 51 (2006) 477.
[8] J. M. P. Carmelo, L. M. Martelo, K. Penc, Nucl. Phys. B 737 (2006) 237.
[9] J. M. P. Carmelo, D. Bozi, K. Penc, J. Phys.: Cond. Mat. 20 (2008) 415103;
D. Bozi, J. M. P. Carmelo, K. Penc, P. D. Sacramento, J. Phys.: Cond. Mat. 20 (2008) 022205.
[10] Karlo Penc, Karen Hallberg, Frédéric Mila, Hiroyuki Shiba, Phys. Rev. Lett. 77 (1996) 1390;
Karlo Penc, Karen Hallberg, Frédéric Mila, Hiroyuki Shiba, Phys. Rev. B 55 (1997) 15 475.
[11] A. Imambekov, L. I. Glazman, Science 323 (2009) 228;
A. Imambekov, L. I. Glazman, Phys. Rev. Lett. 100 (2008) 206805.
[12] A. Imambekov, L. I. Glazman, Phys. Rev. Lett. 102 (2009) 126405.
[13] Thomas L. Schmidt, Adilet Imambekov, Leonid I. Glazman, Phys. Rev. Lett. 116 (2010) 116403;
Thomas L. Schmidt, Adilet Imambekov, Leonid I. Glazman, Phys. Rev. B, 82 (2010) 245104;
A. Shashi, L. I. Glazman, J.-S. Caux, A. Imambekov, Phys. Rev. B, 84 (2011) 045408.
[14] Adilet Imambekov, Thomas L. Schmidt, Leonid I. Glazman, Rev. Mod. Phys. 84 (2012) 1253.
[15] R. G. Pereira, S. R. White, I. Affleck, Phys. Rev. Lett. 100 (2008) 027206; S. R. White, I. Affleck, Phys. Rev. B 77 (2008) 134437.
[16] R. G. Pereira, S. R. White, I. Affleck, Phys. Rev. B 79 (2009) 165113.

[17] Fabian H. L. Essler, Phys. Rev. B 81 (2010) 205120.
[18] R. G. Pereira, K. Penc, S. R. White, P. D. Sacramento J. M. P. Carmelo, Phys. Rev. B 85 (2012) 165132.
[19] B. Sriram Shastry, Phys. Rev. Lett. 56 (1986) 1529;
B. Sriram Shastry, Phys. Rev. Lett. 56 (1986) 2453.
[20] B. Sriram Shastry, J. Stat. Phys. 50 (1988) 57.
[21] J. M. P. Carmelo, T. Prosen, D. K. Campbell, Phys. Rev. B 63 (2001) 205114.
[22] M. Sing, U. Schwingenschlögl, R. Claessen, P. Blaha, J. M. P. Carmelo, L. M. Martelo, P. D. Sacramento, M. Dressel, C. S. Jacobsen, Phys. Rev. B 68 (2003) 125111.
[23] J. M. P. Carmelo, K. Penc, L. M. Martelo, P. D. Sacramento, J. M. B. Lopes dos Santos, R. Claessen, M. Sing, U. Schwingenschlögl, Europhys. Lett. 67 (2004) 233.
[24] J. M. P. Carmelo, K. Penc, P. D. Sacramento, M. Sing, R. Claessen, M. Sing, J. Phys.: Cond. Mat. 18 (2006) 5191.
[25] Dionys Baeriswyl, José Carmelo, Kazumi Maki, Synth. Met. 21 (1987) 271.
[26] D. Jaksch, P. Zoller, Ann. Phys. 315 (2005) 52.
[27] F. H. L. Essler, V. E. Korepin, K. Schoutens, Phys. Rev. Lett. 67 (1991) 3848.
[28] O. J. Heilmann, E. H. Lieb, Ann. N. Y. Acad. Sci. 172 (1971) 583.
[29] E. H. Lieb, Phys. Rev. Lett. 62 (1989) 1201.
[30] C. N. Yang, Phys. Rev. Lett. 63 (1989) 2144;
C. N. Yang, S. C. Zhang, Mod. Phys. Lett. B 4 (1990) 759;
S. C. Zhang, Phys. Rev. Lett. 65 (1990) 120.
[31] J. M. P. Carmelo, S. Östlund, M. J. Sampaio, Ann. Phys. 325 (2010) 1550.
[32] Stellan Östlund, Eugene Mele, Phys. Rev. B 44 (1991) 12413.
[33] J. M. P. Carmelo, J. M. Román, K. Penc, Nucl. Phys. B 683 (2004) 387.
[34] J. M. P. Carmelo, P. D. Sacramento, Phys. Rev. B 68 (2003) 085104.
[35] J. M. P. Carmelo, P. D. Sacramento, N. M. R. Peres, Phys. Rev. Lett. 84 (2000) 4673;
J. M. P. Carmelo, J. M. E. Guerra, J. M. B. Lopes dos Santos, A. H. Castro Neto, Phys. Rev. Lett. 83 (1999) 3892;
J. M. P. Carmelo, P. Horsch, A. A. Ovchinnikov, D. K. Campbell, A. H. Castro Neto, N. M. R. Peres, Phys. Rev. Lett. 81 (1998) 489;
J. M. P. Carmelo, A. H. Castro Neto, Phys. Rev. Lett. 70 (1993) 1904;
J. M. P. Carmelo, A. H. Castro Neto, D. K. Campbell, Phys. Rev. Lett. 73 (1994) 926 and Erratum 74 (1995) 3089;
J. M. P. Carmelo, P. Horsch, A. A. Ovchinnikov, Phys. Rev. Lett. 68 (1992) 871.
[36] J. M. P. Carmelo, A. H. Castro Neto, D. K. Campbell, Phys. Rev. B 50 (1994) 3667;
J. M. P. Carmelo, A. H. Castro Neto, D. K. Campbell, Phys. Rev. B 50 (1994) 3683;
J. M. P. Carmelo, P. Horsch, D. K. Campbell, A. H. Castro Neto, Phys. Rev. B (RC) 48 (1993) 4200.
[37] N. Andrei, J. Lowenstein, Phys. Rev. Lett. 43 (1979) 1698.
[38] N. Andrei, J. Lowenstein, Phys. Lett. 91B (1980) 401.
[39] L. D. Faddeev, L. A. Takhtajan, Phys. Lett. 85A (1981) 375;
L. D. Faddeev, L. A. Takhtadzhyan, J. Math. Sciences 24 (1984) 241.
[40] N. Andrei, *Integrable Models in Condensed Matter Physics, in Series on Modern Condensed Matter Physics - Vol. 6*, 458, World Scientific, Lecture Notes of ICTP Summer Course, Editors: S. Lundquist, G. Morandi, Yu Lu [cond-mat/9408101].
[41] F. H. L. Essler, V. E. Korepin, Phys. Rev. Lett. 72 (1994) 908.
[42] F. H. L. Essler, V. E. Korepin, Nucl. Phys. B 426 (1994) 505.
[43] F. H. L. Essler, H. Frahm, F. Göhmann, A. Klümper, V. E. Korepin, *The one-dimensional Hubbard model* (Cambridge University Press, Cambridge, UK, 2005).
[44] K.-V. Pham, M. Gabay, P. Lederer, Phys. Rev. B 61 (2000) 16 397.
[45] John R. Taylor, *Scattering theory: the quantum theory of nonrelativistic collisions* (Robert E. Krieger Publishing Company, Malabar, Florida, 1987).
[46] Gerald D. Mahan, *Many-particle physics* (Kluwer Academic/Plenum Publishers, New York, 2000), Chapter 4; F. G. Fumi, Philos. Mag. 46 (1955) 1007.
[47] J. Carmelo and A. A. Ovchinnikov, J. Phys.: Condens. Matter 3 (1991) 757.
[48] J. M. P. Carmelo, P. Horsch, A. A. Ovchinnikov, Phys. Rev. B 45 (1992) 7899.
[49] H. C. Ohanian, *Principles of quantum mechanics* (Prentice Hall, Englewood Cliffs, New Jersey, 1993).
[50] J. M. P. Carmelo, S.-J. Gu, P. D. Sacramento, arXiv:1209.1276.
[51] S.-J. Gu, N. M. R. Peres, J. M. P. Carmelo, J. Phys.: Condens. Matter 19 (2007) 506203.
[52] J. M. P. Carmelo, P. Horsch, A. A. Ovchinnikov, Phys. Rev. B 46 (1992) 14 728.
[53] V. E. Korepin, Teor. Mat. Fiz. 41 (1979) 169 (1979); V. E. Korepin, Commun. Math. Phys. 76 (1980) 165.
[54] Holger Frahm, V. E. Korepin, Phys. Rev. B 42 (1990) 10553.
[55] N. M. R. Peres, J. M. P. Carmelo, D. K. Campbell, A. W. Sandvik, Z. Phys. B 103 (1997) 217.
[56] J. M. P. Carmelo, K. Penc, J. Phys.: Cond. Mat. 18 (2006) 2881.